




ERRATA.

VOL., 1939.

Page. Line.  
1849 4 for "2:3-Dimethyl" read "2:4-Dimethyl."

VOL., 1942.

- 534 The second formula for *basseol* should be 
- 537 In the figure the numbers II and III should be interchanged.  
543 20 The oxidation of methyl *siarasinolate* by the *Oppenbauer reagent* (see p. 541) is as follows: Methyl *siarasinolate* (1 g.), aluminium *tert.*-butoxide (3.3 g.), cyclohexanone (10 c.c.), and toluene (50 c.c.) were boiled for 18 hours, the mixture diluted with water and extracted with ether, the extract dried and evaporated, and the residue heated to 100° in a high vacuum. The residue crystallised from dilute acetone in plates, m. p. 214—215°,  $[\alpha]_D^{20} + 57^\circ$  ( $c = 1.75$  in chloroform), giving a yellow colour with tetranitromethane (Found: C, 76.9; H, 10.1.  $C_{31}H_{48}O_4$  requires C, 76.8; H, 10.0%).
- 544 23 for " $C_{29}H_{44}$ " read " $C_{29}H_{48}$ ."  
545 16\* The *dehydro-ester* was prepared as follows: Methyl acetyldihydro-*a*-*elemolate* (700 mg.) was boiled for 16 hours with freshly sublimed selenium dioxide (350 mg.) in AnalaR acetic acid (42 c.c.). The solution was evaporated to dryness under reduced pressure, the residue dissolved in ether, and the solution washed with water, sodium hydroxide, again with water, dried, and evaporated. The residue was dissolved in benzene, and the solution percolated through a short column of alumina. The solid recovered on evaporation of the benzene solution crystallised from methyl alcohol in flattened needles (400 mg.), m. p. 128—129° (Found: C, 77.1, 77.4; H, 10.4, 10.2.  $C_{33}H_{52}O_4$  requires C, 77.3; H, 10.2%). As the intensity of the light absorption was less than had been expected, the compound was subjected to a lengthy repurification by chromatographic adsorption in petroleum solution, but was recovered unchanged and behaved like a pure compound.
- 550 41 for " $\beta$ -*Elemolic acid*" read " $\beta$ -*Elemonic acid*."  
724 23 for "50°" read "5°."  
725 9 for "*NN'*-dimethyldiacridine" read "*NN'*-dimethyldiacridene."  
19 for "*NN'*-Dimethyldiacridine" read "*NN'*-Dimethyldiacridene."  
5\* for "*NN'*-Dimethyldiacridine" read "*NN'*-Dimethyldiacridene."  
741 22 for "*Helv. Chim. Acta*, 1942, 25, 775" read "*Helv. Chim. Acta*, 1942, 25, 1236."
- in formula (III) ring D should be 
- 742 formula (VII) should be 
- 750 add H in formula (I).  
752 30\* for "*-d*-galactoside" read "*l*-galactoside."  
39\* for "←" read "→" in bottom row of formulae.

VOL., 1943.

52 34\* for "below" read "above."  
80 2 for "phosphoric oxide" read "phosphoric oxide (14 g.)."

\* From bottom.