

Chemical Communications

NUMBER 2/1967

25 JANUARY

Pentafluorophenylarsenic Derivatives

By M. GREEN and D. KIRKPATRICK

(Department of Inorganic Chemistry, The University, Bristol 8)

THERE has recently been considerable interest in pentafluorophenylphosphorus compounds, but little has been reported on the chemistry of the corresponding arsenic derivatives. We report some of our results concerning $(C_6F_5)_2As$ and C_6F_5As derivatives, which show interesting differences on comparison with the corresponding phosphorus derivatives.

The arsenic chlorides, $C_6F_5AsCl_2$ (b.p. $52^\circ/0.1$ mm.) and $(C_6F_5)_2AsCl$ (b.p. $82^\circ/0.1$ mm.), were made by reaction of C_6F_5MgBr with $AsCl_3$ or $Me_2N,AsCl_2$, respectively. Treatment of pentafluorophenylarsonous dichloride with mercury gave a benzene-soluble compound, which after recrystallisation from benzene-petroleum ether was characterised by elemental analyses and molecular weight measurements [Found: 952 ($CHCl_3$), 968 (mass-spectroscopy); Calc. for $(C_6F_5As)_4$: 968] as the tetramer $(C_6F_5As)_4$, m.p. 141° . The molecular formula agrees with the ^{19}F n.m.r. spectrum, which showed the expected pattern for a C_6F_5X derivative with peaks centred at 125.9, 152.1, and 161.8 p.p.m. of intensities 2:1:2. The *ortho*-fluorines at low field showed evidence of ^{75}As quadrupolar broadening.

The phosphorus compounds, $C_6F_5PCl_2$ and $C_6F_5PBr_2$, form the pentamer $(C_6F_5P)_5$ on reaction with mercury and it is interesting that the only previously reported² As_4 ring compound is $(CF_3As)_4$. The corresponding methylarsenic and phenylarsenic homocyclic compounds have been shown by

X-ray crystallography to be $(MeAs_5)$ and $(PhAs_6)$, respectively.³

Bis(pentafluorophenyl)arsinous chloride reacts with mercury to afford tetrakis(pentafluorophenyl)diarsine, $(C_6F_5)_2AsAs(C_6F_5)_2$, which after recrystallisation from benzene was characterised by elemental analyses and molecular weight measurements in chloroform. However, the analytical sample melted over a wide range, and examination of the ^{19}F n.m.r. spectrum showed the presence of five peaks at 133.5, 146.6, 152.0, 160.0, and 162.8 p.p.m., which is clearly inconsistent with the molecular formula $(C_6F_5)_2AsAs(C_6F_5)_2$. Sublimation of the diarsine *in vacuo* resulted in the separation of two isomers of tetrakis(pentaphenyl)diarsine. The ^{19}F n.m.r. spectrum of each isomer showed three peaks of relative intensity 2:1:2, which occurred at 133.5, 152.0, and 162.8 p.p.m. for the isomer subliming at 140° , and at 133.5, 146.6, and 160.0 p.p.m. for the isomer subliming at 230° . Examination of models suggests that the two isomers are rotamers, possibly *trans*- and *gauche*-forms of $(C_6F_5)_2AsAs(C_6F_5)_2$. The implied high energy barrier for the interconversion of the two isomers probably arises because of a balance of two factors. (i) Steric hindrance due to the *ortho*-fluorines, and (ii) the As-As bond distance, which is of the correct length to allow the formation of a *gauche*-form, but not of great enough length to allow free rotation. Alternative routes to the diarsine, *e.g.*, the reaction of $(C_6F_5)_2AsCl$ with $(CH_3)_3SiH$ or the reaction of

$(\text{C}_6\text{F}_5)_2\text{As}\cdot\text{S}\cdot\text{As}(\text{C}_6\text{F}_5)_2$ with mercury gave similar mixtures of the two isomers.

Reaction of the As—As compounds with aqueous

alkali gave a quantitative yield of pentafluorobenzene, and reaction with an excess of chlorine proceeded slowly to give $\text{C}_6\text{F}_5\text{Cl}$.

(Received, December 14th, 1966; Com. 993.)

¹ A. H. Cowley and R. P. Pinnell, *J. Amer. Chem. Soc.*, 1966, **88**, 4533.

² A. H. Cowley, A. B. Burg, and W. R. Cullen, *J. Amer. Chem. Soc.*, 1966, **88**, 3178.

³ J. H. Burns and J. Waser, *J. Amer. Chem. Soc.*, 1957, **79**, 859; K. Hedberg, E. W. Hughes, and J. Waser, *Acta Cryst.*, 1961, **14**, 369; S. E. Rasinnsen and J. Danielson, *Acta Chem. Scand.*, 1960, **14**, 1862.