

A New Synthesis of 1,1,3,3-Tetramethyl-2-selenourea

By DANIEL L. KLAYMAN* and ROBERT J. SHINE

(Division of Medicinal Chemistry, Walter Reed Army Institute of Research, Washington, D.C. 20012)


THE synthesis of 1,1,3,3-tetramethyl-2-selenourea (I) was recently accomplished by Jensen, Felbert, and Kägi¹ by refluxing 1,1,3,3-tetramethylurea in anhydrous benzene with phosphorus pentaselenide for three days. Compound (I) was obtained as an air- and moisture-sensitive material in 5.6% yield.

We have developed a simple preparation of the selenourea (I) in good yield from 1,1,3,3-tetramethyl-2-thiourea (II). The latter was treated with methyl iodide in ethanol solution to give a

quantitative yield of S-1,1,3,3-pentamethylisothiuronium iodide (III).² This S-methyl derivative was dissolved in 1 equiv. aqueous ethanolic sodium hydrogen carbonate, added immediately to an aqueous-ethanolic solution containing *ca.* 2 equiv. sodium hydroselenide,[†] and left at room temperature for *ca.* 17 hr. The solution was slowly acidified with 6N-sulphuric acid while nitrogen was passed through to displace the hydrogen selenide and methyl mercaptan liberated. The small quantity

† Hydrogen selenide, prepared by the dropwise addition of 6N-sulphuric acid to 2 equiv. of powdered Al_2Se_3 , was passed into an aqueous-ethanolic solution of 2 equiv. of sodium hydrogen carbonate at 0° to give a solution of sodium hydroselenide.

of elemental selenium present was removed by filtration; the colourless filtrate was concentrated to near dryness under reduced pressure at *ca.* 80°



and cooled. White selenourea (I) (70%) was obtained which, after recrystallization from hexane, was surprisingly stable in air, m.p. 79–81° (lit.,¹ m.p. 78–79°). The selenourea was further identified by elemental analysis and its i.r. spectrum (KBr pellet) which showed bands at 1517, 1360, 1375, 1085, and 920 cm^{-1} , in agreement with those reported by Jensen and Nielsen.³ The n.m.r. spectrum[‡] of the material exhibited a single peak at δ 3.16 (NMe).

This method has been applied to the preparation of other selenoureas by the displacement of methyl mercaptan from the S-methyl derivative of thiourea, as well as mono-, di-, and tri-substituted thioureas by the action of hydroselenide.

(Received, February 5th, 1968; Com. 137.)

‡ Determined on a Varian A-60 spectrometer as a 25% solution in CDCl_3 using Me_4Si as an internal standard.

¹ K. A. Jensen, G. Felbert, and B. Kägi, *Acta Chem. Scand.*, 1965, **20**, 281.

² H. Lecher and C. Heuck, *Annalen.*, 1924, **438**, 169.

³ K. A. Jensen and P. H. Nielsen, *Acta Chem. Scand.*, 1966, **20**, 597.