

## A Novel Series of Transition-metal Chelates of Diphenylphosphinothiolythiourea Anion


By IWAO OJIMA, TOSCHITAKE IWAMOTO, TAKAHARU ONISHI, NAOKI INAMOTO,\* and KENZI TAMARU  
(Department of Chemistry, Faculty of Science, The University of Tokyo, Hongo, Tokyo, Japan)

**Summary** Two thiourea derivatives, 3-(diphenylphosphinothioyl)-1-phenylthiourea and 3-(diphenylphosphinothioyl)-1,1-diethylthiourea, behave as bidentate ligands towards Ni<sup>II</sup> and Pd<sup>II</sup> to produce new square-planar complexes, while the latter ligand gives a tetrahedral complex with Co<sup>II</sup>.

THE thiourea derivatives, 3-(diphenylphosphinothioyl)-1-phenylthiourea (PTTU-A) and 3-(diphenylphosphinothioyl)-1,1-diethylthiourea (PTTU-B), obtained by addition of diphenylphosphinothioly isothiocyanate with aniline and diethylamine, respectively, react with bivalent metal halides to give a series of novel metal chelates.

A solution of PTTU-B (4 mmoles) in dichloromethane (20 ml.) was allowed to react with nickel(II) chloride hexahydrate (2 mmoles) in ethanol at room temperature. The precipitate thus formed was recrystallized from dichloromethane-ethanol solution. A crystalline bis-[3-(diphenylphosphinothioyl)-1,1-diethylthioureato]nickel(II) (Ni-PTTU-B) was obtained in nearly quantitative yield. The Ni<sup>II</sup> and Pd<sup>II</sup> complexes of PTTU-A and the Pd<sup>II</sup> and Co<sup>II</sup> complexes of PTTU-B were prepared in a similar way. Ni-PTTU-A: dark green needles, m.p. 193° (dec.),  $\lambda_{\max}$

(CH<sub>2</sub>Cl<sub>2</sub>) 15.7 (log  $\epsilon$  2.36), 18.0  $\kappa\kappa$ (2.30). Pd-PTTU-A: orange-yellow prisms, m.p. 215° (dec.). Ni-PTTU-B: dark green needles, m.p. 200° (dec.),  $\lambda_{\max}$  (CH<sub>2</sub>Cl<sub>2</sub>) 15.5(2.24) and 17.6  $\kappa\kappa$ (2.24). Pd-PTTU-B: pale-yellow prisms, m.p. 244° (dec.). Co-PTTU-B: bright green needles, m.p. 205–206°,  $\chi_g = 11.9$  c.g.s. e.m.u. (room temp.),  $\lambda_{\max}$  (CH<sub>2</sub>Cl<sub>2</sub>) 4.65 (1.50), 7.87(2.41), 14.3(2.57), and 15.7  $\kappa\kappa$ (2.54).



M-PTTU-A: R<sup>1</sup>=Ph, R<sup>2</sup>=H  
M-PTTU-B: R<sup>1</sup>=R<sup>2</sup>=Et  
M=Ni, Pd, Co

The elemental analyses for these complexes were in good agreement with calculated values. All these complexes have the same metal : ligand ratio (1 : 2). Their solubilities in benzene were not high enough for us to determine their molecular weights except for the case of Co-PTTU-B,

$M$  (found) 751; calc. for monomer: 753. Except for Co-PTTU-B, they are all diamagnetic at room temperature, which suggests that the complexes of Ni<sup>II</sup> and Pd<sup>II</sup> have square-planar configurations. Their crystal structures appear to be isomorphous according to the powder X-ray diffraction patterns.

The effective magnetic moment of Co-PTTU-B,  $\mu_{\text{eff}}$  = 4.50 B.M., corresponds to the high-spin state of  $d^7$  Co<sup>II</sup> in a tetrahedral configuration. Its powder X-ray diffraction pattern was quite different from those of Ni- and Pd-PTTU-B. The electronic spectrum also supported the tetrahedral configuration of Co<sup>II</sup> in Co-PTTU-B. The values of  $B$  (684 cm.<sup>-1</sup>) and  $10 Dq$  (4788 cm.<sup>-1</sup>) calculated on the basis of the Tanabe-Sugano diagram, are in reasonable agreement with those reported on several tetrahedral Co<sup>II</sup> complexes.<sup>1</sup> Cobalt(II) in Co-PTTU-B has a tetrahedral configuration, in contrast with the case of bis(dithioacetyl-acetonato)cobalt(II), which has a square-planar structure.<sup>2</sup>

In the i.r. spectra of the chelates with the square-planar

structure, two metal-sulphur stretching vibrations were expected to appear in the far-i.r. region.<sup>3</sup> The bands most sensitive to metals, appearing in the region 420—320 cm.<sup>-1</sup>, are tentatively assigned to these vibrations in view of the covalent nature of the metal-sulphur bond and the mass

*Characteristic bands in the far-i.r. spectra (cm.<sup>-1</sup>)*

	P=S	M-S(1)	M-S(2)
Ni-PTTU-A .. ..	584	417	358
Pd-PTTU-A .. ..	582	412	336
Ni-PTTU-B .. ..	580	391	340
Pd-PTTU-B .. ..	579	380	329

effect (Table). The tentative assignments of P=S stretching bands are also given in the Table.

We thank Professor Ichiro Nakagawa for his advice and discussion on the i.r. spectroscopic results.

(Received, October 20th, 1969; Com. 1590.)

<sup>1</sup> A. B. P. Lever, "Inorganic Electronic Spectroscopy," Elsevier, Amsterdam, 1968, pp. 322—328.

<sup>2</sup> R. Beckett and B. F. Hoskins, *Chem. Comm.*, 1967, 909.

<sup>3</sup> M. Mikami, I. Nakagawa, and T. Shimanouchi, *Spectrochim. Acta*, 1967, **23**, A, 1037; K. Nakamoto, J. Fujita, R. A. Condrate, and Y. Morimoto, *J. Chem. Phys.*, 1963, **39**, 423.