

Formation of Chlorobenzene by the Reaction of Phenylthallium(III) Compounds with Copper Chloride

By KATSUHIKO ICHIKAWA* and YOSHIHIRO IKEDA

(Department of Hydrocarbon Chemistry, Faculty of Engineering, Kyoto University, Yoshida, Kyoto, Japan)


and SAKAE UEMURA

(Institute for Chemical Research, Kyoto University, Uji, Kyoto, Japan)

Summary The reaction of phenylthallium(III) compounds with CuCl_2 and CuCl in AcOH , MeCN , and tetrahydrofuran gives mainly chlorobenzene.

REACTIONS of organometallic compounds with metal salts are of current interest.¹ In the case of aromatic thallium compounds, it was reported recently that arylthallium bis(trifluoroacetate) reacts with lead tetra-acetate to form the aryl acetate and with potassium cyanide (under u.v. irradiation) to form the aryl cyanide.²

We report that thallium attached to an aromatic ring PhTlXY (I) can be replaced by chlorine by treatment with CuCl_2 or CuCl in AcOH , MeCN containing LiCl and tetrahydrofuran.



- (I) a; X = OAc, Y = ClO_4^\dagger
 b; X = Y = Cl
 c; X = Ph, Y = Cl
 d; X = Y = OAc
 e; X = Y = CF_3CO_2

AcOH solutions of phenylthallium(III) compounds and copper salts were kept at various temperatures for various reaction periods. The main organic product was chlorobenzene which was identified by g.c., i.r., and n.m.r. spectroscopy. Small amounts of benzene and biphenyl were also obtained. The results are shown in the Table.

The nature of X and Y has little effect on the yields of

[†] Prepared by the reaction of aromatic substrates with thallium(III) acetate in AcOH containing HClO_4 .⁵


chlorobenzene, except in the case $X = Y = \text{OAc}$ (Id). Both CuCl_2 and CuCl were effective, but $\text{Cu}(\text{OAc})_2$ was inactive even in the presence of LiCl . The effective

Although the elucidation of the detailed mechanism requires further experiment, it is clear that the present reaction is different from that reported by McKillop *et al.*,³

Phenylthallium(III) compound (10 mmol)	Copper salt (mmol)	Solvent (50 ml)	Reaction temp (°)	Reaction time (h)	Products (yield, %) ^a		
					Benzene	Chlorobenzene	Biphenyl
	$\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$						
PhTl(OAc)(ClO ₄) (Ia)	10	AcOH	117	1	4	56	trace
"	10	"	117	5	4	59	2
"	10	"	117	10	2	59	trace
"	20	"	117	5	2	60	trace
"	10	" ^b	117	5	3	12	9
"	10	" ^c	117	5	1	53	trace
"	10	MeCN ^d	84	5	2	28	3
"	10	THF	67	5	—	20	22
	CuCl						
" ^e	20	AcOH	117	5	4	42	trace
	$\text{Cu}(\text{OAc})_2$						
" ^e	20	"	117	5	13	—	2
"	—	"	117	5	24	—	trace
	$\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$						
PhTiCl ₂ (Ib)	10	"	117	5	2	67	trace
"	1	MeCN ^d	84	20	2	32	21
Ph ₂ TlCl (Ic)	10	AcOH	117	5	10	45	trace
PhTl(OAc) ₂ (Id)	10	AcOH	117	5	19	16	4
PhTl(CF ₃ CO ₂) ₂ (Ie)	10	"	117	5	5	30	trace

^a Based on phenylthallium(III) compound; ^b NaOAc (10 mmol) added; ^c LiCl (40 mmol) added; ^d LiCl (10 mmol) added; ^e 20 mmol; ^f LiCl (20 mmol) added.

reagent, therefore, appears to be the copper chlorides and not Cl^- . Addition of NaOAc to the reaction mixture resulted in a decreased yield of chlorobenzene and an increased yield of biphenyl, showing that the nature of the reaction had changed probably because of the formation of $(\text{CuClOAc})_2$.³ In AcOH solution, PhTl(OAc)(ClO₄) did not react with the copper salts at 70°, while a 20% yield of chlorobenzene was obtained at 67° in tetrahydrofuran. This shows that the solvent has an important effect on the reaction.



($\text{ArTlI}_2 \rightarrow \text{ArI}$), since PhTiCl₂ (Ib) remained unchanged even after a prolonged reaction period. An $\text{S}_{\text{N}}2$ -type substitution of aliphatic thallium(III) salts with halide ion⁴ is not applicable to the present case. Formation of an aromatic copper compound by metal exchange organothallium and copper salt also appears unlikely, since the yields of biphenyl to be expected from that intermediate were not obtained.

A four-centre reaction is the most plausible explanation of the results. The fact that greater amounts of PhCl were formed than of CuCl_2 consumed in the reaction of PhTiCl₂ (Ib) supports this mechanism.

(Received, November 24th, 1970; Com. 2036.)

¹ R. F. Heck, *J. Amer. Chem. Soc.*, 1968, **90**, 5518, 5539; P. M. Henry, *Tetrahedron Letters*, 1968, 2285; M. O. Unger and R. A. Fouty, *J. Org. Chem.*, 1969, **34**, 18; H. Stangl and R. Jira, *Tetrahedron Letters*, 1970, 3589.

² E. C. Taylor, H. W. Altland, R. H. Danforth, G. McGillivray, and A. McKillop, *J. Amer. Chem. Soc.*, 1970, **92**, 3520.

³ S. Uemura, T. Hiramoto, Y. Takagaki, and K. Ichikawa, *J. Chem. Soc. Japan, Ind. Chem. Sect.*, 1969, **72**, 2390.

⁴ M. D. Johnson, *Chem. Comm.*, 1970, 1037.

⁵ K. Ichikawa, S. Uemura, T. Nakano, and E. Uegaki, *Bull. Chem. Soc. Japan*, in the press.