

## Thermal Rearrangements of Some $\alpha$ -Aroyl Epoxides into Dioxoles


By ALBERT ROBERT\* and BRUNO MOISAN

(Groupe de recherches de physicochimie structurale Université de Rennes, Avenue du Général Leclerc, Rennes, France)

**Summary** Thermal rearrangement of  $\alpha$ -cyano- $\alpha$ -oxo-epoxides quantitatively affords the corresponding dioxoles, which are formed by initial cleavage between the two carbon atoms of the starting epoxide ring.

SEVERAL examples of thermal rearrangements of vinylaziridines into  $\Delta^2$ - or  $\Delta^3$ -pyrrolines have been reported,<sup>1</sup> and more recently it has been shown that thermolysis of vinyl epoxides gives rise to dihydrofurans *via* carbonyl ylides.<sup>2</sup> A similar isomerization has been observed with aziridines substituted by oxo,<sup>3</sup> ester, amide, or nitrile groups,<sup>4,5</sup> but there are no reports of similar reactions of  $\alpha$ -oxo-epoxides.

We have shown that thermolysis of epoxides  $\alpha\alpha$ -disubstituted with two electron-withdrawing groups gives carbonyl ylides which can be trapped by various dipolarophiles.<sup>6-9</sup> These results suggest that the thermolysis of  $\alpha$ -oxo-epoxides (**1**) may be similar to that of  $\alpha$ -oxo-aziridines; thermolysis of (**1**) provides, in fact, a new route to the dioxoles (**2**).




$R^1 = \text{Ph}$  or  $p\text{-ClC}_6\text{H}_4$ ,  $R^2 = \text{H}$  or  $\text{Ph}$ ,  $\text{Ar} = \text{Ph}$  or  $p\text{-MeC}_6\text{H}_4$

The new epoxides (**1**) were prepared<sup>10</sup> by Grignard reaction of  $\text{ArMgBr}$  with the dicyano-epoxides (**3**), and were thermolysed in boiling toluene under nitrogen. The dioxoles (**2**) were obtained in quantitative yield and gave satisfactory elemental analyses.

The structures (**2**) were based on n.m.r., u.v., and i.r. spectra. It is of interest that the i.r. spectra contain bands due to a conjugated nitrile group and also medium intensity absorptions which may be attributed to a cyclic acetal [1018—1093 and 1308  $\text{cm}^{-1}$  (Nujol)]<sup>6,11</sup> but lack  $\text{C}=\text{O}$  absorption bands.

It has been established that epoxides gem-disubstituted with electron-withdrawing groups give carbonyl ylides,<sup>7-9</sup> and so, as with  $\alpha$ -oxo-aziridines,<sup>3</sup> it is reasonable to suppose that isomerization of the epoxides (**1**) proceeds *via* a stabilised 1,3-dipolar intermediate (Scheme).



SCHEME

1,3-Dipolar cycloadditions are very sensitive to steric effects, *e.g.*, gem-dicyano-epoxides substituted by two phenyl groups undergo 1,3-cycloadditions to only a limited extent,<sup>9</sup> so it is likely that steric effects would not be so important in the intramolecular cyclisation leading to (**2**).

Isomerization of the diphenyl epoxides (**1**;  $R^1 = R^2 = \text{Ph}$ ) is faster than that of the monoaryl epoxides (**1**;  $R^2 = \text{H}$ ) (Table), which suggests that the rate determining step is

TABLE  
Preparation of dioxoles (**2**)

Dioxoles ( <b>2</b> )	$R^2$	Time <sup>a</sup>	M.p. ( $^{\circ}\text{C}$ )	
$\text{Ar} = \text{Ph}$	$R^1 = \begin{cases} \text{Ph} \\ p\text{-ClC}_6\text{H}_4 \\ \text{Ph} \end{cases}$	H	24 h	77—78
	H	24 h	79—80	
	Ph	1 h	118—119	
$\text{Ar} = p\text{-MeC}_6\text{H}_4$	$R^1 = \begin{cases} \text{Ph} \\ p\text{-ClC}_6\text{H}_4 \\ \text{Ph} \end{cases}$	H	24 h	74—75
	H	24 h		
	Ph	1 h	127—128	

<sup>a</sup> Time required for quantitative isomerization.

formation of the carbonyl ylide (**4**). A second phenyl group may stabilize the 1,3-dipolar intermediate and so favour its formation.

It is thus not surprising that 1,3-dipolar cycloaddition was not observed, even when the epoxides (**1**) were thermo-

lysed in presence of a good dipolarophile such as *p*-nitrobenzaldehyde;<sup>7</sup> the dioxole (**2**) was then formed, by the intramolecular process.

(Received, 20th December 1971; Com. 2157.)

<sup>1</sup> H. W. Heine, *Angew. Chem. Internat. Edn.*, 1962, **1**, 528; P. Scheiner, *J. Org. Chem.*, 1967, **32**, 2628; R. S. Atkinson and C. W. Rees, *Chem. Comm.*, 1967, 1232.

<sup>2</sup> J. C. Paladini and J. Chucho, *Tetrahedron Letters*, 1971, 4383.

<sup>3</sup> A. Padwa and W. Eisenhard, *Chem. Comm.*, 1968, 380; J. E. Baldwin, R. G. Pudussery, A. K. Quereschi, and B. Sklarz, *J. Amer. Chem. Soc.*, 1968, **90**, 5325; F. Texier and R. Carrie, *Compt. rend.*, 1970, **271**, 958.

<sup>4</sup> A. Foucaud and M. Baudru, *Compt. rend.*, 1970, **271**, 1613.

<sup>5</sup> M. Baudru, Thèse de 3ème cycle, Université de Rennes, 1970.

<sup>6</sup> A. Robert, J. J. Pommeret, and A. Foucaud, *Compt. rend.*, 1970, **270C**, 1739.

<sup>7</sup> A. Robert, J. J. Pommeret, and A. Foucaud, *Tetrahedron Letters*, 1971, 231.

<sup>8</sup> J. J. Pommeret and A. Robert, *Compt. rend.*, 1971, **272C**, 333.

<sup>9</sup> J. J. Pommeret and A. Robert, *Tetrahedron Letters*, 1971, 2405.

<sup>10</sup> J. M. Normant and J. Cantacuzene, *Tetrahedron Letters*, 1971, 2405.

<sup>11</sup> J. Gelas, S. Michaud, and R. Rambaud, *Tetrahedron Letters*, 1970, 1533.