

Lamellicolic Anhydride—a Heptaketide Naphthalic Anhydride from *Verticillium lamellicola*

By NORMAN J. MCCORKINDALE,* ALLAN MCRITCHIE, and STEPHEN A. HUTCHINSON

(Joint Mycology Laboratories, Departments of Chemistry and Botany, The University of Glasgow,
Glasgow G12 8QQ)

Summary The structure of lamellicolic anhydride from *Verticillium lamellicola* has been established using a number of transformations highly characteristic of the 2,7-dihydroxy-1,8-naphthalic anhydride system present, and by conversion into the naphthalic anhydride from *Penicillium herquei*.

THE groups of fungal metabolites related to atrovénin and to duclauxin appear to have in common a heptaketide skeleton¹ which is modified by prenylation in the former case and by dimerisation in the latter.² We report the isolation from a strain of the fungus *Verticillium lamellicola* (F. E. V. Smith) of a number of metabolites which are based on the simple heptaketide skeleton.†


Broth extracts of *V. lamellicola* have weak antibacterial and antifungal activity³ due to the presence of traces of an antibiotic which as yet has not been identified. Close to this in polarity (but virtually inactive) was the major metabolite, lamellicolic anhydride (I), R_F 0.35 in MeOH-CHCl₃ (1:9), which formed yellow needles from MeOH-CHCl₃, decomposing over 300°, C₁₃H₈O₈ (M^+ at 260.0323). The solubility of this in aqueous sodium hydrogen carbonate, its twin i.r. absorption at 1700 and 1650 cm⁻¹, and

complex u.v. spectrum apparently showing a reversible hypsochromic shift on basification are all characteristic of the 1,8-naphthalic anhydride system present in (I).⁴ The presence in (I) of three phenolic OH groups was indicated by the formation of the tri-*O*-acetyl derivative (II), m.p. 177–179° and tri-*O*-methyl derivative (III), m.p. 290° (decomp.), ν_{\max} 1750, 1715 cm⁻¹.


Aryl methoxy-groups *peri* to carbonyl groups are often readily cleaved by magnesium iodide-etherate⁵ and with this reagent, (III) afforded the monomethoxy-derivative (IV), m.p. 260° (decomp.), also obtained as a minor product in the methylation of (I). Treatment of (III) with hot aqueous methylamine confirmed the presence of the chelated anhydride system in a remarkable manner, the aminoimides (V), m.p. 225–228°, and (VI), m.p. 244–246°, being formed (NH resonances at τ -1.5 and -0.3, respectively).

The environments of the aryl protons in (I) and (III) were confirmed by the detection of 6-H-*C*-Me coupling and, in the latter case, by nuclear Overhauser effects. Thus in CF₃CO₂H the signal at τ 2.73 (6-H) was enhanced 21 and 11% per irradiation at τ 6.94 (*C*-Me) and 5.73 (*O*-Me), respectively, while the signal at τ 3.14 (3-H) was enhanced 0 and 37, respectively, by irradiation at the same frequencies. Successive treatment of (I) with aqueous NaOH


† Biosynthetic evidence in support of this assertion will be reported elsewhere.




- (I) $R^1 = R^2 = H$
 (II) $R^1 = R^2 = Ac$
 (III) $R^1 = R^2 = Me$
 (IV) $R^1 = H, R^2 = Me$




- (V) $R = NHMe$
 (VI) $R = OMe$




- (VII) $R = CO_2Me$
 (VIII) $R = H$



- (IX) $R = H$
 (X) $R = Me$



(XI)



(XII)

(N₂) and diazomethane gave (VII), m.p. 126—128°, and (VIII), m.p. 93—95°, in which the new aryl protons introduced by decarboxylation could be seen to be *meta* coupled (J 3 Hz) to the appropriate proton originally present. If, after treatment of (I) with alkali, the solution was allowed to stand in air, the orange quinone (IX), m.p. 250° (decomp.) was obtained.⁶ The 8-H proton appeared in the n.m.r. spectrum of the corresponding dimethyl ether (X), m.p. 172—173°, as a doublet (J 2 Hz) at τ 2.42, and the singlet corresponding to 3-H (τ 3.9) showed substantial enhancement upon irradiation at *ca.* τ 6.1 (OMe). The quinone (IX) was also isolated from the broth extracts. As yet we have been unable to disprove that this is formed naturally, at least to some extent, during the fermentation process.

The evidence summarised above conclusively establishes the structure of lamellicolic anhydride (I) and this has been confirmed by conversion into the anhydride (XI),⁷ a sample of which was prepared from herqueinone⁸ provided by Professor R. Thomas. Conclusive proof has been provided independently⁹ by synthesis of (III) and (I) *via* the tetramethoxyphenalenone (XII).¹⁰ Two minor metabolites closely related to (I) will be discussed elsewhere.

All new compounds gave the requisite spectral and analytical data.

(Received, 4th December 1972; Com. 2018.)

- ¹ R. Thomas, *Biochem. J.*, 1961, **69**, 807; U. Sankawa, H. Taguchi, Y. Ogihara, and S. Shibata, *Tetrahedron Letters*, 1966, 2883.
² W. B. Turner, 'Fungal Metabolites,' Academic Press, New York and London, 1971, p. 150.
³ U.K. Patent Application No. 36,963/1954.
⁴ R. G. Cooke, *Chem. and Ind.*, 1955, 142; D. H. R. Barton, P. de Mayo, G. A. Morrison, and H. Raistrick, *Tetrahedron*, 1959, **6**, 48.
⁵ B. W. Bycroft and J. C. Roberts, *J. Chem. Soc.*, 1963, 4868.
⁶ P. M. Baker and B. W. Bycroft, *Chem. Comm.*, 1968, 71.
⁷ N. Narasimhachari and L. C. Vining, *Canad. J. Chem.*, 1963, **41**, 641.
⁸ J. S. Brooks and G. A. Morrison, *J.C.S. Perkin I*, 1972, 421.
⁹ B. W. Bycroft, personal communication.
¹⁰ B. W. Bycroft and A. J. Eglington, *Chem. Comm.*, 1968, 72.