

## Neutral Conversion of Aldoximes into Nitriles at Low Temperature

By H. GRANT FOLEY and DAVID R. DALTON\*

(Department of Chemistry, Temple University, Philadelphia, Pennsylvania 19122)

*Summary* (*E*)- and (*Z*)-Aldoximes can be induced to lose water to yield nitriles at, or slightly above, room temperature in dichloromethane solution by treatment with 1,1'-dicarbonylbi-imidazole.

THE dehydration of aldoximes to nitriles often involves heating and/or acidic or basic catalysis, which may be detrimental to complex molecules.

In the course of our investigations on the structures of some unusual nitrones<sup>1</sup> we examined the reaction between (*E*)- and (*Z*)-aldoximes and 1,1'-dicarbonylbi-imidazole. When, for example, one equivalent of the carbonylbi-imidazole in dichloromethane solution was added to one equivalent of (*Z*)-*p*-chlorobenzaldoxime in the same solvent at room temperature, immediate vigorous evolution of carbon dioxide occurred and, after extraction of the

TABLE

*Nitriles obtained from selected oximes on treatment with carbonylbi-imidazole*

	Oxime <sup>a</sup>	Nitrile	Yield (%)
(1)	( <i>E</i> )- <i>p</i> -Nitrobenzaldoxime <sup>b</sup>	<i>p</i> -Nitrobenzotrile	99
(2)	( <i>E</i> )- <i>p</i> -Chlorobenzaldoxime	<i>p</i> -Chlorobenzotrile	95
(3)	( <i>Z</i> )- <i>p</i> -Chlorobenzaldoxime <sup>c</sup>	<i>p</i> -Chlorobenzotrile	98
(4)	2,2-Dimethylpropanaldoxime <sup>d</sup>	2,2-Dimethylpropionitrile	95

<sup>a</sup> The oximes were prepared by standard procedures. <sup>b</sup> Heated under reflux for 6 h in dichloromethane. <sup>c</sup> Prepared according to the method of C. R. Hauser and D. S. Hoffenberg, *J. Org. Chem.*, 1955, **20**, 1491. <sup>d</sup> This mixture of (*E*)- and (*Z*)-oximes reacted explosively with carbonylbi-imidazole when mixed neat. The crude mixture of oximes was used directly.

imidazole (or filtration through an alumina column), *p*-chlorobenzotrile was isolated as the only product. On the other hand, (*E*)-*p*-methoxybenzaldoxime required, for 75% conversion into the corresponding nitrile, heating under reflux in dichloromethane solvent for 3 h; conditions which we consider mild.

The Table presents a series of typical oximes which have

been successfully converted, by the method indicated, into the corresponding nitriles in the isolated yields shown.†

One of us (H.G.F.) acknowledges financial aid from the Fund for Scientific Education, Inc., Philadelphia, Pennsylvania.

(Received, 1st June 1973; Com. 790.)

† The products were compared to known materials. Satisfactory analyses have been obtained. The yields reported have been obtained more than once for each compound shown in the Table.

<sup>1</sup> D. R. Dalton, K. N. Trueblood, and M. R. Murphy, *Tetrahedron Letters*, 1973, 779.