



## Pyrolytic Generation of Acyloxycarbenes and their Rearrangement to 1,2-Diketones

By ROGER F. C. BROWN,\* FRANK W. EASTWOOD, and GABRIELLE L. McMULLEN  
(Department of Chemistry, Monash University, Clayton, Victoria, 3168, Australia)

*Summary* Acyloxycarbenes are generated by thermal fragmentation of 5-benzoyloxy-1,3-dioxan-4,6-diones (**1** and **2**) and by thermal  $\alpha$ -elimination of benzoic acid from 3-benzoyloxyphthalide (**6**).

IN an attempt to generate the elusive parent methyleneketene,  $\text{CH}_2=\text{C}=\text{O}$ , by a route akin to that used for higher members of the series<sup>1</sup> we examined the flash vacuum pyrolysis of the benzoate (**1**). Pyrolysis of (**1**) (460 °C, 0.5 mmHg) through silica gave a yellow liquid containing



acetone and 1-phenylpropane-1,2-dione (**4**; 90% yield). Similarly, pyrolysis of the phenyl-substituted benzoate (**2**) (460 °C, 0.05 mmHg) gave acetone and benzil (**5**; 81%). We propose that these diketones are formed by acyl migration in the intermediate acyloxycarbenes (**3**), which in turn arise from ketens PhCO<sub>2</sub>CR=CO by very ready decarbonylation.

Acyloxycarbenes are little known except for the cyclic species (**7**), generated photochemically in solution by irradiation of the strained diketone (**8**)<sup>2,3</sup> and studied by Staab and Ipaktschi.<sup>3</sup> We have now obtained this diketone (**8**; 33%) by pyrolysis of 3-benzoyloxyphthalide (**6**)<sup>4</sup> (560 °C, 0.1 mmHg). This process presumably involves  $\alpha$ -elimination<sup>5</sup> of benzoic acid (91% crude yield) and ring contraction of the resulting carbene (**7**). The diketone (**8**) was accompanied by a small hydrocarbon fraction which contained biphenylene, probably formed by decarbonylation of (**8**).<sup>6</sup>

This work was supported by the Australian Research Grants Committee.

(Received, 31st January 1975; Com. 115.)

<sup>1</sup> R. F. C. Brown, F. W. Eastwood, and K. J. Harrington, *Austral. J. Chem.*, 1974, **27**, 2372.

<sup>2</sup> H. A. Staab and J. Ipaktschi, *Chem. Ber.*, 1968, **101**, 1457.

<sup>3</sup> R. F. C. Brown and R. K. Solly, *Tetrahedron Letters*, 1966, 169.

<sup>4</sup> D. D. Wheeler, D. C. Young, and D. S. Erley, *J. Org. Chem.*, 1957, **22**, 547.

<sup>5</sup> P. C. Oele and R. Louw, *Tetrahedron Letters*, 1972, 4941.

<sup>6</sup> M. P. Cava in 'Aromaticity,' Chem. Soc. Special Publication No. 21, 1967, p. 168.