Phenyl Vinyl Sulphoxide, a Vinyl Carbonium Ion Synthetic Equivalent


By G. A. KOPPEL* and M. D. KINNICK

(The Lilly Research Laboratories, Eli Lilly and Company, Indianapolis, Indiana 46206)

Summary Reaction of ethyl 2-oxocyclohexanecarboxylate (I) with phenyl vinyl sulphoxide (II) and sodium hydride followed by pyrolysis of the resulting phenylsulphonylethyl compound (III) gives the 1-vinyl compound (IV).

In view of the isolation and structural proof of the antitumour sesquiterpene vernolepin,¹ which has an angular vinyl substituent, we have investigated methods of introducing a terminal vinyl group.² The early introduction of such a group as a protected equivalent was considered to be a crucial step in our syntheses. We thought that the work of Tsuchihashi and his co-workers³ demonstrating that p-tolyl vinyl sulphoxides undergo Michael reactions and that of Trost and Saltzmann⁴ using the pyrolysis of sulphoxides in generating olefins could be combined, providing the synthetic equivalent of the vinyl carbonium ion, $(A) \rightarrow (B)$. A method for vinyl group synthesis has been reported recently,⁵ and we report here the reaction of compound (I) with phenyl vinyl sulphoxide (II), which was prepared by treatment of phenyl disulphide with vinyllithium and subsequent oxidation with peroxy-acid.

Treatment of (I) (2 equiv.) with NaH (1 equiv.) and phenyl vinyl sulphoxide (1 equiv.) in tetrahydrofuran at reflux afforded after workup and preparative thick-layer chromatography a 50% yield of (III) as a mixture of diastereoisomers; τ (CDCl_a) 2.5 (m, Ph), 5.8 (q, CO₂Et),



6.8-8.8 (side-chain and ring H); m/e 223 (M^+ + 1) and 125 (100%). Subsequent pyrolysis of (III) in refluxing toluene for 72 h afforded the previously reported⁵ compound (IV) (60%).

(Received, 6th March 1975; Com. 283.)

- ⁴ B. M. Trost and T. N. Salzmann, J. Amer. Chem. Soc., 1973, 95, 6840. ⁵ T. Oishi, H. Takechi, and Y. Ban, Tetrahedron Letters, 1974, 3757.

¹S. M. Kupchan, R. J. Hemingway, D. Werner, A. Karim, A. T. McPhail, and G. A. Sim, J. Amer. Chem. Soc., 1968, 90, 3596.

³ Two syntheses of the bicyclic vernolepin skeleton have been reported, see P. A. Grieco and K. Hiroi, Tetrahedron Letters, 1973, 1831; R. D. Clark and C. H. Heathcock, ibid., 1974, 2027.

G. Tsuchihashi, S. Mitamura, S. Inoue, and K. Ogura, Tetrahedron Letters, 1973, 323.