

Synthesis of 2,3-Dihydro-1*H*-phenothiazin-4(10*H*)-ones

By SEIJI MIYANO,* NOBUHIRO ABE, and KUNIHIRO SUMOTO

(Faculty of Pharmaceutical Sciences, Fukuoka University, Fukuoka, Japan)


Summary 2,3-Dihydro-1*H*-phenothiazin-4(10*H*)-ones (**4**)—(**7**) were prepared by condensation of *o*-aminobenzenethiol (**1**) with cyclohexane-1,3-diones (**2**) in Me₂SO.

CURRENT interest¹ in the nucleophilic reactions of enamino-ketone groups prompts us to report a novel synthesis of the 2,3-dihydro-1*H*-phenothiazin-4(10*H*)-ones (**4**)—(**7**). The

dimethyl compound (**6**), for example, was prepared by heating a mixture of (**1**) (1.25 g) and dimedone (**2**; R¹ = R² = Me) (1.4 g) in Me₂SO (5 ml) at 155 °C for 0.5—1 h. The product (**6**), m.p. 262—263 °C (decomp.) (from MeOH), separated as orange plates on cooling. Compounds (**4**) (42% yield), (**5**) (66%), and (**7**) (77%) were prepared similarly.†

Since compound (**1**) is readily oxidized to bis(*o*-aminophenyl) disulphide under the reaction conditions, and since we have shown that this disulphide also undergoes condensation with the diones (**2**) to give compounds (**4**)—(**7**), we suggest that the enamino-ketones (**3**) are intermediates in the reaction.

This one-step synthesis is more convenient than the two-stage synthesis which starts with *o*-nitrobenzenesulphenyl chloride and cyclohexane-1,3-diones.²



(Received, 17th July 1975; Com. 819.)

† All products were adequately characterized by elemental analysis and spectroscopic methods.

¹ G. H. Alt and A. J. Speziale, *J. Org. Chem.*, 1964, **29**, 794; H. J. Teuber and R. Braun, *Chem. Ber.*, 1967, **100**, 1353; H. J. Teuber, E. Worbs, and D. Cornelius, *ibid.*, 1968, **101**, 3918; C. Ruangsiyanand, H.-J. Rimak, and F. Zymalkowski, *ibid.*, 1970, **103**, 2403; S. Miyano and N. Abe, *Chem. Pharm. Bull. (Japan)*, 1972, **20**, 1588; Y. Yoshimoto, N. Ishida, and T. Hiraoka, *Tetrahedron Letters*, 1973, **39**, and references therein.

² V. I. Shvedov, L. B. Altukhova, V. M. Lyubchanskaya, and A. N. Grinev, *Khim. Geterotsikl. Soedin.*, 1972, 1901 (*Chem. Abs.*, 1973, **77**, 43395k).