


Novel Type of Metallation Reaction: Cyclometallation of Heterocyclic Amines by Dicyclopentadienylmethyltitanium

By BERT KLEI and JAN H. TEUBEN*


(Laboratory of Inorganic Chemistry, The University, Nijenborgh 16, 9747 AG Groningen, The Netherlands)

Summary In a new type of metallation reaction 2-substituted pyridines and quinolines are metallated at the 6- and 2-position, respectively by dicyclopentadienylmethyltitanium prepared *in situ*.




SCHEME 1

METALLATION of heterocyclic amines by several transition metals has been reported. The cyclometallated products formed normally consist of five-membered chelate rings [*e.g.* compounds (1)—(3)¹⁻⁵]. However, similar metallations have not been described for Ti. The activating properties of compounds Cp₂TiR towards unsaturated substrates⁶ prompted us to investigate cyclometallation reactions of these compounds with heterocyclic amines analogous to those giving (1)—(3).




We now report an unusual cyclometallation of 2-substituted pyridines and quinolines (*e.g.*, 2-methyl-, 2-phenyl-, or 2-vinyl-pyridine, quinoline, or 8-methylquinoline) by Cp₂TiMe prepared *in situ* (Cp = η⁵-C₅H₅).[†] The metallation takes place at the 6-position of the pyridines and at the 2-position of the quinolines, yielding products probably containing a three-membered titanacycle in which the ligands act as bidentate three-electron donors [*e.g.* compounds (4) and (5)].



In a typical reaction Cp₂TiCl (made by reduction of Cp₂-TiCl₂ with 1 equiv. of Pr¹MgCl⁷) was treated with 1 equiv. of MeLi at -78 °C. After 1 h the Cp₂TiMe formed was treated with a slight excess of the heterocyclic amine. A dark green mixture formed, presumably owing to co-ordination of the Lewis base to the intermediate Cp₂TiMe. On warming to room temperature the colour changed to brown-purple while methane (g.l.c.) was evolved (see Scheme 1; pyridines are used as an example; the same Scheme holds for the quinolines). Methane formation is a normal consequence of metallation which is also observed for other cyclometallations.⁸


[†] The identity of Cp₂TiMe as the reactive species was not completely established. It is possible that addition compounds of Cp₂TiMe and Mg or Li compounds are the true active species.

The purple crystalline compounds (isolated in *ca.* 25% yield by crystallization from *n*-pentane) are stable at room temperature and probably have structures (4) and (5). This was established by elemental analysis, i.r. spectroscopy (characteristic absorptions of η⁵-C₅H₅ groups; specific C-H out of plane deformation modes for the ligands), molecular weight (all compounds are monomeric), and magnetic moments (close to the spin-only value of 1.73 B.M. expected for a Ti^{III} compound Cp₂TiR). The existence of a Ti-C σ bond at the 6-position in the pyridines and at the 2-position in the quinolines was demonstrated by reactions of the compounds with I₂ and D₂O-DCl. In all cases the expected 6-iodo- or 6-deuterio-pyridines and 2-iodo- or 2-deuterio-quinolines, respectively were obtained quantitatively (Scheme 2), as established by i.r., mass, and ¹H-



SCHEME 2

n.m.r. spectroscopy. The presence of a co-ordinative Ti-N bond is indicated by the stability of the compounds which is comparable to that of other compounds Cp₂TiR in which Ti and R form a metallocycle as in (6)⁹ and far greater than that of compounds Cp₂TiR with R = alkyl or aryl where Ti is co-ordinatively unsaturated.¹⁰ The new compounds do not react with N₂ even at very low temperatures indicating that in this case Ti is co-ordinatively saturated by internal co-ordination. A related three-membered, internally co-ordinated, titanacycle has recently been described by Fachinetti and Floriani in their work on Cp₂Ti-acyl com-



pounds containing Ti^{IV} .¹¹ X-Ray studies showed that this compound had the structure (7).

We thank Prof. Dr. F. Jelinek and Dr. H. J. de Liefde Meijer for useful discussions. The investigation was supported by the Netherlands Foundation for Chemical

Research (SON) with financial aid from the Netherlands Organization for the Advancement of Pure Research (ZWO).

(Received, 6th April 1978; Com. 364.)

¹ M. I. Bruce, B. L. Goodall, and I. Matsuda, *Austral. J. Chem.*, 1975, **28**, 1259.

² A. Kasahara, *Bull. Chem. Soc. Japan*, 1968, **41**, 1272.

³ R. J. Foot and B. T. Heaton, *J.C.S. Chem. Comm.*, 1973, 838.

⁴ G. E. Hartwell, R. V. Lawrence, and M. J. Smas, *J.C.S. Chem. Comm.*, 1970, 912.

⁵ M. Nonoyama, *J. Organometallic Chem.*, 1974, **74**, 115.

⁶ F. W. van der Weij and J. H. Teuben, *J. Organometallic Chem.*, 1975, **105**, 203; E. J. M. de Boer and J. H. Teuben, *ibid.*, in the press.

⁷ J. H. Teuben, *J. Organometallic Chem.*, 1973, **57**, 159.

⁸ M. I. Bruce, M. Z. Iqbal, and F. G. A. Stone, *J. Chem. Soc. (A)*, 1970, 3204.

⁹ D. Ytsma, J. G. Hartsuiker, and J. H. Teuben, *J. Organometallic Chem.*, 1974, **74**, 239.

¹⁰ J. H. Teuben, *J. Organometallic Chem.*, 1974, **69**, 241.

¹¹ G. Fachinetti, C. Floriani, and H. Stoeckli-Evans, *J.C.S. Dalton*, 1977, 2297.