

## On the Proposed Intermediacy of $\beta$ -Hydroxyvaline- and Thiazepinone-containing Peptides in Penicillin Biosynthesis


By GULAM BAHADUR, JACK E. BALDWIN,\* TERENCE WAN, and MANKIL JUNG  
(Dyson Perrins Laboratory, Oxford University, South Parks Road, Oxford OX1 3QY)

and SIR EDWARD P. ABRAHAM, JOYCE A. HUDDLESTON, and ROBERT L. WHITE  
(Sir William Dunn School of Pathology, Oxford University, South Parks Road, Oxford OX1 3RE)

**Summary** The two thiazepinone-containing peptides (5) and (7), as well as the  $\beta$ -hydroxyvaline peptide (3), have been synthesised and tested as substrates for isopenicillin N synthesis in a cell-free extract from *Cephalosporium acremonium*; none of these compounds behaved as substrates, in contradiction to a recent proposal.

RECENTLY it has been proposed<sup>1</sup> that the established conversion<sup>2</sup> of the tripeptide (1) into isopenicillin N (2) might proceed by way of intermediates (3), (4), and (5), a process suggested to be in accord with the established stereochemical course of the conversion of (1) into (2), *i.e.* overall retention of configuration at carbons 2 and 5 in (2),<sup>3</sup> Scheme 1. We have synthesised the proposed intermediates (3) and (5) and have tested them as putative precursors in a highly active cell-free synthetase system from *Cephalosporium acremonium* for the conversion of (1) into (2).


Thus D- $\beta$ -hydroxyvaline ( $[\alpha]_D^{20} - 13.3^\circ$ , *c* 1, 5N-HCl), prepared and resolved by known methods,<sup>4</sup> was converted into its benzyl ester through esterification of its *N*-benzyloxy-carbonyl derivative (Cs salt, benzyl bromide, dimethyl-



SCHEME 1. R = L- $\alpha$ -amino- $\delta$ -adipyl.

formamide) and subsequent deprotection with hydrogen bromide in acetic acid.† Reduction ( $\text{Na}, \text{NH}_3$ ) restored the amino-acid with no change in optical activity. This ester was coupled (dicyclohexylcarbodi-imide, 2-hydroxybenzotriazole) with *N*-benzyloxycarbonyl-*S*-benzyl-*L*-cysteine to afford the dipeptide (m.p. 93 °C; 90%), which was then deprotected (40% HBr, glacial AcOH), and finally coupled with 1-benzyl-*N*-benzyloxycarbonyl-*L*- $\alpha$ -aminoadipic acid to give the tripeptide (m.p. 122–124 °C; 12%), which was deprotected ( $\text{Na}, \text{NH}_3$ ) and purified with Hopkin's reagent ( $\text{Hg}^{2+}$ - $\text{H}_2\text{SO}_4$ ) to yield the tripeptide (3).‡

The nucleus of (5), *i.e.* the cyclic peptide (6,  $\text{R} = \text{PhCH}_2\text{O}_2\text{C}$ ), was synthesized by a known procedure.<sup>5</sup> After esterification to give the benzyl ester (triethylamine, benzyl bromide; 25 °C; m.p. 126.5–127.5 °C; 86%) and deprotection (HBr, glacial AcOH, 94%) the free amine (6;  $\text{R} = \text{H}$ ), m.p. 128.5–129 °C, was liberated ( $\text{NaHCO}_3$ ; 96%). This substance was acylated (*N*-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline; tetrahydrofuran; 25 °C; 89%) with 1-benzyl-*N*-benzyloxycarbonyl-*L*- $\delta$ -aminoadipic acid to afford the fully protected cyclic peptide which was deprotected ( $\text{H}_2$ ; 10% Pd/C; MeOH, 25 °C) to give the cyclic peptide (6;  $\text{R} = \text{L-}\alpha\text{-amino-}\delta\text{-adipyl}$ ), m.p. 194 °C (decomp.) (67%). Oxidation (*m*-chloroperbenzoic acid;  $\text{CH}_2\text{Cl}_2$ ; 98%) of the fully protected cyclic peptide gave the (*R*)-sulphoxide which was deprotected, as before, to give the peptide (5) (78%). The (*R*) configuration of the sulphoxide has been previously assigned and corresponds to that implicated in the proposed biosynthetic scheme.¶ The cyclic peptide (6) has been discussed previously as a possible intermediate in penicillin biosynthesis<sup>6</sup> but it was not at that time realised that the aminoadipic acid moiety was essential for penicillin biosynthesis and, furthermore, no cell-free extract with synthetic activity was then available. Consequently, the earlier experiments<sup>6</sup> are invalid. We therefore decided also to test (7) [ $\text{R} = \text{L-}\alpha\text{-amino-}\delta\text{-adipyl}$ ] in a cell-free system.



SCHEME 2

An active cell-free extract containing all co-factors<sup>7</sup> was prepared from *C. acremonium* as previously described. Solutions of the substances (3), (5), and (7) respectively (10  $\mu\text{l}$ ; 40 mM) were each added to the complete system (390  $\mu\text{l}$ ),<sup>7</sup> the mixtures incubated at 27 °C (60 min; shaken at 100 cycles/min), and the reactions stopped by addition of acetone (930  $\mu\text{l}$ ). After centrifugation the supernatants were concentrated in a stream of air to their original volume (400  $\mu\text{l}$ ). All bioassays and chromatographic separations were conducted on these solutions. The controls contained either the tripeptide substrate (1) or no peptide, the latter showing no background level of antibiotic content. The results are presented in the Table. There is no indication

TABLE. Production of antibiotic activity on incubation of substrate (1) and compounds (3), (5), and (7) with the extract<sup>7</sup> from *C. acremonium*.

Compound <sup>a</sup>	Conversion (%) <sup>b</sup>	Product
(1)	77	(2)
(3)	0	—
(5)	0	—
(7)	0	—

<sup>a</sup> Concentration *ca.* 1 mM. <sup>b</sup> Assay against *Staphylococcus aureus* (WCTC-6571) and *Salmonella typhi* (strain Mrs. S). Detection limit *ca.* 2% of (2).

of conversion of compounds (3), (5), or (7) into antibiotics. The incubation mixtures from (3), (5), and (7) were subjected to electrophoresis (paper; pH 6.5) before and after treatment with  $\beta$ -lactamase I (*Bacillus cereus*, 569/H/9) and showed no production of isopenicillin N from (3), (5), or (7), but essentially contained unchanged substrates. Furthermore, chromatography on paper (descending;  $\text{Bu}^n\text{OH-HOAc-H}_2\text{O}$ , 4:1:4) showed only unchanged starting material before or after  $\beta$ -lactamase treatment. When (5) and (7) (1.1 mM) were first incubated in the extract for 15 min there was no detectable inhibition of the synthesis of (2), after the subsequent addition of (1) (0.6 mM). However (3), under these conditions, appeared to behave as a weak inhibitor (*ca.* 9% of normal conversion).

We conclude that none of the proposed intermediates (3), (5), or (7) are acceptable as substrates for the penicillin synthetase system from *C. acremonium*.

(Received, 14th August 1981; Com. 997.)

† It was not possible to benzylate (benzene, benzyl alcohol, toluene-*p*-sulphonic acid) this amino-acid directly.

‡ Homogeneous by t.l.c. ( $\text{SiO}_2$ , 75% aq. PrOH) and electrophoresis on paper (pH 4.5).

§  $\alpha$ -Amino- $\delta$ -adipyl = 5-amino-5-carboxypentanoyl.

¶ The previous study<sup>1</sup> showed that oxidation of the fully protected thiazepinone peptide under these conditions gave only the *R*-sulphoxide. The new data reported were in agreement with those for our compound, both in chemical shift and coupling constants.

<sup>1</sup> S. Wolfe, R. J. Bowers, S. K. Hasan, and P. M. Zazmaier, *Can. J. Chem.* 1981, **59**, 406.

<sup>2</sup> J. O'Sullivan, R. C. Bleaney, J. A. Huddleston, and E. P. Abraham, *Biochem. J.*, 1979, **184**, 421; T. Konomi, S. Herchen, J. E. Baldwin, M. Yoshida, N. A. Hunt, and A. L. De Main, *ibid.*, p. 427.

<sup>3</sup> Reviewed by D. J. Aberhart, *Tetrahedron*, 1977, **33**, 1545.

<sup>4</sup> H. C. Beyerman, L. Maat, D. de Rijke, and J. P. Visser, *Rec. Trav. Chim. Pays-Bas*, 1967, **86**, 1057 and references cited therein; G. W. Edwards and L. M. L. Minthorn, *Can. J. Biochem.*, 1968, **46**, 1227.

<sup>5</sup> N. J. Leonard and R. Y. Ning, *J. Org. Chem.*, 1966, **31**, 3928.

<sup>6</sup> H. R. V. Arnstein and M. E. Clubb, *Biochem. J.*, 1958, **68**, 528; B. Sjöberg, H. Thelin, L. Nathorst Westfelt, E. E. van Tamelen, and E. R. Wagner, *Tetrahedron Lett.*, 1965, 281; N. J. Leonard, and G. E. Wilson, Jr., *J. Am. Chem. Soc.*, 1964, **86**, 5307.

<sup>7</sup> E. P. Abraham, J. A. Huddleston, G. S. Jayatilake, J. O'Sullivan, and R. L. White in 'Recent Advances in the Chemistry of  $\beta$ -Lactam Antibiotics,' (Proceedings of the 2nd International Symposium), ed. G. I. Gregory, Royal Society of Chemistry, 1981, p. 125.