


## Synthetic Reactions in Polyethylene Glycol. Diazotization and Sandmeyer Reactions of Anilines in Polyethylene Glycol–Methylene Dichloride

Nobutaka Suzuki,\*<sup>a</sup> Yoshihiro Kaneko,<sup>a</sup> Tateo Nomoto,<sup>b</sup> and Yasuji Izawa<sup>a</sup>

<sup>a</sup> Department of Industrial Chemistry, Faculty of Engineering, and <sup>b</sup> Department of Chemistry, Faculty of Education, Mie University, Tsu, Mie 514, Japan

Diazotization and the Sandmeyer reactions of representative arylamines proceed smoothly in polyethylene glycol–CH<sub>2</sub>Cl<sub>2</sub> and have been found to be effective for the preparation of aryl halides and cyanides.

Chemistry in cyclic polyethers (crown ethers) has attracted widespread interest because of the rather high selectivity of many reactions of activated anion species. The properties of polyethylene glycol (PEG) make it an excellent substitute for crown ethers in many reactions,<sup>1,2</sup> and its use as a co-solvent has been reported. We have now found that diazotization of arylamines and Sandmeyer substitution reactions of the resulting diazonium compounds take place with NaNO<sub>2</sub>–HCl in PEG–CH<sub>2</sub>Cl<sub>2</sub><sup>3,4</sup> and with Cu<sub>2</sub>X<sub>2</sub>, Cu–MX, or K<sub>2</sub>[Cu(CN)<sub>4</sub>·NH<sub>3</sub>] in PEG–CH<sub>2</sub>Cl<sub>2</sub> systems, respectively, providing the first example of the cyano substitution of diazonium salts using the isolated solid K<sub>2</sub>[Cu(CN)<sub>4</sub>·NH<sub>3</sub>] complex (Scheme 1).



a; Ar = Ph  
 b; Ar = *p*-PhCOC<sub>6</sub>H<sub>4</sub>  
 c; Ar =  $\alpha$ -naphthyl  
 d; Ar = *p*-ClC<sub>6</sub>H<sub>4</sub>  
 e; Ar = *p*-MeOC<sub>6</sub>H<sub>4</sub>  
 f; Ar = *o*-MeOC<sub>6</sub>H<sub>4</sub>  
 g; Ar = *m*-MeOC<sub>6</sub>H<sub>4</sub>  
 h; Ar = *p*-PhC $\equiv$ CC<sub>6</sub>H<sub>4</sub>

X = Cl, Br, I, or CN      Y = Cl, Br, or I

**Scheme 1.** Reagents: i, NaNO<sub>2</sub>–HY in PEG–CH<sub>2</sub>Cl<sub>2</sub> at 0 °C; ii, MX in PEG–CH<sub>2</sub>Cl<sub>2</sub> at 0 °C to room temperature.

The arylamine (1) (0.1 mmol; 10 mmol dm<sup>-3</sup>) mixed with the acid HY (0.1 cm<sup>3</sup> of 1 mol dm<sup>-3</sup>; 0.1 mmol) was diazotized by addition of NaNO<sub>2</sub> (0.1 mmol) dissolved in PEG 200 (2 g; 1 mol dm<sup>-3</sup>)–CH<sub>2</sub>Cl<sub>2</sub> at 0 °C. At 0 °C the resulting diazotized solution was added to the solid reagent MX (0.1 mmol to slight excess), and the mixture set aside at room temperature. Samples were withdrawn at intervals and, for the reactions involving PEG, washed with 1M–NaOH. The products were identified by g.l.c. comparison with authentic samples and also by isolation<sup>‡</sup> (i.r. spectroscopy). For comparison, some experiments in aqueous solution were carried out without the PEG–CH<sub>2</sub>Cl<sub>2</sub> system. Table 1 shows the results.

On diazotization in PEG–CH<sub>2</sub>Cl<sub>2</sub>, the starting materials (1) were consumed completely in 20 min. Three equivalents of the acid were necessary.

Aniline (1a) was chlorinated quantitatively in PEG–CH<sub>2</sub>Cl<sub>2</sub>, whereas it did not give chlorobenzene in aqueous solution under similar conditions (Cu<sub>2</sub>Cl<sub>2</sub> and/or Cu powder).<sup>5</sup> Chlorination of (1b)–(1d) was accompanied by reduction to give ArH (4).<sup>6</sup> Tetraglyme suppressed significantly the side reactions to give (4), when it was employed instead of PEG. Bromination proceeded in good yields with HBr, but yields were lower than in the chlorinations with HCl. The extent of iodination exceeded the consumption of the starting materials; we assume that this resulted from the

<sup>†</sup> Final concentration in a 10 ml solution of PEG (2 g)–CH<sub>2</sub>Cl<sub>2</sub>–water (0.1 ml).

<sup>‡</sup> For isolation 100 ml of the solutions (the same concentration) were employed.

**Table 1.** Diazotization and Sandmeyer reactions in PEG-CH<sub>2</sub>Cl<sub>2</sub>.

Aniline (1) (10 mm)	HY <sup>a</sup>	Consumption of (1) (%)	MX <sup>b</sup>	% Yield of products (%) <sup>c</sup>		
				Ar-X (3)	Ar-H (4)	Ar-Cl
<b>Chlorination:</b>						
(1a)	3M A	96	D	100 [93] <sup>h</sup>	—	—
(1a) <sup>d</sup>	3M A	96	D (or E)	1 (17)	—	—
(1a) <sup>d,e</sup>	conc. A <sup>f</sup>	100	D (or E)	70 (74)	—	—
(1b)	3M A	100	D	19	78	—
(1b)	5M A	100	D	20	66	—
(1c)	3M A	100	D	7	25	—
(1d)	3M A	99	D	35	56	—
<b>Bromination:</b>						
(1a)	3M A	96	F	11	—	43
(1a)	4M B	97	F	92 [86] <sup>h</sup>	—	—
(1e)	4M B	100	F	32	14	—
<b>Iodination:</b>						
(1a)	3M A	99	G	57	—	2
(1a)	3M C	45	G	107	—	—
(1a)	3M C <sup>g</sup>	56	G	163 [85] <sup>h</sup>	—	—
<b>Cyanation:</b>						
(1a)	3M A	88	H	15	—	75
(1a)	3M A	100	J/D; J/K; or J/E	0	—	40
(1a)	3M A	95	L	11	—	19
(1a)	3M A	100	M	67 [62] <sup>h</sup>	—	16
(1c)	3M A	100	M	46 [40] <sup>i</sup>	32	1
(1d)	3M A	100	M	51 [48] <sup>i</sup>	25	20
(1f)	3M A	97	M	93 [88] <sup>i</sup>	7	—
(1g)	3M A	100	M	39 [30] <sup>i</sup>	12	—
(1h)	3M A	99	M	38 [35] <sup>j</sup>	53 [48] <sup>i</sup>	—

<sup>a</sup> A: HCl; B: HBr; C: HI (0.1 ml). <sup>b</sup> D: Cu<sub>2</sub>Cl<sub>2</sub>; E: copper powder; F: Cu<sub>2</sub>Br<sub>2</sub>; G: KI; H: Cu<sub>2</sub>(CN)<sub>2</sub>; J: KCN; K: Ni(NO<sub>3</sub>)<sub>2</sub>·6H<sub>2</sub>O; L: 0.39 M K<sub>2</sub>[Cu(CN)<sub>4</sub>·NH<sub>3</sub>]; M: solid K<sub>2</sub>[Cu(CN)<sub>4</sub>·NH<sub>3</sub>]. <sup>c</sup> Based on the consumption of (1). <sup>d</sup> In aq. soln. <sup>e</sup> 333 mm. <sup>f</sup> 2.5 ml. <sup>g</sup> 0.2 ml. <sup>h</sup> Isolated yield based on the starting (1). <sup>i</sup> Isolated yield after t.l.c.

known iodination of the benzene present in the PEG § by I<sub>2</sub>-nitric acid<sup>7</sup> produced *in situ* from HI-nitrous acid (the HI used was the commercial product which showed a colour due to iodine).

Cyanation in PEG-CH<sub>2</sub>Cl<sub>2</sub> did not proceed under the normal reaction conditions examined.<sup>8,9</sup> Only chlorination or bromination products were obtained depending on the conditions. High yields of cyanation products were obtained when excess of the dry K<sub>2</sub>[Cu(CN)<sub>4</sub>·NH<sub>3</sub>] complex<sup>10</sup> (not isolated; K<sub>2</sub>SO<sub>4</sub> also present) was employed as the cyanation reagent.

Received, 17th May 1984; Com. 689

## References

- N. Suzuki, Y. Ayaguchi, and Y. Izawa, *Bull. Chem. Soc. Jpn.*, 1982, **55**, 3349; N. Suzuki, Y. Ayaguchi, T. Tsukanaka, and Y. Izawa, *ibid.*, 1983, **56**, 353; N. Suzuki, Y. Ayaguchi, K. Shimazu, T. Ito, and Y. Izawa, in 'Polymer Science and Technology,' vol. 24, 'Crown Ethers and Phase Transfer Catalysis in Polymer Science,' eds. C. E. Carraher and L. Mathias, 1984, Plenum Press, New York, pp. 385-96; N. Suzuki, T. Tsukanaka, T. Nomoto, Y. Ayaguchi, and Y. Izawa, *J. Chem. Soc., Chem. Commun.*, 1983, 515.

- E. Santaniello, A. Manzocchi, and P. Sozzani, *Tetrahedron Lett.*, 1979, 4581; E. Santaniello, P. Terraboschi, and P. Sozzani, *J. Org. Chem.*, 1981, **46**, 4584; E. Santaniello, A. Fiecchi, A. Manzocchi, and P. Ferraboschi, *ibid.*, 1983, **48**, 3074; K. Sukata, *J. Synth. Org. Chem. Jpn.*, 1981, **39**, 443.
- For leading reviews on synthetic applications of diazonium compounds, see (a) N. Kornblum, *Org. React.*, 1944, **2**, 262; (b) D. S. Wulfman, in 'The Chemistry of Diazonium and Diazo Groups,' Part 1, ed. S. Patai, 1978, Wiley, New York, p. 247.
- Diazonium compounds prepared in organic solvents: in xylene (yield, 70%), T. L. Fletcher and M. J. Namkung, *Chem. Ind. (London)*, 1961, 179; in Me<sub>2</sub>SO (yield, 50-70% for chlorination; 55-97% for cyanation), M. Kobayashi, E. Yamada, M. Matsui, and N. Kobori, *Org. Prep. Proc. Int.*, 1969, **1**, 221; M. Kobayashi, H. Minato, E. Yamada, and N. Kobori, *Bull. Chem. Soc. Jpn.*, 1970, **43**, 215; in CH<sub>2</sub>Cl<sub>2</sub> (75%), C. G. Swain and R. J. Rogers, *J. Am. Chem. Soc.*, 1975, **97**, 779; in 18-crown-6-CHCl<sub>3</sub> (82%), S. H. Korzeniowski and G. W. Gokel, *Tetrahedron Lett.*, 1977, 3519; J. R. Beadle and G. W. Gokel, *ibid.*, 1984, **25**, 1681.
- L. Gattermann, *Ber.*, 1890, **23**, 1218.
- This reduction is carried out in EtOH classically or with hypophosphorous acid: see ref. 3(b), p. 286; S. Yasui, K. Nakamura, and A. Ohno, *J. Org. Chem.*, 1984, **49**, 878.
- F. B. Dains and R. Q. Brewster, *Org. Synth.*, Coll. Vol. I, 1956, 323.
- J. A. Mcrae, *J. Am. Chem. Soc.*, 1930, **52**, 4550; L. A. Bigelow, *Org. Synth.*, Coll. Vol. I, 1956, 135.
- K. Friedrich and K. Wallanfels, in 'The Chemistry of the Cyano Group,' ed. Z. Rappoport, 1970, Interscience, New York, p. 67.
- H. Hagenest and F. W. Stauf, US. Pat., 1 879 209; *Chem. Abstr.*, 1933, **27**, 997; US. Pat., 1 962 559; *Chem. Abstr.*, 1934, **28**, 4848; ref. 9, p. 87.

§ The PEG 200 used was dried azeotropically with benzene. A series of blank tests in the absence of aniline confirmed that only the iodination gave such a halogenation product under otherwise identical conditions; the other halogenations did not give such a product.