



ether are allowed to warm from  $-78\text{ }^{\circ}\text{C}$  to room temperature, the solution turns brown and  $\text{EtOCH}(\text{Me})\text{CH}_2(p\text{-C}_6\text{H}_4\text{R})$  (**2**) is formed in high yield ( $>90\%$ ) (Scheme 1). At ambient temperature the reaction is complete within *ca.* 2–3 min. The compounds (**2**) which correspond to the product of the insertion of the benzylidene ligand into the  $\alpha\text{-C-H}$  bond of diethyl ether were identified by g.c.,  $^1\text{H}$  n.m.r., and mass spectroscopy and, in the case of (**2a**), additionally by comparison with an authentic sample.<sup>8</sup> Besides (**2**), small amounts of the *cis*- and *trans*-stilbenes and *p*- $\text{MeC}_6\text{H}_4\text{R}$  as well as  $(\text{CO})_6\text{W}$  and small amounts of the  $\mu$ -benzylidene complexes  $[(\text{CO})_5\text{W}]_2[\mu\text{-CH}(p\text{-C}_6\text{H}_4\text{R})]$ <sup>7</sup> were obtained. Neither the  $\beta\text{-C-H}$  insertion product nor benzyl ethyl ether (which constitutes one of the major products when phenyldiazomethane is photolysed in diethyl ether at  $0\text{ }^{\circ}\text{C}$ <sup>9</sup>) and *p*-methylbenzyl ethyl ether, respectively, could be observed. The formation of more than 0.2% of each compound would have been detected.

The analogous reaction of (**1**) with tetrahydrofuran (THF) yields  $\alpha$ -benzyltetrahydrofuran and  $\alpha$ -(*p*-methylbenzyl)tetrahydrofuran ( $>90\%$ ), respectively. The  $(\text{CO})_5\text{W}$ -fragment is stabilized by THF giving  $(\text{CO})_5\text{W}(\text{THF})$  (Scheme 1). Again, the  $\beta\text{-C-H}$  insertion product could not be detected.

The kinetics of the reaction of (**1b**) with THF in THF-octane at  $15.1\text{ }^{\circ}\text{C}$  were studied by following the disappearance of the  $\nu(\text{CO})\text{-A}_1$ -absorption of (**1b**) at  $2072\text{ cm}^{-1}$  in the i.r. spectrum. Pseudo-first order conditions  $\{[(\mathbf{1b})]_0 : [\text{THF}]_0 \text{ ranging from } 1 : 307 \text{ to } 1 : 1500\}$  were employed. The rate constants were reproducible to at least 7%. The reaction follows an additive rate law:  $-\text{d}[(\mathbf{1b})]/\text{d}t = k_1[(\mathbf{1b})] + k_2[(\mathbf{1b})][\text{THF}]^2$ . At  $15.1\text{ }^{\circ}\text{C}$ ,  $k_1 = 1.6 \times 10^{-4}\text{ s}^{-1}$  and  $k_2 = 4.6 \times 10^{-4}\text{ dm}^6\text{ mol}^{-2}\text{ s}^{-1}$  {from a plot of  $k_{\text{obs}}$  vs.  $[\text{THF}]^2$ ; seven pairs of  $k_{\text{obs}}$  and  $[\text{THF}]^2$ , correlation coefficient better than 0.996}. When THF was replaced by  $[\text{H}_8]\text{THF}$ , an isotope effect of  $k_2^{\text{H}}/k_2^{\text{D}} = 1.7$  was observed.

The first-order term coincides with the independently

determined rate of thermolysis of (**1b**) in octane ( $k_{\text{obs}} = 1.64 \times 10^{-4}\text{ s}^{-1}$ , at  $15.1\text{ }^{\circ}\text{C}$ ) giving rise to the formation of the *cis*- and *trans*-stilbenes, *p*- $\text{C}_6\text{H}_4\text{Me}_2$ , and  $[(\text{CO})_5\text{W}]_2[\mu\text{-CH}(p\text{-C}_6\text{H}_4\text{Me})]$  which are the characteristic thermolysis products of (**1b**) in inert solvents.<sup>7</sup>  $\alpha$ -(*p*-Methylbenzyl)tetrahydrofuran is produced by an associative mechanism (third-order term) thus accounting for the regio-specificity of the insertion. This reaction is initiated by a nucleophilic attack of THF at the strongly electrophilic benzylidene carbon of (**1b**). The yield of (**3b**) drops drastically with decreasing concentration of THF, in agreement with the second-order THF term in the rate expression.

When similar conditions are employed, (**1a**) reacts with THF *ca.* three times faster than (**1b**).  $\text{Et}_2\text{O-THF}$  competition experiments show that the insertion of ' $\text{CH}(p\text{-C}_6\text{H}_4\text{Me})$ ' into the  $\alpha\text{-C-H}$  bond of THF is favoured by a factor of ten.

We thank the Deutsche Forschungsgemeinschaft for financial support of this work.

Received, 31st December 1984; Com. 1794

## References

- 1 J. A. Connor and P. D. Rose, *J. Organomet. Chem.*, 1970, **24**, C45; E. O. Fischer and K. H. Dötz, *ibid.*, 1972, **36**, C4.
- 2 J. A. Connor, P. D. Rose, and R. M. Turner, *J. Organomet. Chem.*, 1973, **55**, 111.
- 3 F. R. Kreissl, E. O. Fischer, C. G. Kreiter, and H. Fischer, *Chem. Ber.*, 1973, **106**, 1262.
- 4 U. Schubert and E. O. Fischer, *Chem. Ber.*, 1973, **106**, 3882.
- 5 E. O. Fischer and V. Kiener, *Angew. Chem.*, 1967, **79**, 982; *Angew. Chem., Int. Ed. Engl.*, 1967, **6**, 961.
- 6 E. O. Fischer and G. Kreis, *Chem. Ber.*, 1973, **106**, 2310.
- 7 H. Fischer, S. Zeuner, and K. Ackermann, *J. Chem. Soc., Chem. Commun.*, 1984, 684.
- 8 A. E. Tschitschibabin and S. A. Jelgasin, *Ber. Dtsch. Chem. Ges.*, 1914, **47**, 1843.
- 9 H. Tomioka, *J. Am. Chem. Soc.*, 1979, **101**, 256.