

Synthesis and Chemical Modification of Homoseryl Peptides†

Kleomenis Barlos,*^a Petros Mamos,^b Dionysios Papaioannou,^a Chariklia Sanida,^a and Christos Antonopoulos^a

^a Department of Chemistry, and ^b Department of Medicine, University of Patras, Patras, Greece

The readily synthesised *N,O*-ditritylhomoserine (**4**) was used for the efficient incorporation of homoserine (**1**) into peptides; the derived homoseryl peptides were transformed into peptides of canaline and 1,4-diaminobutyric acid using the Mitsunobu reaction.


Homoserine, Hse, (**1**) is a naturally occurring amino acid with important biological properties.¹ Because of the strong tendency of its unprotected hydroxy derivatives towards lactonisation,² few homoseryl peptides have been synthesised³ to date. Incorporation of (**1**) into peptides is of interest because its side chain hydroxy function could be transformed into derivative groups, thus providing a number of modified peptides useful in structure-activity related investigations. However realisation of this projected application of (**1**), characterising the so-called pluripotential amino acids,⁴ requires derivatives of (**1**) with extremely labile hydroxy protecting groups. We report here on the simple synthesis of

one such derivative, namely (**4**), which allows for efficient incorporation of (**1**) into peptides, and provide first examples of the use of (**1**) as a pluripotential amino acid.

Treatment of the readily available *N*-tritylhomoserine (**2**)⁵ with a slight excess of trityl chloride and triethylamine in CHCl₃ gave (**4**), *via* an intramolecular trityl group migration which occurs rapidly on the initially formed *N*-tritylhomoserine trityl ester (**3**). Compound (**4**), isolated as its corresponding diethylammonium salt {m.p. 172–174 °C, [α]_D²⁵ +18.6 (*c* 2, MeOH)} in 74% yield, was further treated with dicyclohexylcarbodiimide (DCC) and 1-hydroxybenzotriazole (HOBt)⁶ to provide the active ester (**5**) {oil, [α]_D²⁵ +47.5 (*c* 3, CHCl₃)} in 93% yield. Coupling⁷ of (**5**) with leucinamide and glycineamide yielded the dipeptides (**6**) {m.p. 211–212 °C, [α]_D²⁵ –15.7 (*c* 1, CHCl₃)} and (**7**) {m.p. 194–196 °C, [α]_D²⁵ –39.7 (*c* 1, CHCl₃)} in 75% and 87% yield respectively.

Treatment of (**6**) or (**7**) with 25% CF₃CO₂H in CH₂Cl₂ for 5

† All optically active amino acids and derivatives referred to in this communication are of the *L* configuration. New compounds gave analytical and spectral data in agreement with the proposed structures.



Scheme 1. Reagents: i, TrtCl , NEt_3 ; ii, HOBt , DCC ; iii, Gly-NH_2 , NEt_3 ; iv, 1.5% $\text{CF}_3\text{CO}_2\text{H}$ in EtOH-CHCl_3 (75:25); v, Trt-Phe-OBt , NEt_3 ; vi, PPh_3 , diethylazodicarboxylate, *N*-hydroxyphthalimide; vii, PPh_3 , diethylazodicarboxylate, phthalimide.

min at 0°C , in an attempt to remove both trityl groups, resulted in fragmentation of the dipeptides to homoserine lactone and the corresponding amino acid amides. In contrast, complete deprotection without fragmentation occurs when (6) or (7) is treated with 1.5% $\text{CF}_3\text{CO}_2\text{H}$ in EtOH-CHCl_3 (75:25) for 5 h at room temperature, and the reaction mixture is concentrated *in vacuo*. The trifluoroacetate salts of (8) {powder, m.p. 72°C (decomp.), $[\alpha]_{\text{D}}^{25} +11.4$ (*c* 0.6, MeOH)} and (9) {powder, $[\alpha]_{\text{D}}^{25} -2.1$ (*c* 1, MeOH)} were thus isolated in 93% and 85% yield respectively.

Coupling of (8) with *N*-tritylphenylalanine 1-hydroxybenzotriazolyl ester⁶ gave the homoserine tripeptide (10) {m.p. 112°C (decomp.), $[\alpha]_{\text{D}}^{25} -6.9$ (*c* 1, MeOH)} in 72% yield. Condensation of (10) with *N*-hydroxyphthalimide⁸ or phthalimide⁹ using triphenylphosphine and diethylazodicarboxylate¹⁰ gave the protected canaly tripeptide (11) {m.p. $175-177^\circ\text{C}$, $[\alpha]_{\text{D}}^{25} -11.8$ (*c* 1, CHCl_3)} and the 1,4-diaminobutryl peptide (12) {m.p. 125°C , $[\alpha]_{\text{D}}^{25} +10.5$ (*c* 2, CHCl_3)}, in 40% and 58% yield respectively after silica gel column chromatography. Canaline (13) is a structural analogue of ornithine with interesting biological properties.¹¹ Using a similar procedure to that described herein, derivatives of (13), suitably protected for peptide synthesis, have also been prepared.¹² Selective removal of the phthalyl-protected group can be readily achieved on treating (11) with a mixture of acetic acid and hydrazine hydrate (1:1) in methanol at 45°C for 4 h. The side chain-unprotected tripeptide (14) thus obtained in 68% yield {m.p. $198-199^\circ\text{C}$, $[\alpha]_{\text{D}}^{25} -4.5$ (*c* 1, MeOH)} can be further detriylated with a 5 molar excess of HCl in diethyl ether to afford the dihydrochloride of the canaline containing tripeptide (15) {m.p. 190°C (decomp.)},

$[\alpha]_{\text{D}}^{25} -0.7$ (*c* 0.8, MeOH)} in 72% yield. To our knowledge, (15) represents the first synthetic peptide of canaline, (13), reported to date.

Received, 18th April 1986; Com. 512

References

- 1 A. Lehninger, 'Biochemistry,' 2nd edn., Worth Publishers Inc., New York, 1975.
- 2 M. D. Armstrong, *J. Am. Chem. Soc.*, 1949, **71**, 3399; J. P. Greenstein and M. Winitz, 'Chemistry of the Amino Acids,' John Wiley and Sons, New York, vol. III, p. 2612; M. Flavin and C. Slaughter, *Biochemistry*, 1965, **4**, 1370.
- 3 M. Manning, A. Turan, J. Haldar, and W. Sawyer, in 'Proceedings of the Fifth American Peptide Symposium,' eds. M. Goodman and J. Mienhofer, John Wiley and Sons, New York, 1977, p. 201; A. Buku, R. Altman, and T. Wieland, *Liebigs Ann. Chem.*, 1976, 417.
- 4 D. Roberts, K. Suba, J. Samanen, and D. S. Kemp, *Tetrahedron Lett.*, 1980, **21**, 3435.
- 5 K. Barlos and D. Theodoropoulos, *Z. Naturforsch., Teil B.*, 1982, **37**, 886.
- 6 K. Barlos, D. Papaioannou, and D. Theodoropoulos, *Int. J. Pept. Protein Res.*, 1984, **23**, 300.
- 7 K. Barlos, D. Papaioannou, and C. Sanida, *Liebigs Ann. Chem.*, 1984, 1308.
- 8 E. Grochowski and J. Jurczak, *Synthesis*, 1977, 277.
- 9 O. Mitsunobu, M. Wada, and T. Sano, *J. Am. Chem. Soc.*, 1972, **94**, 679.
- 10 O. Mitsunobu, *Synthesis*, 1981, 1.
- 11 G. A. Rosenthal, *Life Sci.*, 1978, **23**, 93.
- 12 K. Barlos, D. Papaioannou, and C. Sanida, *Liebigs Ann. Chem.*, 1986, 287.