

## Biosynthesis of the Insect Antifeedant Steroid Nic-1: Origins of the Aromatic Ring-D

Harjit K. Gill, Roland W. Smith, and Donald A. Whiting\*


Department of Chemistry, The University, Nottingham NG7 2RD, U.K.

Isotope administration experiments with *Nicandra physaloides* plants using [3'-C<sup>2</sup>H<sub>3</sub>]- and [3'-<sup>14</sup>CH<sub>3</sub>]-mevalonic acid, analysed by <sup>2</sup>H n.m.r. and by degradation, respectively, show that the aromatic ring-D of Nic-1 (**1**) is formed by ring-D expansion in a steroid precursor with oxidative inclusion of the c/D angular methyl.

---


*Nicandra physaloides* (the Peruvian 'shoofly' plant) contains a group of highly oxidised 24-methylsteroids (nicandrenoids)<sup>1</sup> related to the withanolides.<sup>2</sup> The major metabolite Nic-1 (**1**) exhibits antifeedant properties towards the tobacco hornworm. A very unusual feature of the structure of Nic-1 is the

aromatic D-ring, carrying a side chain displaced from its customary site in 'normal' steroids *e.g.* Nic-3 (**2**). The co-occurrence of Nic-1 and Nic-3 suggests late stage D-aromatization. Two plausible biosynthetic hypotheses may be entertained (Scheme 1): in path (a) the c/D angular methyl is



oxidised and incorporated into ring-D with subsequent aromatisation, while in path (b) oxidative elision of C-18 is envisaged (as, e.g. the removal of the C-14 methyl in lanosterol) followed by insertion of a C<sub>1</sub> unit from *S*-adenosylmethionine (SAM), and aromatisation. We report here experiments which distinguish these pathways and establish the origin of the aromatic ring. We thus prepared [3'-<sup>14</sup>CH<sub>3</sub>]-mevalonic acid (MVA), using a scaled-down version of the Macmillan-Scott procedure developed for [3'-<sup>13</sup>CH<sub>3</sub>]-MVA,<sup>3</sup> and administered a sample to *N. physaloides* plants, seven weeks old, using the wick method. [<sup>14</sup>CH<sub>3</sub>]-SAM was similarly administered. Nic-1 was isolated in each experiment and oxidised (potassium permanganate) to trimellitic acid (3). Table 1 shows the results after recrystallisation to constant activity. It is clear that although [<sup>14</sup>CH<sub>3</sub>]-SAM is incorporated into Nic-1, no activity appears in ring-D, but is presumably located only at C-28. However on incorporation of [3'-<sup>14</sup>CH<sub>3</sub>]-MVA, 26% of the activity appears in the trimellitic acid fragment: since four sites (C-19, C-18, C-21, and C-26 or C-27) in a steroid precursor such as (2) would be expected to be labelled, this experiment is consistent with retention of C-18 as part of the aromatic fragment.

To determine precisely the site in ring-D supplied by the MVA methyl we turned to deuterium labelling. Thus [3'-C<sup>2</sup>H<sub>3</sub>]-MVA was prepared from deuterioacetic acid, using the same procedure as above, and a large sample (408 mg) was applied to 23 plants. After 7 days metabolism, Nic-1 was isolated without carrier. A <sup>2</sup>H n.m.r. spectrum of this sample (160 mg, 12 680 scans, CH<sub>2</sub>Cl<sub>2</sub>-10% CFCl<sub>3</sub>) showed a clear absorption at δ 7.01, (with other expected signals and no <sup>2</sup>H impurities); this signal corresponds to that of the 18-H (s) in the <sup>1</sup>H n.m.r.



Scheme 1. Possible pathways for ring-D aromatisation.

Table 1. Incorporation of <sup>14</sup>C-labelled species into (1) and (3) and subsequent specific activities.

	Absolute incorporation/ %	Specific activity/ d.p.m. mmol <sup>-1</sup>	
		(1) <sup>a</sup>	(3) <sup>b</sup>
[3'- <sup>14</sup> CH <sub>3</sub> ]-MVA	0.10	444 × 10 <sup>3</sup>	116 × 10 <sup>3</sup>
[ <sup>14</sup> CH <sub>3</sub> ]-SAM	0.002	241 × 10 <sup>3</sup>	0

<sup>a</sup> After dilution. <sup>b</sup> As trimethyl ester.

The evidence thus points to a biosynthetic pathway such as (a), Scheme 1, in which C-18 is oxidised and incorporated into ring-D, while retaining one hydrogen, at least in part. A possible mechanism would involve a fused cyclopropane intermediate and would initially parallel the C-19 transformation in the lanosterol-cycloartenol conversion,<sup>4</sup> but this remains to be determined.

Received, 17th June 1986; Com. 838

## References

- M. J. Begley, L. Crombie, P. J. Ham, and D. A. Whiting, *J. Chem. Soc., Perkin Trans. 1*, 1976, 296, 304.
- E. Glotter, I. Kirson, D. Lavie, and A. Abraham in 'Bioorganic Chemistry,' ed. E. E. van Tamelen, Academic Press, New York, 1978, Vol. II, p. 57; I. Kirson and E. Glotter, *J. Natl. Prod.*, 1981, **44**, 633.
- P. Lewer and J. MacMillan, *J. Chem. Soc., Perkin Trans. 1*, 1983, 1417; E. Bardshiri, T. J. Simpson, A. I. Scott, and K. Shishido, *J. Chem. Soc., Perkin Trans. 1*, 1984, 1765.
- E. Caspi, *Tetrahedron*, 1986, **42**, 3, and references therein.