

## Fluoro(nitrile)xenon(II) Cations, $\text{RC}\equiv\text{N}-\text{XeF}^+ \text{AsF}_6^-$ ( $\text{R} = \text{H}, \text{CH}_3, \text{CH}_2\text{F}, \text{C}_2\text{H}_5, \text{C}_2\text{F}_5, \text{C}_3\text{F}_7,$ or $\text{C}_6\text{F}_5$ ); Novel Examples of Xenon–Nitrogen Bonds and $^{129}\text{Xe}-^{13}\text{C}$ , $^{129}\text{Xe}-^1\text{H}$ , and $^{129}\text{Xe}-^{14}\text{N}$ Nuclear Spin–Spin Couplings



Adel A. A. Emara and Gary J. Schrobilgen\*

Department of Chemistry, McMaster University, Hamilton, Ontario L8S 4M1, Canada


A new class of Xe–N bonded compound results from the interaction of the electron lone pair of a nitrile with the Lewis acid,  $\text{XeF}^+$ ; the cationic adducts,  $\text{RC}\equiv\text{N}-\text{XeF}^+$ , have been derived from the interaction of the appropriate nitrile with either  $\text{XeF}^+\text{AsF}_6^-$  or  $\text{Xe}_2\text{F}_3^+\text{AsF}_6^-$  in anhydrous HF at low temperature and characterized in the solid state by Raman spectroscopy (for  $\text{R} = \text{H}$  and  $\text{Me}$ ) and in HF solution by  $^{129}\text{Xe}$ ,  $^{19}\text{F}$ ,  $^{14}\text{N}$ ,  $^{13}\text{C}$ , and  $^1\text{H}$  n.m.r. spectroscopy.

Herein we report a new ligand group for xenon which is bonded to the noble gas atom through nitrogen. In choosing a likely ligand precursor for a Xe–C or Xe–N bond, the very weak protonic acid HCN was considered and initially found to be unreactive towards  $\text{XeF}_2$  in  $\text{SO}_2\text{ClF}$  at room temperature by the usual HF displacement to give the Xe–C and Xe–N bonded compounds,  $\text{FXe}(\text{CN})$  and  $\text{Xe}(\text{CN})_2$ . Although the conventional HF displacement is not a viable route to xenon(II) cyanides and isocyanides, we have found that HCN, and nitriles in general, behave as nitrogen bases towards the  $\text{XeF}^+$  cation.

The reactions of  $\text{XeF}^+\text{AsF}_6^-$  and  $\text{Xe}_2\text{F}_3^+\text{AsF}_6^-$  with HCN,  $\text{CH}_3\text{CN}$ ,  $\text{CH}_2\text{FCN}$ ,  $\text{C}_2\text{H}_5\text{CN}$ ,  $\text{C}_2\text{F}_5\text{CN}$ ,  $\text{C}_3\text{F}_7\text{CN}$ , and  $\text{C}_6\text{F}_5\text{CN}$  were carried out by combining stoichiometric amounts of the reactants in anhydrous HF and warming to  $-20$  to  $-10^\circ\text{C}$  to effect reaction and dissolution in the solvent (HCN reactions were also conducted in  $\text{SO}_2\text{ClF}$  solvent). The reactions proceed according to equations (1) and (2).



Multinuclear magnetic resonance spectra were recorded for the nitrile cations in HF solvent, and in the case of the  $^1\text{H}$  and  $^{129}\text{Xe}$  n.m.r. spectra of the  $\text{HC}\equiv\text{N}-\text{XeF}^+$  cation, in  $\text{BrF}_5$  solvent. As every element in the  $\text{RC}\equiv\text{N}-\text{XeF}^+$  cations studied possesses at least one nuclide which is suitable for observation by n.m.r. spectroscopy, n.m.r. studies using both naturally abundant and  $^{13}\text{C}$  enriched compounds were undertaken and have provided unambiguous proof for the structures of a majority of the cations in solution (Table 1). In the case of  $\text{HC}\equiv\text{N}-\text{XeF}^+$ , the chemical shifts of all five nuclei and eight of the ten possible spin–spin couplings that have been observed are listed in Table 1. The  $^{129}\text{Xe}$ ,  $^{14}\text{N}$ ,  $^{13}\text{C}$ , and  $^1\text{H}$  spectra are illustrated in Figures 1 and 2. The couplings,  $^1J(^{129}\text{Xe}-^{14}\text{N})$ ,  $^2J(^{129}\text{Xe}-^{13}\text{C})$ , and  $^3J(^{129}\text{Xe}-^1\text{H})$ , represent the first examples of nuclear spin–spin couplings observed between these nuclides. Owing to the cylindrical symmetry of the  $\text{C}\equiv\text{N}-\text{Xe}-\text{F}$  moiety in the new cation series, low viscosity of the HF solvent medium, and the small quadrupole moment of  $^{14}\text{N}$ , quadrupole relaxation of the  $^{129}\text{Xe}-^{14}\text{N}$  coupling is found to be minimal giving rise to slightly quadrupole collapsed 1:1:1 triplets in the  $^{129}\text{Xe}$  spectra and  $^{129}\text{Xe}$  satellites in the  $^{14}\text{N}$  spectra (Figures 1 and 2). The similarities of  $^1J(^{129}\text{Xe}-^{14}\text{N})$  and  $^1J(^{129}\text{Xe}-^{19}\text{F})$  values to those of the  $\text{HC}\equiv\text{N}-\text{XeF}^+$  cation




**Figure 1.** N.m.r. spectra of  $\text{HC}\equiv\text{N-XeF}^+\text{AsF}_6^-$ : (a)  $^{129}\text{Xe}$  spectrum for a 99.2%  $^{13}\text{C}$  enriched sample recorded in HF solvent at  $-10^\circ\text{C}$ ; (b)  $^1\text{H}$  spectrum for a natural abundance sample recorded in  $\text{BrF}_5$  solvent at  $-58^\circ\text{C}$  at 1.8790 T; (c)  $^{13}\text{C}$  spectrum for a 99.2%  $^{13}\text{C}$  enriched sample recorded in HF at  $-10^\circ\text{C}$ . Asterisks (\*) denote  $^{129}\text{Xe}$  satellites.

also allow analogous structures to be assigned to the alkyl-, fluoroalkyl- and pentafluorobenzonitrile cations.

The observation of  $^1J(^{129}\text{Xe}-^{14}\text{N})$  in both the  $^{14}\text{N}$  and  $^{129}\text{Xe}$  spectra is particularly noteworthy and provides conclusive proof that in each case the nitrogen atom is directly bonded to xenon. This is confirmed by comparing their respective reduced coupling constants,  $^1K(\text{Xe}-\text{N})$  (calculated using the expression given in refs. 1–3; (see Table 1), with that of  $\text{FXeN}(\text{SO}_2\text{F})_2$  [ $^1J(^{129}\text{Xe}-^{15}\text{N})$  307 Hz;  $^1K(\text{Xe}-\text{N})$   $0.913 \times 10^{22}$   $\text{NA}^{-2} \text{m}^{-3}$ ].<sup>1</sup> In addition, the small value of the  $J$ -coupling,  $^{129}\text{Xe}-^{13}\text{C}$ , observed for the  $^{13}\text{C}$  enriched sample of  $\text{H-C}\equiv\text{N-XeF}^+$ , and a one-bond  $^{13}\text{C}-^1\text{H}$  coupling, also confirm that the xenon atom is not bonded to carbon in this species.

Assuming that the Xe-N spin-spin couplings in xenon-nitrogen compounds are dominated by the Fermi contact term, a comparison of  $^1K(\text{Xe}-\text{N})$  values for  $\text{R-C}\equiv\text{N-XeF}^+$  with that of the trigonal planar  $\text{sp}^2$ -hybridised nitrogen atom in  $\text{FXeN}(\text{SO}_2\text{F})_2$ <sup>1</sup> allows assessment of the relative degrees of hybridisation for the nitrogen orbitals used in bonding to



**Figure 2.** N.m.r. spectra of  $\text{CH}_3\text{C}\equiv\text{N-XeF}^+\text{AsF}_6^-$  recorded in HF solvent at  $-10^\circ\text{C}$ : (a) and (b) are  $^{129}\text{Xe}$  spectra, where (a) is natural abundance and (b) is 99.7%  $^{13}\text{C}$  enriched at the 2-carbon; (c) natural abundance  $^{14}\text{N}$  spectrum. Asterisks (\*) denote  $^{129}\text{Xe}$  satellites.

xenon. The ratios of  $[^1K(\text{Xe}-\text{N})]_{\text{sp}}$  to  $[^1K(\text{Xe}-\text{N})]_{\text{sp}^2}$  are 1.42–1.53 for the cations listed in Table 1, in excellent agreement with the theoretical ratio, 1.50, calculated from the predicted fractional s-characters of the nitrogen orbitals used in bonding to xenon.

In the case of the perfluoroalkyl derivatives ( $\text{R} = \text{C}_2\text{F}_5$  and  $\text{C}_3\text{F}_7$ ), the Xe-N bonds are found to be labile on the n.m.r. time scale in HF solvent at temperatures down to  $-30^\circ\text{C}$  as a result of a decrease in base strength for  $\text{RC}\equiv\text{N}$ : with increasing fluorine substitution at C-2.

The salts  $\text{HC}\equiv\text{N-XeF}^+\text{AsF}_6^-$  and  $\text{CH}_3\text{C}\equiv\text{N-XeF}^+\text{AsF}_6^-$  were isolated from HF solvent by pumping the solutions under vacuum at  $-50$  to  $-30^\circ\text{C}$  to give white solids whose Raman spectra were recorded at  $-196^\circ\text{C}$  (514.5 nm excitation). The Raman spectrum of  $\text{HC}\equiv\text{N-XeF}^+\text{AsF}_6^-$  was assigned in detail while only key frequencies of  $\text{CH}_3\text{C}\equiv\text{N-XeF}^+\text{AsF}_6^-$  have presently been assigned. In addition to the three Raman active

**Table 1.** N.m.r. parameters<sup>a</sup> for RC≡N-XeF<sup>+</sup> cations.Chemical shifts<sup>b</sup>

R	$\delta(^{129}\text{Xe})$	$\delta(^{19}\text{F})^c$	$\delta(^{14}\text{N})$	$\delta(^{13}\text{C})$	$\delta(^1\text{H})$
H <sup>d</sup>	-1552 -1569 <sup>e</sup>	-198.4	-235.4	104.1	6.01 5.67 <sup>f</sup>
CH <sub>2</sub> F	-1541	-198.4 (XeF) -241.7 (CF)	-229.2		5.44
CH <sub>3</sub>	-1708	-185.5	-251.1	115.3 (CN) <sup>g</sup> 0.6 (CH <sub>3</sub> ) <sup>h</sup>	2.41
C <sub>2</sub> H <sub>5</sub>	-1717	-184.6	-251.9		1.29 (CH <sub>3</sub> ) 2.80 (CH <sub>2</sub> )
C <sub>6</sub> F <sub>5</sub> <sup>i</sup>	-1426				

Coupling Constants<sup>j</sup>

	HC≡N-XeF <sup>+</sup> c	CH <sub>2</sub> FC≡N-XeF <sup>+</sup>	CH <sub>3</sub> C≡N-XeF <sup>+</sup> f,g	C <sub>2</sub> H <sub>5</sub> C≡N-XeF <sup>+</sup>	C <sub>6</sub> F <sub>5</sub> C≡N-XeF <sup>+</sup>
<sup>1</sup> J ( <sup>129</sup> Xe- <sup>14</sup> N)	334	333	313	311	
<sup>1</sup> J ( <sup>129</sup> Xe- <sup>19</sup> F)	6150 (6181) <sup>e</sup>	6163	6020	6017	6610
<sup>1</sup> J ( <sup>14</sup> N- <sup>13</sup> C)	22				
<sup>1</sup> J ( <sup>13</sup> C- <sup>1</sup> H)	308		141		
<sup>2</sup> J ( <sup>129</sup> Xe- <sup>13</sup> C)	84		79		
<sup>2</sup> J ( <sup>19</sup> F- <sup>1</sup> H)		44			
<sup>2</sup> J ( <sup>19</sup> F- <sup>14</sup> N)			18		
<sup>3</sup> J ( <sup>129</sup> Xe- <sup>1</sup> H)	26.8				
<sup>3</sup> J ( <sup>19</sup> F- <sup>13</sup> C)	18		19		
<sup>3</sup> J ( <sup>1</sup> H- <sup>1</sup> H)				7.5	
<sup>4</sup> J ( <sup>19</sup> F- <sup>1</sup> H)	2.6				
<sup>1</sup> K (Xe-N)	1.393 × 10 <sup>22</sup>	1.389 × 10 <sup>22</sup>	1.305 × 10 <sup>22</sup>	1.297 × 10 <sup>22</sup>	

<sup>a</sup> Spectra were recorded in anhydrous HF at -10 °C using 9 mm o.d. FEP sample tubes at 5.8719 T, and spectrometer frequencies (MHz): <sup>129</sup>Xe 69.563, <sup>19</sup>F 235.361, <sup>14</sup>N 18.075, <sup>13</sup>C 62.915, <sup>1</sup>H 250.132. <sup>b</sup> Samples were referenced externally at 24 °C with respect to the neat liquid references: XeOF<sub>4</sub> (<sup>129</sup>Xe), CFCl<sub>3</sub> (<sup>19</sup>F), CH<sub>3</sub>NO<sub>2</sub> (<sup>14</sup>N), SiMe<sub>4</sub> (<sup>13</sup>C and <sup>1</sup>H). A positive chemical shift denotes a resonance occurring to high frequency of the reference compound. <sup>c</sup> All <sup>19</sup>F spectra displayed a broad saddle-shaped feature at ca. -68 p.p.m. arising from the partially quadrupole collapsed <sup>75</sup>As-<sup>19</sup>F coupling of the octahedral AsF<sub>6</sub><sup>-</sup> anion. <sup>d</sup> Sample prepared from 99.2% <sup>13</sup>C enriched H<sup>13</sup>CN. <sup>e</sup> Measured in BrF<sub>5</sub> solvent at -58 °C; the <sup>1</sup>J(<sup>129</sup>Xe-<sup>14</sup>N) coupling was found to be completely quadrupole collapsed under these conditions. The sample was prepared by redissolving solid HC≡N-XeF<sup>+</sup>AsF<sub>6</sub><sup>-</sup>, prepared in HF solvent, in BrF<sub>5</sub> at -50 °C. <sup>f</sup> Measured in a 5 mm o.d. precision glass tube in BrF<sub>5</sub> solvent at -58 °C, at 80.022 MHz (1.8790 T) for <sup>1</sup>H. The sample was prepared by redissolving solid HC≡N-XeF<sup>+</sup>AsF<sub>6</sub><sup>-</sup>, prepared in HF solvent, in BrF<sub>5</sub> at -50 °C. <sup>g</sup> Sample prepared from 99% <sup>13</sup>C enriched CH<sub>3</sub><sup>13</sup>CN. <sup>h</sup> Sample prepared from 99.7% <sup>13</sup>C enriched <sup>13</sup>CH<sub>3</sub>CN. <sup>i</sup> Decomposition occurred at -10 to -20 °C, preventing fuller characterisation. <sup>j</sup> *J* values in Hz; *K* values in NA<sup>-2</sup>m<sup>-3</sup>.

modes consistent with an octahedral AsF<sub>6</sub><sup>-</sup> anion at 679(50),  $\nu_1(a_{1g})$ ; 581(13),  $\nu_2(e_g)$ ; 371(15),  $\nu_5(t_{2g})$  cm<sup>-1</sup>; and the formally Raman inactive modes  $\nu_3(t_{1u})$  at 692(12) and  $\nu_4(t_{1u})$  at 244(1), 269(6), 280(14) cm<sup>-1</sup>; the most prominent features of the linear HC≡N-XeF<sup>+</sup> cation spectrum are the factor-group-split C≡N stretch at 2159(41) and 2163(18) cm<sup>-1</sup> and a pair of intense lines at 559(100) and 569(94) cm<sup>-1</sup> assigned to the factor-group-split Xe-F stretch. <sup>13</sup>C enrichment (99.2%) confirms the assignment of  $\nu(\text{C}\equiv\text{N})$  (isotopic shift, 31.6 cm<sup>-1</sup>) and the split, doubly degenerate bend  $\delta(\text{C}\equiv\text{N}-\text{Xe})$  at 327(4), 334(2) cm<sup>-1</sup> (isotopic shift, 5.5 cm<sup>-1</sup>); also *cf.* the N≡C-I bend in ICN at 304 cm<sup>-1</sup>.<sup>4</sup> A low-frequency shoulder on  $\nu_5$  of the anion at 368 cm<sup>-1</sup> was tentatively assigned to the Xe-N stretch (*cf.*  $\nu(\text{Xe}-\text{N})$  of FXeN(SO<sub>2</sub>F)<sub>2</sub> at 422 cm<sup>-1</sup>)<sup>1</sup> and a weak band at 3141(4) cm<sup>-1</sup> was assigned to  $\nu(\text{C}-\text{H})$  (*cf.* the C-H stretching frequency of gaseous HC≡N, 3311 cm<sup>-1</sup>).<sup>5</sup> The remaining features were assigned to the doubly degenerate bends  $\delta(\text{F}-\text{Xe}-\text{N})$  [116(33), 133(10), 157(5), 180(2) cm<sup>-1</sup>] and  $\delta(\text{H}-\text{C}\equiv\text{N})$  [706(1) cm<sup>-1</sup>]. Preliminary assignments of some key frequencies for the CH<sub>3</sub>C≡N-XeF<sup>+</sup> cation spectrum were made by comparison with those of CH<sub>3</sub>C≡N<sup>6</sup> and HC≡N-XeF<sup>+</sup>:  $\nu_{\text{asym}}(\text{CH}_3)$ , 3013(6), 3021(7), 3027(4) cm<sup>-1</sup>;  $\nu_{\text{sym}}(\text{CH}_3)$ , 2944(19), 2949(19) cm<sup>-1</sup>;  $\nu_{\text{asym}}(\text{CH}_3)$ , 2335(16) cm<sup>-1</sup>;  $\nu(\text{XeF})$ , 559(100), 570(74), 571(34) cm<sup>-1</sup>;  $\delta(\text{F}-\text{Xe}-\text{N})$ , 160(7), 170(6) cm<sup>-1</sup>.

Additional examples of nitriles and other inorganic and organic nitrogen bases are under active investigation as potential electron-pair donors towards noble gas-cations, as well as representative X-ray crystal structures containing the RC≡N-XeF<sup>+</sup> cations.

This research was sponsored by the United States Air Force Astronautics Laboratory, Edwards Air Force Base, California (Contract F49620-87-C-0049) and a Natural Sciences and Engineering Research Council of Canada (N.S.E.R.C.C.) operating grant.

Received, 9th June 1987; Com. 797

## References

- J. F. Sawyer, G. J. Schrobilgen, and S. J. Sutherland, *Inorg. Chem.*, 1982, **21**, 4064.
- G. A. Schumacher and G. J. Schrobilgen, *Inorg. Chem.*, 1983, **22**, 2178.
- R. Faggiani, D. K. Kennepohl, C. J. L. Lock, and G. J. Schrobilgen, *Inorg. Chem.*, 1986, **25**, 563.
- S. Hemple and E. R. Nixon, *J. Chem. Phys.*, 1967, **47**, 4273.
- H. C. Allen, E. D. Tidwell, and E. K. Plyler, *J. Chem. Phys.*, 1956, **25**, 302.
- J. E. Griffiths in 'Vibrational Spectra and Structure,' ed. J. R. Durig, Elsevier, New York, 1977, vol. 6, p. 333.