


Corrigenda

Synthesis of a New Bifunctional Hemilabile Ligand by Insertion of PhNCO into a Co-ordinated β -Phosphinoketonate. Crystal Structure of



Salah-Eddine Bouaoud, Pierre Braunstein, Daniel Grandjean, Dominique Matt, and Dominique Nobel

J. Chem. Soc., Chem. Commun., 1987, 488.

On page 489, in the ^1H n.m.r. data, the assignments for the NMe_2 and NCH_2 signals should be interchanged between isomers (2a) and (2b).

Stabilisation of Mononuclear Palladium(III). The Single Crystal X-Ray Structure of the $[\text{Pd}(\text{L})_2]^{3+}$ Cation (L = 1,4,7-trithiacyclononane)

Alexander J. Blake, Alan J. Holder, Timothy I. Hyde, and Martin Schröder

J. Chem. Soc., Chem. Commun., 1987, 987.

The redox potentials for the complex cation $[\text{Rh}(\mathbf{1})_2]^{3+}$ [$\mathbf{1}$ = 1,4,7-trithiacyclononane] are quoted vs. Ag/Ag^+ and not vs. Fc^0/Fc^+ as indicated. The correct values are $^1E_{\frac{1}{2}} = -0.71$ and $^2E_{\frac{1}{2}} = -1.08$ V vs. Fc^0/Fc^+ .

Stereoselective Synthesis of the Dihydrobenzo[b]furan Segments of the Ephedradine Alkaloids

Raymond Baker, Nigel G. Cooke, Guy R. Humphrey, Stanley H. B. Wright, and Jordan Hirshfield

J. Chem. Soc., Chem. Commun., 1987, 1102.

References 11—13 should be as follows.

11 J. Bartroli, D. A. Evans, and T. L. Shih, *J. Am. Chem. Soc.*, 1981, **103**, 2127.

12 D. A. Evans, J. V. Nelson, T. R. Tabor, and R. Vogel, *J. Am. Chem. Soc.*, 1981, **103**, 3099.

13 Compound (**11a**) was shown to be homogeneous by ^1H n.m.r. spectroscopy and h.p.l.c.