

Synthesis of (\pm)-Nitramine and (\pm)-Isonitramine









William Carruthers and Roger C. Moses

Chemistry Department, The University, Stocker Road, Exeter EX4 4QD, U.K.

The alkaloids (\pm)-nitramine (**1**) and (\pm)-isonitramine (**2**) have been synthesised, employing an intramolecular Mannich reaction to set up the spirocyclic ring system.

Nitramine (**1**) and isonitramine (**2**), isolated recently from plants of the genus *Nitraria*,¹ are of interest because of their unusual spirocyclic structures and their structural relationship to the neurotoxic alkaloid histrionicotoxin (**3**). The structures of the new alkaloids were established by *X*-ray crystallography of crystalline salts, which showed that they are diastereoisomers;² in each isomer the hydroxy substituent is equatorially disposed in the cyclohexane ring. Two syntheses of (\pm)-nitramine³ and one of (\pm)-isonitramine⁴ have been recorded. We report here a direct route to both alkaloids using an intramolecular Mannich reaction to set up the spirocyclic ring system.

The readily available nitrile (**4**) was converted into the acetal (**5**) and thence into the benzylamine (**6**) by sequential LiAlH_4 reduction, benzylation, and LiAlH_4 reduction. Cyclisation was effected by bubbling gaseous formaldehyde into a solution of amine (**6**) in methanol containing hydrochloric acid⁵ and gave the spirocyclic ketone (**7**) in 60% yield. Reduction of the carbonyl group with sodium borohydride afforded a mixture of the alcohols (**8**) and (**9**) (*ca.* 1:1) which were separated by chromatography on silica gel. Debenzylation of (**8**) was smoothly effected with palladium-carbon in boiling methanol in the presence of ammonium formate⁶ and gave (\pm)-nitramine (**1**) as an oil, characterised by its high



Bzl = benzyl

resolution mass spectrum and its ^1H and ^{13}C n.m.r. spectra, which were in close agreement with the published spectra.² Debenzylation of (9) similarly gave (\pm)-isonitramine (2) as a crystalline solid, m.p. 100.5–103°C; its structure was fully supported by the high resolution mass spectrum and the ^1H and ^{13}C n.m.r. spectra. The nitramine and isonitramine were easily distinguished from each other by the C-1 ^{13}C n.m.r. signals at δ 52.32 in the former and δ 60.71 in the latter (lit.² δ 52.0 for nitramine and 60.3 for isonitramine).

We thank the Ministry of Defence for support of this work.

Received, 4th December 1986; Com. 1726

References

- 1 N. Yu Novgorodova, S. Kh. Maekh, and S. Yu Yunusov, *Chem. Nat. Prod.*, 1973, 191; *Khim. Prir. Soedin.*, 1973, 196; Z. Osmanov, A. A. Ibragimov, and S. Yu Yunusov, *Chem. Nat. Prod.*, 1977, 607; *Khim. Prir. Soedin.*, 1977, 720; Z. Osmanov, A. A. Ibragimov, and S. Yu Yunusov, *Chem. Nat. Prod.*, 1983, 121; *Khim. Prir. Soedin.*, 1981, 126.
- 2 A. A. Ibragimov, Z. Osmanov, B. Tashkhodzhaev, N. D. Abdullaev, M. R. Yagudaev, and S. Yu Yunusov, *Chem. Nat. Prod.*, 1982, 458; *Khim. Prir. Soedin.*, 1981, 623; B. Tashkhodzhaev, *Chem. Nat. Prod.*, 1983, 70; *Khim. Prir. Soedin.*, 1981, 75. There seems to be some confusion in the Russian papers in the attribution of crystal structures to nitramine and isonitramine. In the two papers above the structures of the two alkaloids are interchanged, without comment. We have taken the second paper as the definitive one.
- 3 B. B. Snider and C. P. Cartaya-Marin, *J. Org. Chem.*, 1984, **49**, 1688; L. H. Hellberg, C. Beeson, and R. Somanathan, *Tetrahedron Lett.*, 1986, **27**, 3955.
- 4 A. P. Kozikowski and P.-W. Yuen, *J. Chem. Soc., Chem. Commun.*, 1985, 847; *ibid.*, 1985, 1548; see also J. B. Mieczkowski, *Bull. Polish Acad. Sci.*, 1985, **33**, 13.
- 5 Cf. E. Wenkert, K. G. Dave, and R. V. Stevens, *J. Am. Chem. Soc.*, 1968, **90**, 6177.
- 6 A. M. Koskinen and H. Rapoport, *J. Med. Chem.*, 1985, **28**, 1301.