

A Planar Binuclear Phthalocyanine and Its Dicobalt Derivatives

Clifford C. Leznoff,* Herman Lam, Sebastian M. Marcuccio, W. Andrew Nevin, Pavel Janda, Nagao Kobayashi, and A. B. P. Lever*

Department of Chemistry, York University, North York (Toronto), Ontario, Canada M3J 1P3

The synthesis, characterisation, and intramolecular interactions of a metal-free planar binuclear phthalocyanine and its cobalt derivatives are described.

Binuclear phthalocyanines covalently linked by 5,^{1a,b} 4,^{1c} 2,^{1c} 1,^{1d} and 0² atom bridges have been recently described. We now report a new planar binuclear phthalocyanine (**1**) in which two phthalocyanine rings share a common benzene ring. Since this lacks a benzene ring relative to the other known binuclear phthalocyanines, this is referred to as the (-1) linked binuclear phthalocyanine (**1a**).

We have previously studied intramolecular interactions in binuclear phthalocyanines² as a function of bridge link and it was obviously of interest to observe such interactions in (**1**). Indeed (**1a**) and its derivatives show spectroscopic and electrochemical evidence for greater intramolecular interactions than have been seen previously in binuclear phthalocyanines. Compound (**1**) also has value as a building block in making doubly-linked stacked conducting metallomacrocyclic polymers.³


Treatment of 4-neopentoxypthalonitrile^{1a,b} and 1,2,4,5-tetracyanobenzene⁴ with ammonia^{1b} gave the isoindolines (**2**) and (**3**) respectively. A mixed condensation of an excess of (**2**) with (**3**) in 2-*N,N*-dimethylaminoethanol under standard conditions^{1a-d} gave (**1a**) after flash and gel-permeation chromatography.⁵ The dicobalt(II) derivative (**1b**) was prepared by previously described insertion methods¹ in 63% yield from (**1a**). Compounds (**1a**) and (**1b**) gave parent ions in their mass spectra at 1468 and 1581 respectively using the fast atom bombardment (f.a.b.) technique¹ and were fully characterised by i.r., n.m.r., and elemental analysis. The inner NH protons of (**1a**) gave broad absorption peaks between -2 and -3 p.p.m. typical of other binuclear phthalocyanines.¹

Both species (**1a**) and (**1b**) have electronic spectra (Figures 1 and 2) which are quite atypical, with broad absorption in the Q band region⁶ indicative² of intramolecular electronic coupling. The characteristic double Q band of metal-free phthalocyanines² is barely evident even at 10⁻⁷ M concentrations. Q band S₁ emission is still observed but with an unusually broad² excitation spectrum (Figure 1).

Given the flat nature of the species (**1**), aggregation, prevalent in the phthalocyanine series⁷ is important. Both the metal-free and cobalt(II) species aggregate in micromolar concentration solutions. In *o*-dichlorobenzene (DCB), the aggregation constants, for dimer formation, for (**1a**) and (**1b**) are (9.9 ± 0.9) × 10⁴ dm³ mol⁻¹ and (8.6 ± 0.6) × 10⁴ dm³ mol⁻¹ respectively, more than an order of magnitude higher than those for our previously investigated binuclear species,⁸ but an order lower than that for a tetranuclear species.⁵

Cyclic voltammetry of (**1b**) in DCB solution (with 0.3 M tetrabutylammonium perchlorate, TBAP, Pt disc working electrode) consists of a series of broad peaks. Half-wave potentials were obtained by differential pulse voltammetry. Species (**1b**) shows oxidation at +0.17 and +0.59 V and reduction at -0.81, -1.69, and -1.96 V vs. Fc⁺/Fc.⁹ The three most positive waves show evidence for splitting being broad or having well pronounced shoulders. Species (**1b**) is easier to reduce and more difficult to oxidise than previously described cobalt(II) mononuclear and binuclear analogues,^{1,10} being reduced at some 0.3 V less negative a potential.

Oxidised and reduced species were obtained using an optically transparent thin layer electrode (OTTLE) cell



utilising a gold minigrad working electrode. Stepwise oxidation across the first oxidation couple gives a spectrum typical of a phthalocyanine cation radical species^{5,6b,10,11} [Figure 2(a)], corresponding to the oxidation of each of the phthalocyanine rings and with no clear spectroscopic evidence of any one-electron oxidised intermediate. In contrast, reduction over the double peak of the first reduction couple (-0.81 V) occurs *via* two consecutive one-electron steps, corresponding to the consecutive reduction of each of the two cobalt atoms to Co^I [Figure 2(b)]. This is the first report of a mixed valence Co^{II}-Co^I phthalocyanine species, (1c). It displays a specific electronic spectrum [Figure 2(b)] different from either (1b) or the doubly reduced bis-Co^I species, (1d). This is in contrast to cofacial mixed valence Co^{II}-Co^I porphyrin species¹² whose spectra show little evidence for interaction between the two halves of the molecule. The spectrum of (1d) shows some unusual features not seen previously with Co^I phthalocyanine species,^{6b,10,13,14} having a split Q band and an additional strong absorption peak at 825 nm. Both (1c) and (1d) show Q band and MLCT (metal to ligand charge transfer) absorption (near 500 nm) red shifted relative to less electronically coupled species.^{1,10} Polarisation of the OTTL negative of the couple at -1.69 V results in formation of a ring-reduced species, whose quite typical, but red shifted spectrum is also shown in Figure 2(b).

Electrocatalytic reduction of oxygen was examined at electrodes (glassy carbon and ordinary and stress annealed pyrolytic graphite) with adsorbed (1b). Oxygen reduction occurred at -0.34 V vs. saturated calomel electrode (s.c.e.) in cyclic voltammetric curves through pH 1 to 13, and the limiting current corresponding to two electron reduction of oxygen to hydrogen peroxide was observed in rotating disc experiments. The absence of any dependence of the oxygen reduction potential upon pH down to pH 1 is noteworthy.

This system represents an important new class of phthalocyanine species which should prove of importance in photo-




Figure 1. Electronic absorption spectrum of (1a) in DCB (—) at 1×10^{-6} M. Emission (λ_{ex} , 340 nm) (---) and excitation (λ_{em} , 715 nm) (·····) spectra of (1a) in 1:1 ethanol-dichloromethane; a.u. = arbitrary units.




Figure 2. (a) Development of the electronic spectrum with time, during the oxidation of (1b) at +0.55 V vs. Fc⁺/Fc in DCB (0.3 M TBAP) to a Co^{II}-Co^{II} ring oxidised di-cation radical. (b) Electronic absorption spectra of (1b) in DCB, after electrolysis at -0.90 V (—), -1.25 V (---), and -1.85 V (·····) vs. Fc⁺/Fc. The species are, neglecting axial ligands, the mixed valence Co^{II}-Co^I, the Co^I-Co^I, and a Co^I-Co^I ring reduced species, respectively. The insets show the near i.r. region.

voltaic and photo- and electro-catalytic applications, as well as in the field of molecular metals.¹⁵

We are indebted to the Natural Sciences and Engineering Research Council (Ottawa) and the Office of Naval Research (Washington) for financial support. We also thank Penny Seymour for spectroscopic assistance. P. J. is a visiting scholar from the Heyrovsky Institute, Prague, Czechoslovakia, and

N. K. is a visiting Professor from Tohoku University, Sendai, Japan.

Received, 22nd December 1986; Com. 1804

References

- (a) C. C. Leznoff, S. Greenberg, S. M. Marcuccio, P. C. Minor, P. Seymour, A. B. P. Lever, and K. B. Tomer, *Inorg. Chim. Acta*, 1984, **89**, L35; (b) C. C. Leznoff, S. M. Marcuccio, S. Greenberg, A. B. P. Lever, and K. B. Tomer, *Can. J. Chem.*, 1985, **63**, 623; (c) S. M. Marcuccio, P. I. Svirskaya, S. Greenberg, A. B. P. Lever, C. C. Leznoff, and K. B. Tomer, *ibid.*, 1985, **63**, 3057; (d) S. Greenberg, S. M. Marcuccio, C. C. Leznoff, and K. B. Tomer, *Synthesis*, 1986, 406.
 - E. S. Dodsworth, A. B. P. Lever, P. Seymour, and C. C. Leznoff, *J. Phys. Chem.*, 1985, **89**, 5698.
 - C. W. Dirk, T. Inabe, K. F. Schoch, Jr., and T. J. Marks, *J. Am. Chem. Soc.*, 1983, **105**, 1539; B. N. Diel, T. Inabe, N. K. Jaggi, J. W. Lyding, O. Schneider, M. Hanack, C. R. Kannewurf, T. J. Marks, and L. H. Schwartz, *ibid.*, 1984, **106**, 3207.
 - (a) C. Piechocki and J. Simon, *J. Chem. Soc., Chem. Commun.*, 1985, 259; (b) D. Wöhrle and W. Wahl, *Tetrahedron Lett.*, 1979, 227.
 - W. A. Nevin, W. Liu, S. Greenberg, M. R. Hempstead, S. M. Marcuccio, M. Melnik, C. C. Leznoff, and A. B. P. Lever, *Inorg. Chem.*, 1987, **26**, in the press.
 - (a) A. B. P. Lever, *Adv. Inorg. Chem. Radiochem.*, 1965, **7**, 27; (b) P. C. Minor, M. Gouterman, and A. B. P. Lever, *Inorg. Chem.*, 1985, **24**, 1894.
 - (a) K. Bernauer and S. Fallab, *Helv. Chim. Acta*, 1961, **44**, 1287; (b) Y-C. Yang, J. R. Ward, and R. P. Seiders, *Inorg. Chem.*, 1985, **24**, 1765; (c) A. R. Monahan, J. A. Brado, and A. F. DeLuca, *J. Phys. Chem.*, 1972, **76**, 1994.
 - W. A. Nevin, W. Liu, and A. B. P. Lever, *Can. J. Chem.*, 1987, **65**, in the press.
 - Potentials were referenced internally to the ferrocenium/ferrocene (Fc^+/Fc) couple, which lies at +0.16 V vs. saturated calomel electrode (s.c.e.), see R. R. Gagne, C. A. Koval, and D. C. Lisensky, *Inorg. Chem.*, 1980, **19**, 2854; G. Gritzner and J. Kuta, *Electrochim. Acta*, 1984, **29**, 869.
 - W. A. Nevin, M. R. Hempstead, W. Liu, C. C. Leznoff, and A. B. P. Lever, *Inorg. Chem.*, 1987, **26**, 570.
 - T. Nyokong, Z. Gasyna, and M. J. Stillman, *Inorg. Chim. Acta*, 1986, **112**, 11; G. Ferraudi, S. Oishi, and S. Muraldiharan, *J. Phys. Chem.*, 1984, **88**, 5261; D. Dolphin, B. R. James, A. J. Murray, and J. Thornback, *Can. J. Chem.*, 1980, **58**, 1125.
 - (a) J. P. Collman, M. Marrocco, C. M. Elliot, and M. L'Her, *J. Electroanal. Chem.*, 1981, **124**, 113; (b) Y. Le Mest, M. L'Her, J. Courtot-Coupez, J. P. Collman, E. R. Evitt, and C. S. Bencosme, *ibid.*, 1985, **184**, 331.
 - W. A. Nevin, W. Liu, M. Melnik, and A. B. P. Lever, *J. Electroanal. Chem.*, 1986, **213**, 217.
 - M. J. Stillman and A. J. Thompson, *J. Chem. Soc., Faraday Trans. 2*, 1974, **70**, 790.
 - A. B. P. Lever, M. R. Hempstead, C. C. Leznoff, W. Liu, M. Melnik, W. A. Nevin and P. Seymour, *Pure Appl. Chem.*, 1986, **58**, 1467.
-