

Radical Combination in the *ortho* Position of Trityl Radical Observed in Single-electron Transfer Reactions of Trityl Anion

Jürgen Werry, Pen-Yuan Lin, Konstantinos Bellos, Petros Assithianakis and Helmut Stamm*

Pharmazeutisch-Chemisches Institut der Universität Heidelberg, Im Neuenheimer Feld 364, D-6900 Heidelberg, FRG

Single-electron transfer reactions between trityl anion and 1-acyl-2,2-dimethylaziridines provide, among other products, the methallyl amides **7** and the triphenylmethanes **8** carrying an amidoethyl chain attached with the tertiary carbon *ortho* to the triphenylmethane.

Trityl radical (Tr = CPh₃) Tr[•] combines with radicals in the *para* position; only radicals of low steric demand add to the central carbon.¹ Combination in the *ortho* position has not been reported to the best of our knowledge. Space-filling models show that attack upon an *ortho* carbon should be possible for *t*-butyl or radicals of similar steric demand. *para*-Central ambidenticity and steric control² is also known for trityl anion Tr⁻ with possible² or proved^{3,4} involvement of single-electron transfer (SET).

Ketyl **2**, generated by electron attachment to **1**, forms the radical **3** which can react further.⁵ When the electron source is a nucleophile, this SET reaction is not possible unless the competing S_N2 attack is sufficiently slow. Unhindered S_N2 reaction of Tr⁻ with **4** exclusively yields the central products **5**.⁶ We now find that the reaction of Tr⁻ with **1a–c** can yield all three substitution products **6**, **8** and **9**. Compounds **8** are the first *ortho* products that are the major products of reactions of **1a** and **b** while the central product **6** is always the minor product. Compound **8** arises most probably from a combination of Tr[•] with radical **3**.

The main products in all runs are the methallylamides **7** which slowly isomerize to the corresponding enamides under base catalysis.⁵ This isomerization guarantees that **7** or rather the amide anions derived from **7** arise prior to work-up. There can be no doubt that **7** is formed *via* transfer of a hydrogen atom from radical **3** to Tr[•]. Thus, the main products indicate an SET mechanism. The second most important products obtained from **1a,b** are the *ortho* products **8a,b** (runs 1 and 2). The *para* products **9a,b** could only be identified from ¹H NMR spectroscopic signals.† The central product **6a** was not found, *cf.* **6b**. Structure assignment for **8** and **9** was based primarily on the chemical shift data (6.3–6.4 ppm) for the methine proton in the trityl moiety. In the common propeller conformation of triphenylmethanes the methine singlet appears at 5.5 ± 0.1 ppm.^{2,4,7} The downfield shift in **8** indicates a 'steeper' conformation for the propeller blades caused by the *ortho* substituent.⁷ Structure **8** was proved by NOE (**8b**) and by the isolation of the non-aromatized precursor **10a** of **8a** in a nearly pure state, when a repetition of run 1 was quenched after 15 min. Structure **10a** follows from the ¹H NMR multiplets of the

† An example of characterisation for **8a**, **9a** and **10a**. **8a**: m.p. 207–209 °C. IR (KBr) $\nu = 3335$ cm⁻¹ (NH), 1646 (amide I), 1550 (amide II). ¹H NMR (250 MHz, CDCl₃) $\delta = 0.87$ (s, 9H, Bu^t), 1.51 (s, 6H, CMe₂), 3.60 (d, *J* 6.0 Hz, 2H, CH₂), 4.76 (t, br, *J* 6 Hz, 1H, NH), 6.30 (s, 1H, methine-H), 6.98–7.08 (m, 4H, *o*-H of 2 Ph), 7.08–7.13 (m, 1H, *o'*-H of C₆H₄), 7.16–7.32 (m, 8 aromatic H), 7.38–7.45 (m, 1H, *m*-H of C₆H₄).

9a (only in mixture with **8a**): ¹H NMR (250 MHz, CDCl₃) $\delta = 1.05$ (s, 9H, Bu^t), 1.31 (s, 6H, CMe₂), 3.39 (d, *J* 5.9 Hz, 2H, CH₂), 5.24 (s, br, 1H, NH), 5.54 (s, 1H, methine-H), the signals of the aromatic H are hidden under signals of **8a**.

10a: Oil. IR (film) $\nu = 3370$ cm⁻¹ (NH), 1654 (amide I), 1518 (amide II). ¹H NMR (500 MHz, CDCl₃) $\delta = 0.79$ (s, 3H, 1 Me of CMe₂), 0.80 (s, 3H, 1 Me of CMe₂), 1.04 (s, 9H, Bu^t), 2.88 (dd, *J* 13.8, 5.1 Hz, 1H of CH₂), 3.38 (dd, *J* 13.8, 7.5 Hz, 1H of CH₂), 3.60 (d, *J* 6.0 Hz, 1H, 1-H), 5.25 (t, br, *J* 6 Hz, 1H, NH), 5.78 (dd, *J* 9.9, 5.2 Hz, 1H, 4-H).

five spin system 1-H to 5-H (supported by COSY) and from a pronounced diastereotopic NCH₂ group.†


There can be little doubt that **10** and hence **8** arise by combination of the two radicals Tr[•] and **3**, which is probably favoured by a rather low steric demand and high selectivity of **3**. Attachment of a hydrogen atom to **3** (final product **11** formed at the expense of **7**) was observed only when Tr⁻ was generated from triphenylmethane by means of sodium naphthalenide. Dihydronaphthalene resulting from this technique is a good donor of hydrogen atoms.

Reaction of **1c** with Li⁺Tr⁻ markedly changed the composi-

Table 1 Reactions of trityllithium with aziridines **1a–c** in THF^a

Run	Aziridine	Time/ days	Products [yields (%)]
1	1a	6	6a (0) 7a^b (59) 8a (22) 9a (trace)
2	1b	16	6b (5) 7b^b (57) 8b (21) 9b (1)
3	1c	6	6c (4) 7c^b (30) 8c (6) 9c (19) 10c (3) 11c (2)

^a Reactions with two equivalents of Tr⁻ at room temperature. Li⁺Tr⁻ was generated from triphenylmethane (small excess) and butyllithium (hexane) in THF. ^b In part isomerized to the respective enamide RCONHCH=CMe₂: 4% in run 1, 27% in run 2, 5% in run 3.



a; R = Bu^t
b; R = Ph
c; R = 4-biphenyl

tion of the product mixture (run 3). The yield of the *para* product (**9c**) increased at the expense of **7** and **8**. Besides, 2% of **11c** was found. A non-SET ring opening by the nucleophile Tr^- should lead to virtually identical results with **1b** and **1c**. However, the reduction potential⁸ for **1b** and the difference in potentials (*ca.* 0.3 V) for acetophenone and 4-phenylacetophenone⁹ allow us to conclude that the reduction potential of **1c** is close to the potential of bromobenzene. The Tolbert mechanism (radical addition to Tr^-) takes on chain character in the reaction of Tr^- with bromobenzene.⁴ Thus, we assume that the *para* adduct (a radical anion of the *para* analogue of **10** type) formed by addition of **3** to Tr^- is able to sustain a chain reaction by SET to **1c** but not to **1a,b**. SET to **3c** may be responsible for the formation of **11c**.

A tertiary radical of pronounced steric demand greater than **3** cannot add to an *ortho* position of Tr^{\cdot} as is shown by the reaction of Li^+Tr^- with 1-benzoyl-2-*t*-butyl-2-phenylaziridine. The corresponding radical (in which two methyl groups in **3b** are replaced by phenyl and *t*-butyl) adds to the *para* position only (9%).

We thank Professor Dabrowsky and Mr. Großkurth, Max-Planck-Institut für Medizinische Forschung, for the COSY spectrum (500 MHz).

Received, 31st May 1990; Com. 0/02441D

References

- 1 H. Lankamp, W. Th. Nauta and C. McLean, *Tetrahedron Lett.*, 1968, 249; H. A. Staab, H. Brettschneider and H. Brunner, *Chem. Ber.*, 1970, **103**, 1101; W. P. Neumann and R. Stapel, *Chem. Ber.*, 1986, **119**, 2006; M. E. Sigman and J. E. Leffler, *J. Org. Chem.*, 1987, **52**, 1165; W. A. Pryor, K. Smith, J. T. Echols Jr. and D. L. Fuller, *J. Org. Chem.*, 1972, **37**, 1753 and references cited therein.
- 2 H. O. House and P. D. Weeks, *J. Am. Chem. Soc.*, 1975, **97**, 2785 and references cited therein.
- 3 H. E. Zieger, I. Angres and L. Maresca, *J. Am. Chem. Soc.*, 1973, **95**, 8201.
- 4 L. M. Tolbert and D. Martone, *J. Org. Chem.*, 1983, **38**, 1185; L. M. Tolbert, *J. Am. Chem. Soc.*, 1980, **102**, 6808.
- 5 H. Stamm, P. Assithianakis, R. Weiss, G. Bentz and B. Buchholz, *J. Chem. Soc., Chem. Commun.*, 1984, 753. Subsequent work showed the reaction of *N*-acylaziridines with radical anions to be more complicated than depicted in this preliminary publication. Inconsistent results have only recently found an interpretation which rests on other work and the publication of which is in preparation.
- 6 H. Stamm and W. Wiesert, *Chem. Ber.*, 1978, **111**, 502.
- 7 *cf.*: H. Kessler, A. Moosmayer and A. Rieker, *Tetrahedron*, 1969, **25**, 287.
- 8 D. Archier-Jay, N. Besbes, A. Laurent, E. Laurent, S. Lesniak and R. Tardivel, *Bull. Soc. Chim. Fr.*, 1989, 537.
- 9 L. Meites and P. Zuman, *CRC Handbook Series in Organic Electrochemistry*, vol. I, CRC Press, Cleveland, 1976.