

Selectively Dehydroxylated Calix[4]arenes and 1,3-Dithiocalix[4]arenes; Novel Classes of Calix[4]arenes

Yuhua Ting, Willem Verboom, Leo C. Groenen, Jan-Dirk van Loon and David N. Reinhoudt*

Laboratory of Organic Chemistry, University of Twente, P.O. Box 217, 7500 AE Enschede, The Netherlands

Selective modification of calix[4]arenes either by selective dehydroxylation *via* the reductive removal of phosphate(s), or *via* the Newman-Kwart rearrangement of 1,3-bis(dimethylthiocarbamoyl)calix[4]arenes yields calix[4]arenes **3** and 1,3-dithiocalix[4]arenes **6**, respectively.

Calix[4]arenes¹ can easily be functionalised both at the phenolic OH groups (lower rim) and, after removal of the *t*-butyl groups, at the *para* positions of the phenol rings (upper rim). Recently we reported *e.g.* the regioselective introduction of two substituents at the upper rim² and the 1,3-*O*-alkylation of the lower rim.³ Subsequently, we have also studied the selective modification of calix[4]arenes at the diametrical (1,3-) positions, either *via* removal of two phenolic groups or *via* replacement of two of the original oxygen atoms by sulphur. The complete dehydroxylation of *p*-*t*-butylcalix[*n*]arenes (*n* = 4 and 8) by reductive cleavage of the corresponding diethyl phosphate esters, very recently published by Goren and Biali,⁴ prompts us to present our preliminary results on the selective modification of the lower rim.

Using a slightly modified literature procedure^{5,6} we reacted *p*-*t*-butylcalix[4]arene **1a**⁷ with 4 equiv. of diethyl chlorophosphate in tetrahydrofuran (THF) in the presence of K₂CO₃ as a base to give selectively the 1,3-diphosphorylated calix[4]arene **2a**[†],‡ in 74% yield. Analogously **2b** was obtained in 76% yield starting from 1,3-dimethoxycalix[4]arene **1b**³ using NaH as a base. The ¹H NMR spectrum of **2b** shows, in addition to one singlet at δ 3.04 (OCH₃), two ABq patterns at δ 4.48 and 3.77 (*J* 17.3 Hz) and δ 4.46 and 3.25 (*J* 13.1 Hz), which indicates that **2b** is present in the 1,2-alternate conformation.⁸ To the best of our knowledge only Atwood *et al.*⁹ and Gutsche *et al.*¹⁰ have isolated a calix[4]arene in the 1,2-alternate conformation.

Reaction of **2a** with 6 equiv. of potassium in liquid ammonia at -60 °C^{5,6} gave, after chromatography, the bis(dehydroxy-

lated) **3a** and the mono(dehydroxylated) calix[4]arene **3b** as crystalline compounds in yields of 71 and 14%, respectively. § The formation of **3b** is due to partial hydrolysis of one of the phosphate moieties. The singlet at δ 3.92 (**3a**) and the two singlets (1 : 1 ratio) at δ 3.91 and 3.86 (**3b**) of the methylene bridge protons in the ¹H NMR spectra indicate that these molecules are flexible. Reaction of **2b** with potassium in liquid ammonia afforded the bis(dehydroxylated) dimethoxycalix[4]arene **3c** as a 83 : 17 mixture of 1,3-*syn*-(1,3-alternate like) and 1,3-*anti*-configurations¶ in a yield of 55%. Varying the temperature from -20 °C to 60 °C did not influence this ratio. The presence of one singlet at δ 3.02 (OCH₃), and one ABq at δ 4.18 and 3.59 (*J* 14.9 Hz, ArCH₂Ar) and two singlets at δ 1.38 and 1.14 (Bu^t) indicates the predominant 1,3-*syn*-configuration. The 1,3-*anti*-configuration exhibits a corresponding ¹H NMR spectrum, however, with two striking differences, *viz.* a singlet of the methoxy group at δ 3.20 and an aromatic singlet at δ 5.95 (δ 6.30 in the 1,3-*syn*).

Phenols can be transformed into the corresponding thiophenols *via* the corresponding *O*-aryl dialkylthiocarbamates and *S*-aryl dialkylthiocarbamates.¹¹ We have used this Newman-Kwart rearrangement for the transformation of the bis(dehydroxylated) calixarene **3a** and the 1,3-dimethoxycalix[4]arene **1b** to the 1,3-dithiocalix[4]arenes **6a** and **6b**, respectively. Reaction of **3a** and **1b** with 2.5 equiv. of *N,N*-dimethylthiocarbamoyl chloride in *N,N*-dimethylformamide (DMF) in the presence of NaH as a base gave the *O*-dimethylthiocarbamates **4a** and **4b** in yields of 86 and 87%, respectively. Heating compounds **4a** and **b** at 320 °C for 4 h





Fig. 1



Scheme 1

§ The corresponding reaction with the tetraphosphate **2c**⁴ gave the tris(dehydroxylated)-**3d** and the tetrakis(dehydroxylated) calix[4]arene **3e**⁴ in yields of 23 and 41%, respectively. These results demonstrate that in addition to the complete dehydroxylated calix[4]arene **3e**⁴ the mono-, di- and tri-hydroxycalix[4]arenes (**3d**, **3a** and **3b**, respectively) are easily accessible.

¶ *anti* Implies that the two methoxy groups are at opposite sites of the molecule.

† Satisfactory analytical and spectral data were obtained for all new compounds.

‡ For reasons of simplicity the positions of the original OH groups in **1** are numbered as 1, 2, 3 and 4.

under argon afforded the rearranged *S*-dimethylthiocarbamates **5a** and **b** in yields of 64 and 90%, respectively. According to their ¹H NMR spectra || both **4a** and **5a** exist as a 1:1 mixture of the 1,3-*syn*- and 1,3-*anti*-configurations in which the unsubstituted phenyl rings can freely rotate. Both compounds **4b** and **5b** exist in a mixture of the 1,2-alternate and the partial cone conformations, in which the carbamate moieties are in the *anti*-position, in ratios of 5.4:1 and 3.2:1, respectively. The 1,2-alternate conformations display similar patterns in the ¹H NMR spectra as **2b**, whereas the partial cones are recognized by the 2:1:1 ratio of the *t*-butyl signals and one methoxy signal. Reduction of *S*-dimethylthiocarbamate **5a** with LiAlH₄¹² in refluxing THF for 14 h under argon afforded the 1,3-dimercaptocalix[4]arene **6a** in a yield of 85%. In contrast to 1,3-dihydroxycalix[4]arene **3a**, **6a** exists as a 87:13 mixture of 1,3-*syn*- and 1,3-*anti*-configurations as followed from a corresponding ¹H NMR spectrum (SH at δ 2.50 and 2.76, respectively) as that of **3c**. Treatment of *S*-dimethylthiocarbamate **5b** with LiAlH₄ for 2 days afforded the 1,3-dimercapto-2,4-dimethoxycalix[4]arene **6b** as a complicated mixture of conformers. However, methylation of **6b** with methyl iodide in a 1:1 mixture of dichloromethane and acetonitrile in the presence of K₂CO₃ gave the 1,3-dimethoxy-2,4-bis(methylthio)calix[4]arene **6c** in 40% overall yield (from **5b**). The ¹H NMR spectrum shows the characteristic patterns¹ of a partial cone (with a methylthio group in the cavity) and a cone conformation in a ratio of 3:2. Compounds **6b** and **6c** represent the first examples of 'mixed' calix[4]arenes having two different (substituted) heteroatoms at the lower rim.

In principle both types of novel calix[4]arene derivatives **3**

|| For instance, for **5a**, two ABq patterns (ratio 1:1) are present of the methylene bridge protons at δ 4.37 and 3.79 (*J* 15.0 Hz) and 4.30 and 3.87 (*J* 14.7 Hz) and four singlets of the *t*-butyl groups (ratio 1:1:1:1) at δ 1.39, 1.36, 1.19 and 1.14.

and **6** will give rise to a variety of novel applications in calixarene chemistry. We are currently studying the (selective) replacement of the hydroxy functions in calix[4]arenes by other functionalities.

We are grateful to Dr C. S. Wilcox (Pittsburgh, USA) for suggesting to us the dehydroxylation experiments.

Received, 25th June 1990; Com. 0102845B

References

- 1 C. D. Gutsche, *Calixarenes: Monographs in Supramolecular Chemistry*, vol. 1, ed. J. F. Stoddart, RSC, Cambridge, 1989.
- 2 J.-D. van Loon, A. Arduini, W. Verboom, R. Ungaro, G. J. van Hummel, S. Harkema and D. N. Reinhoudt, *Tetrahedron Lett.*, 1989, **30**, 2681; J.-D. van Loon, A. Arduini, L. Coppi, W. Verboom, A. Pochini, R. Ungaro, S. Harkema and D. N. Reinhoudt, *J. Org. Chem.*, in the press.
- 3 P. J. Dijkstra, J. A. J. Brunink, K.-E. Bugge, D. N. Reinhoudt, S. Harkema, R. Ungaro, F. Uguzzoli and E. Ghidini, *J. Am. Chem. Soc.*, 1989, **111**, 7567.
- 4 Z. Goren and S. E. Biali, *J. Chem. Soc., Perkin Trans. 1*, 1990, 1484.
- 5 G. W. Kenner and N. R. Williams, *J. Chem. Soc.*, 1955, 522.
- 6 R. A. Rossi and J. F. Bunnett, *J. Org. Chem.*, 1973, **38**, 2314.
- 7 C. D. Gutsche, M. Iqbal and D. Stewart, *J. Org. Chem.*, 1986, **51**, 742.
- 8 A dynamic ¹H NMR study of flexible tetraalkylated calix[4]arenes shows that in these systems the 1,2-alternate conformation is a 'common' conformation: L. C. Groenen, J.-D. van Loon, W. Verboom, A. Casnati, R. Ungaro, A. Pochini, S. Harkema, F. Uguzzoli and D. N. Reinhoudt, submitted for publication.
- 9 S. G. Bott, A. W. Coleman and J. L. Atwood, *J. Inclusion Phenom.*, 1987, **5**, 747.
- 10 C. D. Gutsche, J. S. Roger, D. Stewart and K. A. See, *Pure Appl. Chem.*, 1990, **62**, 485.
- 11 M. S. Newman and H. A. Karnes, *J. Org. Chem.*, 1966, **31**, 3980.
- 12 Th. N. Sorrell and E. H. Cheesman, *Synth. Commun.*, 1981, **11**, 909.