

Chiral Bicyclic Lactams—An Asymmetric Synthesis of the Framework of the *Lycopodium* Alkaloid Magellanine Containing all Six Adjacent Stereogenic Centres

David A. Sandham and A. I. Meyers*



Department of Chemistry, Colorado State University, Fort Collins, Colorado 80523, USA

A concise bicyclic lactam-mediated asymmetric synthesis of the titled ring system is described.


A plethora of complex polycyclic alkaloids have been isolated from club mosses of the genus *Lycopodium*,¹ posing a diverse array of synthetic challenges for the organic chemist. Of particular interest are the compounds magellanine **1**,^{2a} magellaninone **2**^{2b} and paniculatine **3**^{2c,d} isolated from *L. magellanicum* and *L. paniculatum* and possessing a highly condensed tetracyclic nucleus with five or six adjacent stereogenic centres. Recent synthetic efforts have culminated in the preparation of both **1** and **2** in enantiomerically pure³ and racemic forms.^{4,5}

We now describe a conceptually different approach to this class of compounds, which further underscores the versatility of our chiral bicyclic lactam methodology for the construction of asymmetric quaternary centres.⁶ Our plan to access the stereochemically enriched framework hinged upon a recent disclosure from these laboratories⁷ on the asymmetric synthesis

of angularly-substituted hydrinden-2-ones (Scheme 1). Thus, a standard double metallation-alkylation sequence on the lactam **4** afforded a 5.9:1 ratio of diastereoisomers favouring **6**,[†] which could be isolated pure in good yield. The *endo* facial selectivity of the second alkylation is consistent with a multitude of related alkylations reported by us in recent years.^{6,7}



Scheme 1 Reagents and conditions: i, LDA, DMPU-THF, $-78\text{ }^{\circ}\text{C}$; 4-bromobutene, $-78\text{--}0\text{ }^{\circ}\text{C}$; ii, LDA, DMPU-THF, $-78\text{ }^{\circ}\text{C}$; 3-(bromomethyl)pyridine $-78\text{--}0\text{ }^{\circ}\text{C}$; iii, 9-BBN, THF, $0\text{ }^{\circ}\text{C}$; NaOH, H_2O_2 , $25\text{ }^{\circ}\text{C}$; iv, (COBr)₂, DMF, CH_2Cl_2 , $0\text{--}25\text{ }^{\circ}\text{C}$; v, KH, Bu^tLi, THF, $-78\text{ }^{\circ}\text{C}$; vi, Bu₄NH₂PO₄, THF-H₂O-CH₂Cl₂, reflux; vii, NaOEt, EtOH, $25\text{ }^{\circ}\text{C}$



Scheme 2 Reagents and conditions: i, PhOCOCl, TiCl₄, Pr₂NEt, CH₂Cl₂, $-78\text{ }^{\circ}\text{C}$; ii, DDQ, benzene $25\text{ }^{\circ}\text{C}$; iii, H₂ (1 atm), Pd/C, EtOH, $25\text{ }^{\circ}\text{C}$; iv, L-Selectride, THF, $-78\text{--}20\text{ }^{\circ}\text{C}$; v, MeI, MeCN, reflux; vi, NaBH₄, MeOH, $25\text{ }^{\circ}\text{C}$; vii, H₂ (1 atm), PtO₂, EtOH, $25\text{ }^{\circ}\text{C}$




Fig. 1 ORTEP drawing of the tetracyclic piperidine **15**

Following hydroboration–oxidation, a novel bromination protocol featuring *in situ* formation of (bromomethyl)ene)dimethyliminium bromide⁸ afforded the unstable bromolactam **8**, which was immediately submitted to the halogen-metal exchange, followed by cyclization and hydrolysis of the resultant carbinolamine **9** to remove the chiral auxiliary.⁷ Notably, the use of more standard bromination conditions (Ph₃P–NBS, *etc.*) resulted in formation of intractable mixtures of **8** and triphenylphosphine oxide. The resultant enantiomerically pure diketone **10** then underwent aldol condensation to furnish the pivotal hydrinden-2-one **11**.[†]

For the critical ring closure to assemble the tetracyclic alkaloid framework, we planned to adapt the known regioselective intermolecular 1,4-addition of enolates to activated acyloxypyridinium salts.^{9a–d} Hence, treatment of **11** with phenyl chloroformate (Scheme 2) to activate the pyridine ring towards nucleophilic attack, followed by titanium enolate formation,^{9c,10} resulted in isolation of the 1,4-dihydropyridine **12** as the major component of an inseparable 5:1 mixture of regioisomers. Intramolecular enolate addition to acylpyridinium salts are rather rare.¹¹ Oxidation¹² of the labile mixture **12**, followed by hydrogenation and separation of the pyridine isomers provided a single diastereoisomer of the tetracyclic ketopyridine **13**, the stereochemistry of which was confirmed by NOE experiments.

To set the stereochemistry of the AB ring junction, we envisaged that selective reduction of the carbonyl moiety of **13** would provide a hydroxy group amenable disposed for a directed hydrogenation¹³ of the pyridine ring. Furthermore, examination of molecular models led to the expectation that an appropriately bulky reducing agent would deliver the requisite α -carbinol. Gratifyingly, treatment of **13** with L-Selectride provided a single carbinol diastereoisomer which was quantitatively converted to the pyridinium methiodide salt. Unfortunately, the pyridinium salt proved inert to moderate pressures (3 atm) of hydrogen and recourse was made to an alternative two-stage reduction protocol. Reduction of the methiodide salt with excess NaBH₄ provided a single regioisomer of the tetrahydropyridine **14** in excellent yield, a process hitherto rarely reported with polycyclic substrates.¹⁴ Subsequent hydrogenation of **14** over Adams' catalyst (1 atm) smoothly produced a single diastereoisomer of the tetracyclic piperidine **15**,[†] whose structure was confirmed *via* a single crystal X-ray analysis (Fig. 1).[‡]

In summary, compound **15** has been prepared in 13 steps from the commercially available bicyclic lactam **4**, from which were derived all of the six stereogenic centres of magellanine.

The authors are grateful to Dr Masahiko Seki for some early experiments on related systems. We are also grateful to Susie Miller for the X-ray crystal structure determination and Tamara Joice for the NOE measurements. Financial support for this work was generously provided by the National Institutes of Health and a NATO postdoctoral fellowship (to D. A. S.) from the Engineering and Physical Sciences Research Council, UK.

Received, 29th August 1995; Com. 5105678K

Footnotes

[†] Selected physical and spectroscopic data for **6** (all coupling constants measured in Hz): [α]_D²⁵ + 72.6 (*c* 1.0, CHCl₃); δ_{H} (300 MHz, CDCl₃): 0.80 (d, *J* 6.6, 3 H), 1.01 (d, *J* 6.6, 3 H), 1.37 (s, 3 H), 1.58 (m, 2 H), 1.76 (m, 1 H), 1.89 (d, *J* 14.3, 1 H), 2.26 (d, *J* 14.3, 1 H), 2.08 (m, 2 H), 2.58 (d, *J* 13.5, 1 H), 3.00 (d, *J* 13.5, 1 H), 3.48 (m, 2 H), 3.68 (m, 1 H), 4.95 (d, *J* 10.0, 1 H), 5.03 (d, *J* 17.2, 1 H), 5.77 (ddt, *J* 17.2, 10.0, 6.5, 1 H), 7.14 (dd, *J* 7.7, 4.8, 1 H), 7.45 (dt, *J* 7.7, 2.0, 1 H), 8.40 (m, 2 H); δ_{C} (75 MHz, CDCl₃): 18.88 (q), 20.85 (q), 25.20 (q), 29.09 (t), 34.05 (d), 38.08 (t), 40.24 (t), 41.22 (t), 52.34 (s), 62.45 (s), 69.86 (t), 95.53 (s), 115.19 (t), 122.94 (d), 133.17 (s), 137.51 (d), 137.8 (d), 147.97 (d), 151.49 (d), 182.90 (s); ν_{max} /cm⁻¹ 1704, 1641; MS (EI, 70 eV): 328 (M⁺), 313, 285, 274, 243. For **11**: mp

58.5–59.5 °C; [α]_D²⁵ + 18.2 (*c* 0.72, CH₂Cl₂); δ_{H} (300 MHz, CDCl₃): 1.37–1.55 (m, 2 H), 1.70–1.89 (m, 2 H), 1.95 (d, *J* 18.6, 1 H), 2.03–2.17 (m, 2 H), 2.42 (d, *J* 18.6, 1 H), 2.57 (dt, *J* 5.4, 13.2, 1 H), 2.78 (br d, *J* 13.0, 1 H), 2.90 (ABq, *J* 14.0, $\Delta\delta_{\text{AB}}$ 9.0, 2 H), 5.81 (d, *J* 1.2, 1 H), 7.18 (dd, *J* 7.8, 4.8, 1 H), 7.39 (d, *J* 8.1, 1 H), 8.35 (d, *J* 1.5, 1 H), 8.45 (dd, *J* 4.8, 1.5, 1 H); δ_{C} (75 MHz, CDCl₃): 21.75 (t), 27.98 (t), 28.41 (t), 38.11 (t), 39.23 (t), 47.44 (s), 48.21 (t), 123.20 (d), 128.60 (d), 132.85 (s), 137.26 (d), 148.27 (d), 151.03 (d), 185.39 (s), 206.65 (s); ν_{max} /cm⁻¹ 1693, 1621, 1589; MS (EI, 70 eV): 227 (M⁺), 199, 184, 156, 135. For **15**: mp 177–178 °C (blocks from CH₂Cl₂–hexanes); [α]_D²⁶ – 10.0 (*c* 0.21, CHCl₃); δ_{H} (300 MHz, C₆D₆): 0.70 (br s, disappears on D₂O wash, 1 H), 1.12–1.64 (series of m, 13 H), 1.69 (dd, *J* 6.6, 4.5, 1 H), 1.78–1.96 (m, 4 H), 2.07–2.09 (m, 2 H), 2.13 (s, 3 H), 2.55–2.64 (m, 2 H), 3.99 (app t, *J* 5.6, 1 H); δ_{C} (75 MHz, C₆D₆): 22.38 (t), 25.11 (t), 25.97 (t), 30.31 (t), 36.65 (d), 39.21 (t), 39.88 (t), 40.93 (t), 41.29 (d), 41.57 (d), 47.50 (q), 51.97 (s), 56.45 (t), 58.04 (t), 64.15 (d), 73.14 (d); ν_{max} /cm⁻¹ 3167; MS (FAB): 250 (MH⁺), 180, 179, 111.

[‡] Crystal data for **15**: C₁₆H₂₇NO, *M* = 249.4, monoclinic, *a* = 10.891(2), *b* = 12.437(2), *c* = 11.029(2) Å, β = 106.22(3)°, *V* = 1434.4(4) Å³, space group *P*2₁, *Z* = 4, *D*_c = 1.155 Mg m⁻³, *F*(000) = 552, μ (Mo–K α) = 0.071 mm⁻¹. A clear colourless irregular shaped block of dimensions 0.60 × 0.38 × 0.26 mm was used. Data were measured on a Siemens P4/Unix diffractometer with graphite monochromated MoK α radiation (λ = 0.71073 Å) at 173 K within 3° ≤ 2 θ ≤ 55°. The structure was solved by direct methods [Siemens SHELXTL PLUS (UNIX)] on the non-hydrogen atoms refined anisotropically using full-matrix least-squares giving final *R* = 5.99% and *wR* = 7.92% for 2584 observed reflections and 324 parameters. Atomic coordinates, bond lengths and angles and thermal parameters have been deposited at the Cambridge Crystallographic Data Centre. See Information for Authors, Issue No. 1.

References

- W. A. Ayer and L. S. Trifonov, *Alkaloids*, Academic Press, 1994, **45**, 233 and references cited therein.
- (a) M. Castillo, L. A. Loyola, G. Morales, I. Singh, C. Calvo, H. L. Holland and D. B. Maclean, *Can. J. Chem.*, 1976, **54**, 2983; (b) L. A. Loyola, G. Morales and M. Castillo, *Phytochemistry*, 1979, **18**, 1721; (c) M. Castillo, G. Morales, L. A. Loyola, I. Singh, C. Calvo and H. L. Holland, *Can. J. Chem.*, 1975, **53**, 2513 (d) M. Castillo, G. Morales, L. A. Loyola, I. Singh, C. Calvo and H. L. Holland, *Can. J. Chem.*, 1976, **54**, 2900.
- G. C. Hirst, T. O. Johnson Jr. and L. E. Overman, *J. Am. Chem. Soc.*, 1993, **115**, 2992.
- L. A. Paquette, D. Friedrich, E. Pinard, J. P. Williams, D. R. St. Laurent and B. A. Roden, *J. Am. Chem. Soc.*, 1993, **115**, 4377; J. P. Williams, D. R. St. Laurent, D. Friedrich, E. Pinard, B. A. Roden and L. A. Paquette, *J. Am. Chem. Soc.*, 1994, **116**, 4689.
- Racemic approaches to the skeleton have also been reported: G. Mehta and K. S. Rao, *J. Chem. Soc., Chem. Commun.*, 1987, 1578; G. Mehta and M. S. Reddy, *Tetrahedron Lett.*, 1990, **31**, 2039; M. T. Crimmins and P. S. Watson, *Tetrahedron Lett.*, 1993, **34**, 199.
- A. I. Meyers and D. Romo, *Tetrahedron*, 1991, **47**, 9503.
- A. I. Meyers and L. B. Snyder, *Synlett*, 1991, 863; A. I. Meyers and L. B. Snyder, *J. Org. Chem.*, 1993, **58**, 507.
- M. Yoshihara, T. Eda, K. Sakaki and T. Maeshima, *Synthesis*, 1980, 746. These workers utilized the isolated iminium salt to convert primary alcohols to the corresponding bromides.
- K. Akiba, Y. Nishihara and M. Wada, *Tetrahedron Lett.*, 1983, **24**, 5269; (b) D. L. Comins and J. D. Brown, *Tetrahedron Lett.*, 1984, **25**, 3297; (c) G. Courtois, A. Al-Arnaout and L. Miginiac, *Tetrahedron Lett.*, 1985, **26**, 1027; (d) D. L. Comins and J. D. Brown, *Tetrahedron Lett.*, 1986, **27**, 2219.
- D. A. Evans, D. L. Reiger, M. T. Bilodeau and F. Urpi, *J. Am. Chem. Soc.*, 1991, **113**, 1047.
- Some examples of intramolecular reactions of acylpyridinium salts with enolates have been reported: D. D. Weller and G. R. Luellen, *Tetrahedron Lett.*, 1981, **22**, 4381; D. D. Weller, G. R. Luellen and D. L. Weller, *J. Org. Chem.*, 1983, **48**, 3061.
- M. Wada, Y. Nishihara and K. Akiba, *Tetrahedron Lett.*, 1985, **26**, 3267.
- For an excellent review of substrate-directed chemical reactions, see A. H. Hoveyda, D. A. Evans and G. C. Fu, *Chem. Rev.*, 1993, **93**, 1307.
- (a) M. Ferles, M. Kovarik and Z. Vondrackova, *Collect. Czech. Chem. Commun.*, 1965, **31**, 1348; D. S. Grierson, M. Harris and H-P. Husson, *J. Am. Chem. Soc.*, 1980, **102**, 1064; (b) A. I. Meyers and T. R. Bailey, *J. Org. Chem.*, 1986, **51**, 872.