

Tungsten-mediated Syntheses, Skeletal Rearrangement and Synthetic Applications of η^1 -2,5-Dihydro-3-furanyl Rings

Ming-Huei Cheng,^a Gene-Ming Yang,^a Jin-Feng Chow,^a Gene-Hsian Lee,^b Shie-Ming Peng^b and Rai-Shung Liu*^a

^a Department of Chemistry, National Tsing Hua University, Hsinchu 30043, R. O. China


^b Department of Chemistry, National Taiwan University, Taipei 10764, R. O. China

The syntheses, acid-catalysed rearrangements and demetallation chemistry of $\text{CpW}(\text{CO})_3(\eta^1\text{-}\overline{\text{C}=\text{CR}-\text{CHR}^1-\text{O}-\text{CH}_2})$ ($\text{R} = \text{CHCH}_2$, $\text{C}\equiv\text{CPh}$; $\text{R}^1 = \text{alkyl, phenyl}$; $\text{Cp} = \text{C}_5\text{H}_5$) are reported, and their synthetic utility in organic reactions is demonstrated.

A major goal of organometallic chemistry is to develop the synthetic utility of organometallic compounds in organic reactions.¹ Transition metal-mediated organic reactions may provide an easy and efficient synthesis of useful organic molecules which are not readily produced by common methods. In organic chemistry, dihydrofuran rings are of special interest because they are important constituents of

numerous natural products.² We describe here a novel tungsten-mediated synthesis of new classes of functionalized 2,5-dihydrofurans which provide potential utility in organic synthesis.

Although in a recent paper³ it was reported that a transition-metal- η^1 -allyl compound failed to undergo acid-catalysed cyclization with aldehydes, the reaction between the




Scheme 1

tungsten η^1 -propynyl compounds **1** and **2**⁴ and a mixture of RCHO (R = Me, Me₂CH, Me₃C, Ph, 1.2 equiv.)-BF₃·Et₂O (1.0 equiv.) in cold toluene (-40 °C, 2 h) produced the tungsten- η^1 -2,5-dihydro-3-furanyl compounds in good yields (>80%). The five-membered ring structure is supported by both the long-range coupling parameters $J_{1,2,4-5}$ and $J_{1,3,1.5-2}$ Hz and an X-ray diffraction study[†] of **5**. The mechanism of their formation is believed to follow a [3 + 2] cycloaddition⁵ involving an allene zwitterionic intermediate.


Compounds **7-9** are prone to acid-catalysed skeletal rearrangement. Treatment of **7-9** with anhydrous CF₃SO₃H (0.5 equiv.) in CH₂Cl₂ (1 h, 0 °C) afforded **10-12** in 42-25% yield, characterized as unsaturated ketone derivatives based on mass, IR, ¹H and ¹³C NMR data.[‡] In a proton NOE experiment, irradiation of the CH₂CO signal (δ 2.75) of **10** produced a 1.5% increase in the signal of =CH (δ 7.08) whereas the Me signal was unaffected; therefore the (*E*)-configuration is assigned. In contrast, no skeletal isomerization was observed for the vinyl compounds **3-6**, which gave a dark red insoluble solid of undetermined composition under the same conditions. As shown in Scheme 1, the H⁺ can be envisaged to initiate furan ring opening, regenerating the α -hydroxyallene cation **I**, which undergoes further intramolecular cyclization to give the unstable four-membered ring dihydrooxete **II**. A subsequent acid-catalysed isomerization of **II** is expected to give compounds **10-12**. Our preliminary results reveal that this skeletal rearrangement is also achieved by BF₃·Et₂O (1.0 equiv.) which nevertheless decreases the yields (**10**, 26%; **11**, 18%; **12**, 22%).

[†] Compound **5** crystallizes in the monoclinic space group $P2_1/n$, $a = 7.444(1)$, $b = 29.676(4)$, $c = 8.224(2)$ Å, $\beta = 90.63(2)^\circ$, $Z = 4$, final $R = 0.021$ and $R_w = 0.016$ for 1928 reflections with $I > 2\sigma(I)$ out of 2340 unique reflections.

[‡] Selected spectra data for complex **10**: ¹H NMR (300 MHz, CDCl₃) δ 2.02 (d, J 7.0 Hz, 3 H, Me), 2.75 (s, 2 H, CH₂CO), 5.54 (s, 5 H, C₅H₅), 7.08 (q, J 7.0 Hz, CH=CMe), 7.32 (m, 3 H, Ph-H) and 7.48 (d, 2 H, *o*-Ph-H); ¹³C NMR (75 MHz, CDCl₃) - 5.3 (W-CH₂), 16.8 (Me), 85.6, 97.2 (C≡C), 92.2 (C₅H₅), 123.3, 126.8, 127.9, 131.2 (Ph-C), 129.3, 143.8 (C=C), 204.1 (CO), 217.1 and 228.2 (2 × W-CO); IR (Nujol) $\nu(\text{CO})/\text{cm}^{-1}$ 2019s, 1920s and 1645s. For complex **23**: ¹H NMR (400 MHz, CD₃CN) δ 0.83 (d, 3 H, Me), 0.91 (d, 3 H, Me'), 1.78 (m, 1 H, CHMe₂), 2.23 (m, 1 H, H⁵), 2.68 (ddd, 1 H, H⁶), 2.79 (m, 1 H, H⁴), 3.53 (m, 1 H, H⁹), 3.64 (dd, 1 H, H⁸), 4.02 (dd, 1 H, H¹), 4.07 (dd, 1 H, H²), 4.16 (dd, 1H, H³) and 5.74 (m, 1 H, H⁷), $J_{1,7}$ 3.0, J_{1,CHMe_2} 6.8, $J_{2,3}$ 9.1, $J_{2,4}$ 6.9, $J_{3,4}$ 8.4, $J_{4,8}$ 8.3, $J_{5,6}$ 15.5, $J_{5,7}$ 6.1, $J_{5,9}$ 7.0, $J_{6,7}$ 1.6, $J_{6,9}$ 1.7, $J_{8,9}$ 8.7 and J_{CHMe_2} 6.8 Hz.



Scheme 2 M = CpW(CO)₃; i, Me₃NO·H₂O (5 equiv.), CH₂Cl₂ (23 °C, 5 h); ii, CO (1 atm), (NH₄)₂Ce(NO₃)₆ (3.0 equiv.), MeOH-CH₂Cl₂ (1:1, -78 °C, 4 h)



Scheme 3 TCNE = tetracyanoethylene, MA = maleic anhydride; i, benzene, 2 h, 25 °C, TCNE (1.2 equiv.); ii, MeCN, 80 °C, 10 h, MA (2.0 equiv.)

Demetallation of the organometallics above has been studied to explore their utility in organic reactions. As shown in Scheme 2, treatment of **4-6** with an excess of Me₃NO·H₂O[§] (5.0 equiv.) in CH₂Cl₂ gave the new dienes **13**, **15** and **17**, respectively in reasonable yields. Additional α , β -unsaturated ketones **14**, **16** and **18** were also isolated in pure forms in ca. 10% on a preparative SiO₂-TLC plate.

Similarly, the enyne **19** can be obtained from **9** in 51% yield under the same conditions. Following a known procedure,

[§] Me₃NO·H₂O was prepared by mixing equimolar amounts of anhydrous Me₃NO and H₂O. We have performed an isotopic study involving Me₃NO·D₂O in the reaction. According to ¹H and ¹³C NMR spectra, the three alkenic protons HC=C and =CH₂ were all deuterated equally, *i.e.* ca. 1/3 deuterium content, indicating that D₂O serves as proton source.

Ce^{IV}-promoted⁶ carbonylation of **5** and **9** in MeOH-CH₂Cl₂ (-78 °C, 2 h) smoothly produced the esters **20** and **21** in ca. 55% yield. The dienes **13**, **15** and **17** belong to a novel class[¶] of plane-nonsymmetric endocyclic dienes. We are particularly interested in their Diels-Alder reaction chemistry because it may provide easy access to some natural tetracyclic terpenes^{7,8}. The three dienes above react smoothly with maleic anhydride (MeCN, 80 °C, 12 h) and tetracyanoethylene (benzene, 23 °C) to give the adducts **22-26**. Notably only a single diastereoisomer was observed for TCNE or maleic anhydride adducts **22-24** of **13** and **17**. Good diastereofacial selectivities were observed for the cycloaddition of **15** with TCNE and maleic anhydride. We have carefully determined the stereochemistries of the major (single) isomer of each reaction by means of ¹H NMR NOE-difference spectra.¹¹ The results are given in Scheme 3. In the particular case of **26**, irradiation of the H⁴-signal at δ 2.98 led to an Overhauser increase in the intensities of H¹ and H⁹ by 1.9 and 1.4%, respectively. This information indicates that maleic anhydride approaches the diene *cis* to the phenyl group with its anhydride orientation *endo* to the diene fragment. In contrast, the Me₂CH and Me₃C substituents preferentially form *anti*-cycloaddition adducts **22-24** as deduced from the proton NOE-difference data. The *syn*-directed effect of the phenyl case is surprising. A possible secondary orbital interaction⁹ between the phenyl π-electrons (HOMO) and the frontier

orbitals of the dienes and dienophiles may be responsible for its stereochemistry.

Received, 25th February 1992; Com. 2/01004F

References

- 1 J. P. Collman, L. A. Hegedus, J. R. Norton and R. G. Finke, *Principles and Applications of Organotransition Metal Chemistry*, University Science Books, Mill Valley, CA, 1987; S. G. Davies, *Organotransition Metal Chemistry: Application to Organic Synthesis*, Pergamon, Oxford, 1987.
- 2 D. M. X. Donnelly and M. J. Meegan, in *Comprehensive Heterocyclic Chemistry. The Structure Reactions, Synthesis and Uses of Heterocyclic Compounds*, ed. A. R. Katritzky, C. W. Rees, C. W. Bird and G. W. H. Cheeseman, Pergamon, Oxford, 1984, vol. 4, part 3, ch. 12, p. 657.
- 3 S. C. Jiang and E. Turos, *Tetrahedron Lett.*, 1991, **32**, 4639; G. E. Agoston, M. P. Cabal and E. Turos, *Tetrahedron Lett.*, 1991, **32**, 3001.
- 4 M. H. Cheng, Y. H. Ho, G. H. Lee, S. M. Peng and R. S. Liu, *J. Chem. Soc., Chem. Commun.*, 1991, 697.
- 5 M. Rosenblum, *Acc. Chem. Res.*, 1974, **7**, 122.
- 6 K. M. Nicolas and M. Rosenblum, *J. Am. Chem. Soc.*, 1973, **95**, 4449.
- 7 R. K. Okuda, P. J. Scheuer, J. E. Hochlowski, R. J. Walker and D. J. Faulkner *J. Org. Chem.*, 1983, **48**, 1866.
- 8 J. E. Hochlowski, D. J. Faulkner, L. S. Bass and T. Clardy, *J. Org. Chem.*, 1983, **48**, 1738; P. Karuso, R. C. Cambie, B. F. Bowden and P. R. Bergquist, *J. Nat. Prod.*, 1989, **52**, 289.
- 9 P. C. Jain, Y. N. Mukerjee and N. Anana, *J. Chem. Soc., Chem. Commun.*, 1971, 303; S. Imagaki, H. Fujimoto and K. Fukui, *J. Am. Chem. Soc.*, 1976, **98**, 4054; R. Gleiter and L. A. Paquette, *Acc. Chem. Res.*, 1983, **16**, 328.
- 10 K. M. Sun, R. M. Giuliano and B. Fraser-Reid, *J. Org. Chem.*, 1985, **50**, 4774.
- 11 B. M. Trost, E. D. Edstron and M. B. Carter-Petillo, *J. Org. Chem., J. Org. Chem.*, 1989, **54**, 4489.

¶ In contrast, the derivatives of related endocyclic dienes bearing a dihydrofuran ring such as 4-vinyl-2,5-dihydrofurans¹⁰ and 3,4-bis-(methylidene)-2,5-dihydrofurans¹¹ have been recently reported, see refs. 10 and 11.