

The Fortuitous Discovery of a Synthetic Cationic Molecular Receptor System for Methanol

Billy L. Allwood,^a Luis Méndez,^b J. Fraser Stoddart,*[†] David J. Williams^a and M. Kevin Williams^b

^a Department of Chemistry, Imperial College, London SW7 2AY, UK

^b Department of Chemistry, The University, Sheffield S3 7HF, UK

A single crystal X-ray structural study of a diprotonated bis(1,7-diaza-12-crown-4), crystallised from a methanolic solution, reveals that a methanol molecule is bound to each protonated crown *via* both [O—H...N] and [N+—H...O] hydrogen bonds.

A few years ago, we synthesised¹ a range of potential receptors for urea **1**. One of the design concepts (Fig. 1) was the 1:1 complex **1·2** involving compound **2** as a potential receptor. We have now determined the crystal structure[‡] of the MeOH solvate of **2** and found that the principal interactions are (Fig. 2) intramolecular hydrogen bonds between the amide protons and the oxygen and nitrogen atoms of their adjacent 12-membered rings: it showed no propensity to bind§ urea **1**. Subsequently, we also screened some of the synthetic intermediates (Scheme 1) for their ability to complex with uronium hexafluorophosphate **1·HPF₆**. One of these intermediates—7,7'-bis-benzyl-1,1'-ethylenebis(1,7-diaza-4,10-dioxacyclododecane) **5**—was treated with **1·HPF₆** and the product was recrystallised by diffusion of light petroleum into an EtOAc—MeOH solution.

Here, we report (a) the synthesis of the bisamide **2** *via* the intermediate **5**, (b) their X-ray crystal structures and (c) provide evidence in the solid state for the ability of **5·2HPF₆** to bind *two* MeOH molecules.

Compound **5** was obtained (Scheme 1) in 72% yield from the reaction of *N*-benzyl-1,7-diaza-4,10-dioxacyclododecane² **3** and ethylene glycol bistosylate **4** with Na₂CO₃ in MeCN under reflux for 48 h. It was isolated, characterised¶ and crystallised from CHCl₃—C₅H₁₂. Compound **2** was obtained (Scheme 1) by the hydrogenolysis (H₂, Pd/C) of **5** in MeOH and subsequent condensation (Na₂CO₃, MeCN) of the product with 2,2'-dichloro-*N,N'*-meta-phenylenedimethylene




Fig. 1 The proposed 6-point binding model for the complexation of urea **1** by the bisamide **2**




Fig. 2 The structure of **2·MeOH** in the crystal. Transannular [N...N] and [O...O] distances (Å): [N(1)...N(7)], 4.61; [N(10)...N(16)], 4.48; [O(4)...O(33)], 3.53; [O(13)...O(38)], 3.60. Hydrogen bonding geometries [X—H...Y] expressed as [X...Y], [H...Y] distances (Å), [X—H...Y] angle (°): [O(43)—H...O(41)], 2.73, 1.78, 171; [N(19)—H...N(16)], 2.71, 2.21, 111; [N(19)—H...O(38)], 3.02, 2.16, 149; [N(28)—H...N(1)], 2.73, 2.24, 111; [N(28)—H...O(33)], 3.16, 2.31, 148.

[†] Present address: School of Chemistry, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK.

[‡] Crystal data for **2·MeOH**: C₃₁H₅₄N₆O₇, *M* = 622.8, triclinic, space group *P*1̄, *a* = 8.609(1), *b* = 11.804(2), *c* = 17.811(2) Å, α = 72.93(1), β = 87.60(1), γ = 76.91(1)°, *U* = 1685 Å³, *Z* = 2, *D_c* = 1.23 g cm⁻³, μ = 7 cm⁻¹.

Crystal data for **5**: C₃₂H₅₀N₄O₄, *M* = 554.8, triclinic, space group *P*1̄, *a* = 6.250(1), *b* = 10.715(2), *c* = 12.405(2) Å, α = 94.43(1), β = 99.18(1), γ = 104.66(1)°, *U* = 787.6 Å³, *Z* = 1 (the molecule is positioned about a centre of symmetry), *D_c* = 1.17 g cm⁻³, μ = 6 cm⁻¹.

Crystal data for **5·2HPF₆·2MeOH**: C₃₄H₆₀F₁₂N₄O₆P₂, *M* = 910.8, triclinic, space group *P*1̄, *a* = 10.328(2), *b* = 13.323(3), *c* = 17.004(4) Å, α = 108.19(2), β = 98.70(2), γ = 95.72(2)°, *U* = 2170 Å³, *Z* = 2, *D_c* = 1.39 g cm⁻³, μ = 18 cm⁻¹.

Data for the three structures were measured on a Nicolet R3m diffractometer with Cu-Kα radiation (graphite monochromator) using ω-scans. The structures were solved by direct methods and refined anisotropically to give for **2·MeOH**, *R* = 0.058, *R_w* = 0.063 for 2783 independent observed reflections [|*F_o*| > 3σ(|*F_o*|), 2θ ≤ 100°]; for **5**, *R* = 0.039, *R_w* = 0.056 for 1907 independent observed reflections [2θ ≤ 110°]; for **5·2HPF₆·2MeOH**, *R* = 0.060, *R_w* = 0.066 for 4366 independent observed reflections [2θ ≤ 100°]. All computations were carried out using the SHELXTL program system. Atomic coordinates, bond lengths and angles, and thermal parameters have been deposited at the Cambridge Crystallographic Data Centre. See Notice to Authors, Issue No. 1.

§ The quest for a synthetic molecular receptor for urea has been responsible for some highly innovative and inspired research in the area of supramolecular chemistry. A variety of approaches has been pursued involving (a) crown ethers (e.g. J. W. H. M. Uiterwijk, S. Harkema, D. N. Reinhoudt, K. Daasvatn, H. J. den Hertog, Jnr., and J. Geever, *Angew. Chem. Suppl.*, 1982, 1100; C. J. Staveren, V. M. L. J. Aarts, P. D. J. Grootenhuis, W. J. H. Droppers, J. van Eerden, S. Harkema and D. N. Reinhoudt, *J. Am. Chem. Soc.*, 1988, **110**, 8134); (b) transition metal-crown ether co-complexes (C. J. van Staveren, J. van Eerden, F. C. J. M. van Veggel, S. Harkema and D. N. Reinhoudt, *J. Am. Chem. Soc.*, 1988, **110**, 4994; W. F. Nijenhuis, A. R. van Doorn, A. M. Reichwein, F. de Jong and D. N. Reinhoudt, *J. Am. Chem. Soc.*, 1991, **113**, 3607); (c) acyclic polyethers (F. Vögtle, H. Sieger and W. M. Müller, *Top. Curr. Chem.*, 1981, **98**, 107), and molecular clefts (T. W. Bell and J. Liu, *J. Am. Chem. Soc.*, 1988, **110**, 3673; V. Hedge, P. Madhukar, J. D. Madura and R. P. Thummel, *J. Am. Chem. Soc.*, 1990, **112**, 4549; M. Crego, J. J. Marugán, C. Raposo, M. J. Sanz, V. Alcázar, M. C. Caballero and J. R. Morán, *Tetrahedron Lett.*, 1991, **32**, 4185).

¶ Spectroscopic data for **2**: colourless crystals, m.p. 150–152°C, (Found: *M* (EIMS), 590. Calc. for C₃₀H₅₄N₆O₆: *M*, 590). ¹H NMR (CD₂Cl₂, 250 MHz), δ 2.17–2.90 (20H, br m), 3.15–3.62 (20H, br m), 4.44 (4H, d), 7.12–7.28 (4H, m), 8.68–8.90 (2H, br t).

Spectroscopic data for **5**: colourless crystals, m.p. 73–74°C, (Found: *M* (EIMS), 554. Calc. for C₃₂H₅₀N₄O₄: *M*, 554). ¹H NMR (CDCl₃, 220 MHz), δ 2.68 (4H, s), 2.70–2.80 (16H, m), 3.54–3.72 (16H, m), 4.65 (4H, s), 7.17–7.42 (10H, m).




Fig. 3 The structure of **5** in the crystal. The transannular [N...N] and [O...O] distances are 4.30 and 4.32 Å, respectively.





Fig. 4 (a) Framework and (b) space-filling representations of the structure of **5**·2HPF₆·2MeOH in the solid state. In the space-filling representation, the two MeOH molecules are shaded. Transannular [N...N] and [O...O] distances (Å): [N(1)...N(7)], 4.60; [N(15)...N(21)], 4.49; [O(4)...O(10)], 3.02; [O(18)...O(24)], 3.08. Hydrogen-bonding geometries [X...Y] expressed as [X...Y], [H...Y] distances (Å), [X-H...Y] angle (°): [N(1)-H...O(41)], 2.78, 1.83, 169; [O(41)-H...N(7)], 2.91, 1.82, 168; [N(15)-H...O(43)], 2.73, 1.81, 162; [O(43)-H...N(21)], 2.82, 1.82, 178.

bisacetamide.¹ It was isolated, characterised[¶] and crystallised from an EtOAc-MeOH solution layered with light petroleum in an overall yield of 21%.

The solid-state structure[‡] (Fig. 3) of **5** has crystallographic C₂ symmetry and shows that both 12-membered rings adopt a [66] conformation.^{3,4} In addition to the fact that the oxygen atoms are pointing in opposite directions, the nonbonding electron pairs on the nitrogen atoms are also directed over opposite faces of the macrocycles.

The X-ray crystal structure[‡] (Fig. 4) of the product of the treatment of **5** with **1**·HPF₆ in MeOH revealed that, though no **1**·HPF₆ was present in the crystal lattice, two molecules of MeOH were intimately bound, one to each diaza-12-crown-4 macrocycle in **5**·2HPF₆. Furthermore, selective protonation was observed to have occurred in a constitutionally unsymmetrical manner within **5**. The transfer of a proton from a uronium salt to the diaza-crown nitrogen atoms is not unexpected, taking into account the difference in pK_a values⁵ between urea (ca. 0.1) and tertiary alkyl amines (ca. 10–11). The binding of each MeOH molecule is via both [O-H...N] (2.82, 2.91 Å) and [N⁺-H...O] (2.73, 2.78 Å) hydrogen bonds. An important feature of this association between neutral MeOH molecules and the monoprotonated diaza-12-crown-4 units is an appealing logical variation of the two-point binding model observed⁶ previously for primary alkyl and secondary dialkyl ammonium ions with neutral diaza-12-crown-4 systems. Whereas in the latter situation, positively charged guests are complexed by neutral hosts, in the former, the positive charge is located within the host which thence binds neutral guests.⁷ In this manner, strong association of a pole-dipole type is preserved, i.e. it is immaterial whether the charge resides in the host or in the guest. As a result of alcohol consumption the diaza-12-crown-4 macrocycles both adopt a



Scheme 1

[48] conformation³ wherein both nitrogen atoms within each macrocycle are directed over one face of their parent 12-membered ring. The oxygen atoms, on the other hand, are still essentially positioned 'one up and one down', though one CH₂OCH₂ group within each macrocycle lies close to the mean planes of their associated macrocycles. This results in transannular [O...O] distances of ca. 3 Å.

The fortuitous discovery of a synthetic cationic molecular receptor system for alcohols^{||} is a salutary reminder of

|| Different approaches have been employed for the demanding task of recognising alcohols selectively. These include the involvement of (a) clathrates (e.g. see F. Toda, K. Okada and T. C. W. Mak, *Chem. Lett.*, 1988, 1829; D. Worsch and F. Vögtle, *J. Incl. Phenom.*, 1986, 163; S. A. Bourne, L. R. Nassimbeni, K. Skobridis and E. Weber, *J. Chem. Soc., Chem. Commun.*, 1991, 282; M. Czugler and E. Weber, *J. Incl. Phenom.*, 1991, 10, 355); (b) cyclodextrins (Y. Matsui, K. Mochida, *Bull. Chem. Soc. Jpn.*, 1979, 52, 2808; A'. B. A'ri, J. Szejtli and L. Barcza, *J. Incl. Phenom.*, 1983, 1, 151); (c) cyclophanes (K. Kobiro, M. Takahashi, N. Nishikawa, K. Kakiuchi, Y. Tobe and Y. Odaira, *Tetrahedron Lett.*, 1987, 28, 3825; Y. Kikuchi, Y. Kato, Y. Tanaka, H. Toi and Y. Aoyama, *J. Am. Chem. Soc.*, 1991, 113, 1349); (d) crown ethers (E. Weber, H.-P. Josel, H. Puff and S. Franken, *J. Org. Chem.*, 1985, 50, 3125); (e) vanadium bilayers (J. W. Johnson, A. J. Jacobson, W. M. Butler, S. E. Rosenthal, J. F. Brody and J. T. Lewandowski, *J. Am. Chem. Soc.*, 1989, 111, 381); and (f) other miscellaneous receptors (B. Dung and F. Vögtle, *J. Incl. Phenom.*, 1988, 6, 429; W. Moneta, P. Baret and J.-L. Pierre, *J. Chem. Soc., Chem. Commun.*, 1985, 899; T. I. Richards, K. Layden, E. E. Warminski, P. J. Milburn and E. Haslam, *J. Chem. Soc., Perkin Trans. 1*, 1987, 2765).

Note added in proof: Since this communication was submitted for publication, novel boron-containing crown ethers (M. T. Reetz, C. M. Niemeyer and K. Harms, *Angew. Chem.*, 1991, 103, 1515; *Angew. Chem., Int. Ed. Engl.*, 1991, 30, 1472) have been shown (M. T. Reetz, C. M. Niemeyer and K. Harms, *Angew. Chem.*, 1991, 103, 1517; *Angew. Chem., Int. Ed. Engl.*, 1991, 30, 1474, to bind alcohols, such as MeOH, EtOH, PrOH, and PhCH₂OH, with good selectivities in the presence of amines like PhCH₂NH₂ or PhCHMeNH₂.

the serendipitous element which pervades supramolecular chemistry.

We acknowledge financial support from AFRC and SERC.

Received, 15th October 1991, Com. 1/05233K

References

- 1 M. K. Williams, PhD Thesis, University of Sheffield, 1986.
 - 2 M. J. Calverley and J. Dale, *J. Chem. Soc., Chem. Commun.*, 1981, 1084.
 - 3 J. Dale, *Top. Stereochem.*, 1976, **9**, 270.
 - 4 P. Groth, *Acta Chem. Scand., Sect. A*, 1978, **32**, 279.
 - 5 D. D. Perrin, *Dissociation Constants of Organic Bases in Aqueous Solution*, Butterworths, London, 1965.
 - 6 J. C. Metcalfe, J. F. Stoddart and G. Jones, *J. Am. Chem. Soc.*, 1977, **99**, 8317; J. C. Metcalfe, J. F. Stoddart, G. Jones, W. E. Hull, A. Atkinson, I. S. Kerr and D. J. Williams, *J. Chem. Soc., Chem. Commun.*, 1980, 540.
 - 7 For another example of the use of cationic receptors for the complexation of neutral guests, see B. Odell, M. V. Reddington, A. M. Z. Slawin, N. Spencer, J. F. Stoddart and D. J. Williams, *Angew. Chem., Int. Ed. Engl.*, 1988, **27**, 1547; P. R. Ashton, B. Odell, M. V. Reddington, A. M. Z. Slawin, J. F. Stoddart and D. J. Williams, *Angew. Chem. Int. Ed. Engl.*, 1988, **27**, 1550; T. T. Goodnow, M. V. Reddington, J. F. Stoddart and A. E. Kaifer, *J. Am. Chem. Soc.*, 1991, **113**, 4335; M. V. Reddington, A. M. Z. Slawin, N. Spencer, J. F. Stoddart, C. Vicent and D. J. Williams, *J. Chem. Soc., Chem. Commun.*, 1991, 630; D. Philp, A. M. Z. Slawin, N. Spencer, J. F. Stoddart and D. J. Williams, *J. Chem. Soc., Chem. Commun.*, 1991, 1584.
-