

Generation, Isolation and Photoelectron Spectrum¹ of HC≡C–N≡C

Laszlo Zanathy,^a Hans Bock,^{*a} Dieter Lentz,^{*b} Dagmar Preugschat^b and Peter Botschwina^{*c}

^a Chemische Institute der Universität, Niederurseler Hang, W-6000 Frankfurt 50, Germany


^b Institut für Anorganische Chemie, Freie Universität, Fabeckstr. 34-36, W-1000 Berlin 33, Germany

^c Institut für Physikalische Chemie der Universität, Tammannstr. 6, W-3400 Göttingen, Germany

Thermal fragmentation of pentacarbonyl(1,2-dichlorovinylisocyanide)chromium, (OC)₅Cr(CN–CIC=CHCl), yields acetylene isocyanide, a putative interstellar molecule, which, after purification by fractional condensation, is characterized by its helium(I) photoelectron spectrum.

In contrast to numerous saturated and unsaturated organic cyanides, R–CN (R = H, CH₃) and H–(C≡C)_n–CN (*n* = 1 to 5)² found in interstellar space and especially in the Taurus molecular cloud,² only a few isocyanide isomers, H–NC and H₃C–NC,² have so far been detected. But also on Earth, known nitriles by far outnumber isonitriles and, therefore, new methods for their synthesis or that of their metal complexes are welcome.³ In turn, thermal fragmentation of the latter can yield hitherto unprepared,⁴ although precalculated⁵ molecules such as the title compound.

Advantageously, acetylene isocyanide can be generated, isolated and characterized by its ionization pattern by performing the thermal fragmentation of the pentacarbonyl(1,2-dichlorovinylisocyanide)chromium complex [eqn. (1)] in a building block apparatus connected to a photoelectron spectrometer (Fig. 1), which allows the optimization of all essential reaction conditions by real-time photoelectron spectroscopic (PES) gas analysis.⁶ The chromium complex, evaporated under 10^{–4} mbar (1 bar = 10⁵ Pa) pressure at room temperature, passes through a quartz tube (length 40 cm, diameter 1.5 cm), which is heated to an oven temperature of 510 K. Except for CO, and HC≡C–N≡C, all other products of a nine hour fragmentation are frozen out in the intense cooling trap at 190 K. After closing the left-hand Teflon valve of the 78 K trap (Fig. 1) and on slow warming to 190 K, the liquid acetylene isocyanide can be evaporated into the spectrometer and its PES ionization fingerprint (Fig. 2) recorded.



The helium(I) PE spectrum recorded (Fig. 2) shows three vibrationally fine-structured bands corresponding to vertical ionizations at 11.23, 11.90 and 15.27 eV and, in addition, unresolved bands at 18.0 eV and, presumably, around 19.5 eV. The assignment to the three well-defined valence radical cation states of lowest energy, $\tilde{X}(^2\Pi)$, $\tilde{A}(^2\Sigma^+)$ and $\tilde{B}(^2\Pi)$ is straightforwardly supported by correlated Green function ionization energies^{5b} and can even be achieved applying Koopman's theorem, $IE_{i,v} = -\epsilon_j^{SCF}$, to the eigenvalue pattern from a fully geometry-optimized AM 1 calculation,⁷ with only the 5σ value exhibiting a somewhat larger defect of 1.3 eV (Fig. 2). Especially with the almost 1:1 correspondence between the observed vibrational fine structures [Fig. 2




Fig. 1 Building block apparatus assembled for the 510 K thermolysis of the pentacarbonylchromium complex, for the fractionate condensation of side products in a 190 K intense cooling trap and for the isolation of acetylene isocyanide in a 78 K trap with Teflon valves. The photoelectron spectrometer (PES) connected is used both for the evaporation of the system and for the optimization of the reaction conditions and the product analysis.

(insert)] and those predicted by anharmonic *ab initio* potential energy function calculations (Table 1), there remains hardly any doubt that the molecule generated from the chromium complex by thermal fragmentation and purified by fractionate condensation (Fig. 1) is the required acetylene isocyanide.

As concerns the other molecular properties of HC≡C–N≡C, now available on Earth, further results of the geometry-optimized AM 1 calculations (Scheme 1) are compared with those for the isomer HC≡C–C≡N: Accordingly, their semi-empirically predicted structures, which closely resemble the





Fig. 2 He I PE spectrum of HC≡C–N≡C [calibrated with ²P_{3/2} (Ar) = 15.76 eV] and its assignment based on correlated Green function ionization energies, $IE_{i,v}^{calc.}$ (GF), and on Koopman's correlation, $IE_{i,v} = -\epsilon_j^{AM1}$, with eigenvalues from a fully geometry-optimized AM 1 calculation. Insert: expanded record between 11.1 and 12.60 eV with assignment of the vibrational radical cation states observed according to anharmonic *ab initio* potential energy function calculations,^{5c} (cf. Table 1); the rather weak peak at 11.60 eV (↓) could be owing to traces of the acetylene cyanide isomer formed in the thermal fragmentation (see, text).


Table 1 Assignment of the vibrational fine structures observed for the radical cation states $\tilde{X}(^2\Pi)$ and $\tilde{A}(^2\Sigma^+)$ of HC≡C–N≡C (Fig. 1: insert) including the comparison of experimental and calculated relative intensities $I_{rel.}$ (%) as predicted by the anharmonic *ab initio* potential energy function approximation^{5c}

Assignment	Experimental		Calculated		
	$IE_{i,v}^e/eV$	$I_{rel.}$ (%)	$\Delta IE_{i,v}^e/eV$	$I_{rel.}$ (%)	
$\tilde{X}(^2\Pi)$	0 ₀ ⁰	11.23	100	0	100
	2 ₀ ¹	11.50	90	0.27	76
	2 ₀ ²	11.77	35	0.54	29
	2 ₀ ³	12.04	10	0.81	7
$\tilde{A}(^2\Sigma^+)$		11.90	80	0	80
		12.01	10	0.11	8
		12.16	5	0.27	6
		12.20	4	0.29	6

Structures



Charges



Scheme 1 Comparison of the HC₃N isomers based on selected results of fully geometry-optimized AM 1 calculations (see text)

experimental ones,^{4,8} expectedly only reverse the bond distances in the isomeric subunits CNC and CCN. The estimated difference in the heats of formation, $\Delta\Delta H_f \approx +128 \text{ kJ mol}^{-1}$ for the isocyanide isomer agrees reasonably well with the result from an SCF CI calculation,^{5a} and would yield a plausible explanation for the unassigned weak peak at 11.60 eV in the PE spectrum (Fig. 2: insert, ↓). The charge distributions calculated, expectedly reflect the difference in effective nuclear charges, $Z_{\text{eff}}(\text{N}) > Z_{\text{eff}}(\text{C})$ and, although the values for the individual centres are subject to caution, the difference in the resulting dipole moments of the two isomers approaches the experimentally (exp.) determined value of 0.8 D.^{4,10}

With reference to its considerable dipole moment,⁴ the novel molecule HC≡C–N≡C stands a reasonable chance of being detected radioastronomically in interstellar space (see below).

Our research project has been supported by the Deutsche Forschungsgemeinschaft, the Fonds der Chemischen Industrie and the Deutsche Akademische Austauschdienst. We are most grateful to K. Kawaguchi, M. Oshishi, S. Ishikawa and N. Kaifu from the Nobeyama Radio Observatory, Japan, for informing us about their unpublished results that they had succeeded in detecting the acetylene isocyanide in the TMC-1

cloud in concentrations about 20 to 60 times smaller than that of the isomer acetylene cyanide.

Received, 18th November 1991; Com. 1/058331

References

- For part 127 of the series: Photoelectron Spectra and Molecular Properties, see H. Bock, M. Kremer, M. Dolg and H.-W. Preuß, *Angew. Chem.*, 1991, **103**, 1200; *Angew. Chem., Int. Ed. Engl.*, 1991, **30**, 1186. For a review on the generation of short-lived molecules and their PES characterization see ref. 6.
- For a review see M. Winnewisser, *Chemie in unserer Zeit*, 1984, **18**, 1–16 and 54–61; cf. also J. Harju, M. Sahn, C. Henkel, T. L. Wilson, K. C. Sahn and S. R. Pottasch, *Astron. Astrophys.*, 1990, **223**, 197 or D. J. DeFrees, A. D. McLean and E. Herbst, *Astrophys. J.* 1985, **293**, 236 and references cited therein.
- Cf. e.g. W. P. Fehlhammer, F. Degel, and G. Beck, *Chem. Ber.*, 1987, **120**, 461 or W. P. Fehlhammer and G. Beck, *J. Organomet. Chem.*, 1989, **379**, 97 and references cited therein.
- For additional information see M. Krüger, H. Dreizler, D. Preugschat and D. Lentz, *Angew. Chem.*, 1991, **103**, 1674; *Angew. Chem., Int. Ed. Engl.*, 1991, **30**, 1644.
- (a) N. N. Haese and R. C. Woods, *J. Chem. Phys.*, 1980, **73**, 4521; 1981, **75**, 3160; (b) W. von Niessen, L. S. Cederbaum, J. Schirmer, G. H. F. Dierksen and W. P. Kraemer, *J. Electron Spectrosc. Relat. Phenom.*, 1982, **28**, 45; (c) P. Botschwina and P. Sebald, *J. Mol. Spectrosc.*, 1983, **100**, 1.
- Cf. the review H. Bock, B. Solouki, S. Aygen, M. Bankmann, O. Breuer, R. Dammel, J. Dörr, M. Haun, T. Hirabayashi, D. Jaculi, J. Mintzer, S. Mohmand, H. Müller, P. Rosmus, B. Roth, J. Wittmann and H. P. Wolf, *J. Mol. Structure*, 1988, **173**, 31 and references cited therein, especially H. Bock and B. Solouki, *Angew. Chem.*, 1981, **93**, 425; *Angew. Chem., Int. Ed. Engl.*, 1981, **20**, 427.
- M. J. S. Dewar, E. G. Zebisch, E. F. Healy and J. P. Stewart, *J. Am. Chem. Soc.*, 1985, **107**, 3902.
- M. D. Harmony, V. W. Laurie, R. L. Kuczkowski, R. H. Schwendeman, D. A. Ramsey, F. Lovas, W. J. Lafferty and A. G. Maki, *J. Chem. Phys. Ref. Data* 1979, **9**, 619.
- Cf. the review on photoelectron spectra of cyano compounds by H. Stafast and H. Bock, in *The Chemistry of Functional Groups, Supplement C*, ed. S. Patai and Z. Rappoport, Wiley, Chichester, 1983, pp. 137–185 and references cited therein.
- R. L. de Leon and J. S. Muenter, *J. Chem. Phys.*, 1985, **82**, 1702.