

Addition of MeMgSnBu^n_3 to Terminal Enyne; Preparation of Geometrically Pure 2-Substituted 1-Iodo-1,3-Dienes

Jun'ichi Uenishi,* Reiko Kawahama, Arifumi Tanio and Shoji Wakabayashi



Department of Chemistry, Okayama University of Science, Ridaicho Okayama, 700, Japan

The title iododienes have been prepared from 6-phenylhex-3-en-yne by the addition of methylmagnesiumtributyltin, followed by reactions of stannylmagnesium intermediate **A** with appropriate electrophiles and of the resulting alkylated stannyldiene **B** with iodine.

Palladium catalysed cross-coupling reaction of an iodoalkene with an appropriate alkenylstannane or alkenylborane is one of the common methods to prepare polyenes.¹ For construction of polyene units in terpenoid syntheses, we require 2-alkyl substituted 1-iodo-1,3-diene derivatives. Carbometallation reactions of 1-alkynes developed by Normant and Alexakis,² Negishi³ and others⁴ have been adopted for a preparation of 2-alkyl substituted 1-iodoalkenes. In this type of reaction, a terminal acetylene is regarded as a di-positively charged 1-alkene equivalent. However, introduction of an alkyl group at the 2-position is limited to simple alkyl or allyl groups, because of the non-availability of other organocopper or organoaluminium species. If a terminal acetylene could act as a di-negatively charged 1-alkene equivalent,⁵ a variety of electrophiles could be introduced on the 2-position; the Nozaki addition reaction of methylmagnesiumtributylstannane to terminal acetylenes⁶ can be used for this purpose. After the *syn* addition of the reagent to 1-alkyne, trapping by alkyl halides or aldehydes gives 2-alkyl substituted 1-stannyl-

alkenes.⁷ However, this reaction has not been used for the synthesis of 2-alkyl substituted 1,3-dienes to the best of our knowledge.† We have now applied the Nozaki addition to terminal enynes and to the formation of geometrically pure 1-iodo-2-alkyl-1,3-dienes. In particular, this reaction allows a novel preparation of cyclic β -[2-(1-iodo-1,3-dienyl)]ketones.








Methylmagnesiumtributylstannane was generated by the Nozaki's procedure which on exposure to enyne **1** (Scheme 2), in the presence of a catalytic amount of CuCN at -20°C , produced 2-methylmagnesium-1-tributylstannyldiene intermediate **A**. Treatment of **A** with a large excess of iodomethane formed 2-methyl-1-stannyldiene **B** ($\text{E} = \text{Me}$). Although non-substituted tributylstannyldienes have been used in syntheses,⁹ an attempt to isolate the stannyldiene **B** in pure form failed.‡ Acidic work up of **B** ($\text{E} = \text{Me}$) gave exclusively 2-methyl-1,3-diene in 93% yield. Quenching **B** with iodine



† Addition of $\text{Bu}^n_3\text{SnMgMe}$ to enyne substrates followed by protonation gave (*E*) and (*Z*) mixtures of 1,3-diene. See examples in ref. 8.


‡ 2-Substituted-1-tributylstannyl-1,3-dienes **B** are unstable and easily hydrolysed to give 2-substituted-1,3-dienes, they were always obtained as unseparable mixtures along with hydrolysed products.

Table 1 Reaction of 1 with electrophiles^a

Entry	Electrophile	Compound (E) ^a	Yield (%) ^b
1	Me-I	Me 2	91
2	PhCH ₂ OCH ₂ -Cl	PhCH ₂ OCH ₂ 3	65
3		HOCH ₂ CH ₂ 4	71
4			5 67
5			6 61
6			7 ^c 53

^a All the products exhibited satisfactory spectral data. ^b Isolated yields. ^c Single diastereoisomer.

after careful extraction by anhydrous hexane from the reaction mixture afforded the desired (*E,E*)-1-iodo-2-methyl-1,3-diene 2 in 91% yield. The stereochemistry of the original olefin was maintained without isomerization. Regioselectivity of the addition was exclusively *syn* and showed much more selectivity than the case of simple 1-alkynes.⁷ Reactions with other electrophiles such as benzyloxymethyl chloride, ethylene oxide gave the corresponding 2-substituted 1-iodo-1,3-dienes 3 and 4 in 65 and 71% yields, respectively. These results are summarized in Table 1. More interestingly the reactions with cyclopentenone and cyclohexenone gave the corresponding β -substituted cyclic ketones 5 and 6 in medium yields, but acyclic enones did not react with A. In the cases of substituted cyclic enones, 2-methylcyclopentenone reacted to give dienyl ketone 7 in 53% yield, while 3-methylcyclopentenone and



Scheme 3 Reagents and conditions: i, Bu₃SnMgMe, CuCN (cat.); ii, MeI; iii, I₂

3-methylcyclohexenone were inactive. The reaction of (*Z*)-enyne 8 proceeded stereospecifically and presented (*E,Z*)-1-iodo-2-methyl-1,3-diene 9 in 84% yield. No olefin isomerization occurred in the reaction process as long as the temperature was maintained at -20 °C in all the cases. § This reaction offers versatile preparation of 2-substituted 1-iodo-1,3-dienes from terminal enynes, which will be useful synthetic pieces for polyene synthesis and for Diels-Alder ring construction reactions.

We appreciate financial support by a Grant-in-Aid for Scientific Research from Ministry of Education, Science and Culture, Japan.

Received, 6th May 1993; Com. 3/02610H

References

- 1 J. K. Stille, *Angew. Chem., Int. Ed. Engl.*, 1986, **25**, 508; A. Suzuki, *Acc. Chem. Res.*, 1982, **15**, 178.
- 2 J. F. Normant and A. Alexakis, *Synthesis*, 1981, 841.
- 3 E. Negishi, *Acc. Chem. Res.*, 1987, **20**, 65.
- 4 P. Knochel, in *Comprehensive Organic Synthesis*, ed. M. F. Semmelhack, Pergamon Press, Oxford, 1991, vol. 4, p. 865.
- 5 I. Marek, A. Alexakis and J.-F. Normant, *Tetrahedron Lett.*, 1991, **32**, 6337.
- 6 J. Hibino, S. Matsubara, Y. Morizawa, K. Oshima and H. Nozaki, *Tetrahedron Lett.*, 1984, **25**, 2151.
- 7 S. Matsubara, J. Hibino, Y. Morizawa, K. Oshima and H. Nozaki, *J. Organomet. Chem.*, 1985, **285**, 163.
- 8 J. K. Stille and J. H. Simpson, *J. Am. Chem. Soc.*, 1987, **109**, 2138; R. Aksela and A. C. Oehlschlager, *Tetrahedron*, 1991, **47**, 1163.
- 9 For examples, B. H. Lipshutz and J. I. Lee, *Tetrahedron Lett.*, 1991, **32**, 7211; D. A. Evans and J. R. Gage, *J. Org. Chem.*, 1992, **57**, 1958.

§ Isomerization occurred at a higher temperature and the details will be discussed in a full account.