

Solid-state Interactions in Radicals: Crystal Structures of α - and β -10-Phenyl-5(10H)-phenazinyI

Thomas Barth,^a Basil Kanellakopoulos,^b Claus Krieger^a and Franz A. Neugebauer*^a

^a Abt. Organische Chemie, Max-Planck-Institut für medizinische Forschung, Jahnstr. 29, D-69028 Heidelberg, Germany

^b Institut für Heisse Chemie, Kernforschungszentrum, D-76021 Karlsruhe, Germany

Crystallization of 10-phenyl-5(10H)-phenazinyI affords two modifications in which the radicals are arranged in a chain pattern along the *a*-axis (α) or in radical pairs along the *c*-axis (β) giving rise to considerable spin pairing.


Solid-state properties of organic free radicals, in particular anomalies in the temperature (*T*) dependence of the static magnetic susceptibility (χ) attract considerable interest, an example being Wurster's blue radical cation salts.^{1,2} Similarly the uncharged 10-ethyl-5(10H)-phenazinyI **1**³ shows no Curie-Weiss behaviour, $\chi \cdot T$ falls off with decreasing temperature.^{1,4} A determination of its crystal structure, however, is not feasible owing to experimental difficulties in isolating **1** in a pure state. Further studies, therefore, require a more stable derivative. We report here the synthesis and the crystal structure of 10-phenyl-5(10H)-phenazinyI **2** and comment upon some of its properties.

Reaction of phenazine with phenyl lithium followed by oxidation of the intermediate 5,10-dihydro-5-phenylphenazine with silver tetrafluoroborate afforded 5-phenylphenazinium tetrafluoroborate [7% yield, m.p. 290 °C (decomp.)]. Reduction of this salt with potassium ferrocyanide in water led to **2**, which after careful recrystallization gave black needles [α -**2**, 6%, m.p. 156–157 °C (decomp.)] accompanied by very few prisms [β -**2**, m.p. 156 °C (decomp.)]. Microhydrogenation (5% Pd/BaSO₄, 0.48 mol H₂) and spin concentration deter-

minations in toluene (EPR, 298 K, 98%) confirmed the pure radical nature of **2**.

The well-resolved EPR spectrum of **2** in toluene was analysed by ENDOR[†] and ¹H NMR⁵ studies. Unexpectedly the π -SOMO (singly occupied molecular orbital) of **2** with large positive spin populations at N-5, 2,8-C (*para*) and 4,6,9a,10a-C (*ortho*) and small negative spin populations at 1,3,7,9-C (*meta*) is similar to that of diphenylaminyl.⁶ Determination of spin concentration of **2** in toluene in the 300–220 K temperature range ($c = 10^{-3}$ mol dm⁻³) clearly shows that **2** is monomeric in solution, whereas magnetic measurements in

[†] ENDOR (toluene, 260 K): $a(5\text{-N}) = 6.79$, $a(10\text{-N}) = 4.29$ G (1G = 10⁻⁴ T); (230 K) $a(2,8\text{-H}) = -3.47$, $a(4,6\text{-H}) = -2.70$, $a(\text{H}) \approx 0.21$ G. ¹H NMR (CD₂Cl₂, 295 K): $a(2',6'\text{-H}) = -0.22$, $a(3',5'\text{-H}) = +0.24$, $a(4'\text{-H}) = -0.06$ G; additionally $a(\text{H}) = +0.24$ and $a(\text{H}) = +0.13$ G are found for the hydrogens in 1,9- and 3,7-positions. All assignments are deduced from EPR results of selectively substituted derivatives (G. Peraus, unpublished results). The observed EPR spectrum (toluene, 300 K) is well simulated with the data $a(\text{N}) = 6.80$, $a(\text{N}) = 4.23$, $a(\text{H}) = 3.50$ (2H), $a(\text{H}) = 2.74$ (2H) and $a(\text{H}) = 0.25$ G (8H).



the solid state using the Faraday method indicate a radical content of only 43% at 294 K which further decreases at lower temperatures to reach a limit value of 2% at 70–4 K. The temperature dependence of χ was analysed by the triplet-singlet model⁷ yielding a singlet-triplet separation energy of $\Delta E = 1.1$ kcal mol⁻¹ (1 cal = 4.184 J). In comparison with **1** ($\Delta E = 0.9$ kcal mol⁻¹)⁴ the magnetic behaviour of solid **2** is similar. The striking spin pairing in solid **5(10H)**-phenaziny radicals points to a specific radical pair interaction which may be discernible from the crystal structure.

The molecular structures of α - and β -**2** were established by X-ray diffraction.[‡] In both modifications approximate C_{2v} symmetry is found for the molecules. Owing to spatial requirements the phenyl substituent is forced into an almost perpendicular arrangement with regard to the **5(10H)**-phenaziny plane [Fig. 1(a), torsion angle α -**2**: C(10a)–N(10)–C(1')–C(2') 93.0(1)°; β -**2**: [C(10a)–N(10)–C(1')–C(2')] 94.0(2)°]. Corresponding bond lengths and angles in α - and β -**2** show no significant deviation. As compared to phenazine⁸ (C–N 1.345 Å, C–N–C 116.6°), in the C(4a), N(5), C(5a) segment of α - and β -**2** [Fig. 1(a)] the C–N bond lengths are slightly increased to 1.356(3) Å while the C–N–C angle is unchanged at 116.9(3)°. Within the C(9a)–N(10)–C(10a) segment, however, the C–N–C angle [120.5(2)°] and the N–C bond lengths [1.391(3) Å] are significantly enlarged.

In α -**2** the molecules are packed in a chain pattern along the *a*-axis in which the phenyl substituents of adjacent molecules are alternately displaced by 180° [Fig. 1(b)]. In β -**2**, on the other hand, the radicals are arranged in pairs along the *c*-axis (Fig. 2). The stacking of the molecules in α -**2** corresponds exactly to the packing arrangement of phenazine.⁸ The interplanar distance of 3.29 Å in α -**2**, however, is found to be significantly shorter (phenazine: 3.49 Å). Furthermore, it is notable that only atoms with large spin populations [N(5), C(2), C(4), C(6), C(10a)] are involved in the shortest intermolecular contacts [Fig. 1(b)]. This holds also for β -**2** in which the radicals of each pair are related by a centre of inversion (Fig. 2). Within the radical pair the interplanar distance (3.50 Å) is larger than in α -**2** (3.29 Å).

‡ *Crystal data* for α -**2**, C₁₈H₁₃N₂: Black needles from ethyl acetate; dimensions 0.05 × 0.05 × 0.25 mm, *M* = 257.3, space group *I2/a* (No. 15), symmetry of the molecule in crystal: *C*₂; *a* = 10.226(2), *b* = 10.538(2), *c* = 12.103(2) Å, β = 97.33(2)°, *U* = 1293.5(9) Å³, *Z* = 4, *D*_c = 1.320 g cm⁻³, μ (Mo–K α) = 0.732 cm⁻¹, *F*(000) = 540 e. The intensity data were collected using an Enraf-Nonius CAD-4 circle diffractometer with graphite-monochromated Mo–K α radiation (λ = 0.71069 Å, $\theta/2\theta$ scanning technique). Total number of unique reflections 1141 ($\sin\theta/\lambda \leq 0.62$ Å⁻¹), of which 646 [*I* ≥ 3.0 σ (*I*)] were classified to be observed. The structure was solved by direct-methods. In the refinement anisotropic thermal parameters were applied for all non-hydrogen atoms and isotropic thermal parameters for the hydrogens. The final cycle of refinement converged with *R* = 0.031 and *R*_w = 0.034; max(Δ / σ) = 0.01, $\Delta\rho_{\max}$ = 0.04 e Å⁻³.

For β -**2**, C₁₈H₁₃N₂: Black prisms from ethyl acetate; dimensions 0.08 × 0.01 × 0.15 mm, *M* = 257.3, space group *P2₁/n* (No. 14); *a* = 8.798(1), *b* = 17.197(3), *c* = 9.184(1) Å, β = 109.91(1)°, *U* = 1306.6(7) Å³, *Z* = 4, *D*_c = 1.309 g cm⁻³, μ (Mo–K α) = 0.726 cm⁻¹, *F*(000) = 540 e. Total number of unique reflections 2471 ($\sin\theta/\lambda \leq 0.62$ Å⁻¹), of which 1034 [*I* ≥ 3.0 σ (*I*)] were classified to be observed. For measurement of the intensity data and solution of structure see α -**2**: *R* = 0.039 and *R*_w = 0.028; max(Δ / σ) = 0.01, $\Delta\rho_{\max}$ = 0.11 e Å⁻³.

Atomic coordinates and thermal parameters have been deposited at the Cambridge Crystallographic Data Centre. See Notice to Authors, Issue No. 1.




Fig. 1 Molecular structure of α -**2**: (a) view onto the **5(10H)**-phenaziny plane, showing the atom-labelling scheme, bond distances (Å) and angles (°, standard deviation about 1°). (b) Shortest intermolecular contacts between adjacent radicals d_1 [C(2)⋯N(5)] 3.315(2) Å and d_2 [C(4)⋯C(10a)] 3.347(2) Å.




Fig. 2 Packing diagram of β -**2** projected down the *c*-axis showing the radical pairs with shortest intermolecular contacts d_1 [C(2)⋯N(5)] 3.554(3) Å and d_2 [C(4)⋯C(10a)] 3.539(3) Å.

For the molecular structures of α - and β -**2** three findings are obvious, the unusual short interplanar distance of the 10-phenyl-5(10*H*)-phenaziny radicals in the chain of α -**2**, the formation of radical pairs in β -**2**, and the fact that the short intermolecular contacts in α - and β -**2** only occur between atoms bearing large spin populations. These results support the view that the magnetic anomalies of 5(10*H*)-phenaziny radicals are related to specific intermolecular π - π interactions⁹ which may also determine the crystal structures. Phenazine and α -**2**, however, show the same packing arrangement. Therefore further work is necessary to confirm, whether the molecular structures of **2** are an accidental result or typical for 5(10*H*)-phenaziny radicals.

This work was supported by the Deutsche Forschungsgemeinschaft.

Received, 10th June 1993; Com. 3/03359G

References

- 1 K. H. Hausser, *Z. Naturforsch., Teil A*, 1956, **11**, 20.
- 2 J. Yamauchi and H. Fujita, *Bull. Chem. Soc. Jpn.*, 1990, **63**, 2928, and references therein.
- 3 H. McIlwain, *J. Chem. Soc.*, 1937, 1704.
- 4 O. Serafimov and H. Zimmermann, *Ber. Bunsenges. Phys. Chem.*, 1972, **76**, 904, and references therein.
- 5 K. H. Hausser, H. Brunner and J. C. Jochims, *Mol. Phys.*, 1966, **10**, 253.
- 6 F. A. Neugebauer and S. Bamberger, *Chem. Ber.*, 1974, **107**, 2362.
- 7 B. Bleaney and K. D. Bowers, *Proc. R. Soc. London, Ser. A*, 1952, **214**, 451.
- 8 F. H. Herbstein and G. M. J. Schmidt, *Acta Crystallogr.*, 1955, **8**, 406; identical results were also obtained by K. Wozniak, B. Kariuki and W. Jones, *Acta Crystallogr., Sect. C*, 1991, **47**, 1113.
- 9 E. M. Kosower, *Top. Curr. Chem.*, 1983, **112**, 117; W. Geuder, S. Hünig and A. Suchy, *Tetrahedron*, 1986, **42**, 1665, and references therein.