


## Unusual Reactivity of a Luminescent Bis- $\mu$ -Sulfido Platinum(II) Dimer with Methylene Chloride. X-Ray Structural Characterization of $[\text{Pt}_2(\mu\text{-S})_2(\text{dppy})_4]$ and $[\text{Pt}(\text{dppy})_2(\text{S}_2\text{CH}_2)]$ (dppy = 2-diphenylphosphinopyridine)

Vivian Wing-Wah Yam,\* Phyllis Kok-Yan Yeung and Kung-Kai Cheung

Department of Chemistry, The University of Hong Kong, Pokfulam Road, Hong Kong

Reaction of  $[\text{Pt}(\text{dppy})_2\text{Cl}_2]$  with NaSH in the presence of triethylamine in MeCN produces a luminescent dimeric  $[\text{Pt}_2(\mu\text{-S})_2(\text{dppy})_4]$  complex **1** which is characterized by X-ray crystallography; its unusual reactivity with  $\text{CH}_2\text{Cl}_2$  to give novel monomeric  $[\text{Pt}(\text{dppy})_2(\text{S}_2\text{CH}_2)]$  **2**, identified by both NMR spectroscopy and X-ray crystallography, is described (dppy = 2-diphenylphosphinopyridine).


There has been considerable interest in the chemistry of soluble metal-sulfur complexes.<sup>1</sup> Recently, a tetranuclear  $d^{10}$  copper(I) cluster containing a  $\mu_4$ -sulfido moiety has been shown by us to exhibit rich photophysical and photochemical properties.<sup>2</sup> This, together with our extensive efforts on the design of luminescent polynuclear  $d^8$  and  $d^{10}$  metal complexes and the study of their photoreactivities with electron acceptors and halocarbons,<sup>2-4</sup> has prompted us to extend our studies to polynuclear  $d^8$  metal-sulfur systems which are especially attractive in view of their ability to function as metalloligands in the synthesis of polynuclear heterometallic aggregates.<sup>5</sup> In addition, the ability of  $d^8$  metal systems in the activation of carbon-halogen and carbon-hydrogen bonds has been well documented.<sup>5,6</sup> In this communication, we report on the synthesis, luminescent properties, and X-ray crystal structure of a dimeric  $[\text{Pt}_2(\mu\text{-S})_2(\text{dppy})_4]$  complex **1**; its unusual reactivity with  $\text{CH}_2\text{Cl}_2$  to give novel monomeric  $[\text{Pt}(\text{dppy})_2(\text{S}_2\text{CH}_2)]$  **2**, which has also been identified crystallographically, is also described (dppy = 2-diphenylphosphinopyridine).



Reaction of  $[\text{Pt}(\text{dppy})_2\text{Cl}_2]$  with NaSH in the presence of triethylamine in MeCN in an inert atmosphere of nitrogen gave the dimeric  $[\text{Pt}_2(\mu\text{-S})_2(\text{dppy})_4]$  **1** as an orange solid. Recrystallization from benzene-diethyl ether gave reddish orange crystals of **1** in ca. 55% yield;† the identity of which has been confirmed by single-crystal X-ray diffraction‡ (Fig. 1). Complex **1** shows a flat square-planar geometry with the two platinum atoms bridged by two  $\mu$ -sulfur ligands and the


terminal dppy ligands P-coordinated to the platinum centre in a monodentate fashion. This differs substantially from other related dimeric platinum-sulfido complexes such as  $[\text{Pt}_2(\mu\text{-S})_2(\text{PMe}_2\text{Ph})_4]$  which showed a hinged square-planar geometry with a dihedral angle of  $121^\circ$  (**1**, dihedral angle  $0^\circ$ ). Dihedral angles lying in the range  $131$ – $141^\circ$  have also been reported in other dimeric  $\mu$ -alkylsulfido complexes of platinum. The Pt...Pt distance of  $3.555(1)$  Å is significantly greater than the sum of the covalent radii, indicative of no formal metal-metal bond. The S...S separation of  $3.005(2)$  Å is comparable to those observed in bidentate thioether and phosphine ligands and other related sulfido complexes which are known to function as metalloligands.<sup>5,7</sup>

Dissolution of **1** in dichloromethane followed by vapour diffusion of diethyl ether gave pale yellow crystals of  $[\text{Pt}(\text{dppy})_2(\text{S}_2\text{CH}_2)]$  **2**, which has been characterized both spectroscopically§ and crystallographically.‡ Fig. 2 shows a perspective drawing of **2** with atomic numbering. Complex **2** shows a slightly distorted square-planar geometry with an S-Pt-S angle of  $76.23(5)^\circ$ . A similar S-Pt-S angle of  $74.7^\circ$  has been reported for a related platinum dithiocarbamate complex,  $[\text{Pt}(\text{S}_2\text{CF})(\text{PPh}_3)_2]^+$ .<sup>8</sup> To the best of our knowledge, this represents the first report on the isolation and the X-ray crystal structure of a saturated dithiolato four-membered platinumacycle. Five- and six-membered platinumacycles of dithiolate and diselenate are known.<sup>9</sup> A similar four-membered dithiolato




**Fig. 1** Perspective drawing of compound **1** with the atomic numbering scheme. Hydrogen atoms have been omitted for clarity. Thermal ellipsoids are shown at the 30% probability levels. Selected bond lengths (Å) and angles ( $^\circ$ ): Pt-S 2.333(1), Pt-P(1) 2.275(2), Pt-P(2) 2.279(1), Pt...Pt' 3.555(1), S...S' 3.005(2), S-Pt-P(1) 86.75(5), S-Pt-P(2) 170.05(7), P(1)-Pt-P(2) 102.98(6), S-Pt-S' 80.42(6), Pt-S-Pt' 99.58(6).

palladacycle  $[\text{Pd}(\text{PR}_3)_2(\text{S}_2\text{CHPR}_3)]^+$  has been isolated from the reaction of  $[\text{Pd}(\text{PR}_3)_3\text{H}]^+$  and  $\text{CS}_2$ .<sup>10</sup> The methylene protons in **2** have been unambiguously characterized by  $^1\text{H}$ ,  $^{13}\text{C}$  and  $^{13}\text{C}$  DEPT135 NMR spectroscopy (Fig. 3). The methylene group appears in the  $^{13}\text{C}\{^1\text{H}\}$  NMR spectrum as a singlet at  $\delta$  38.0; this signal is negative in the DEPT spectrum, indicative of an even number of hydrogen atoms. The methylene protons appear as a triplet at  $\delta$  5.54 [ $^4J(\text{P-H}) = 1.8$ ,  $^3J(^{195}\text{Pt-H}) = 42.2$  Hz] with platinum satellites in the  $^1\text{H}$  NMR spectrum. Similar chemical shifts and  $^3J(^{195}\text{Pt-H})$  values have been observed in the  $^{13}\text{C}$  and  $^1\text{H}$  NMR spectra of other related systems.<sup>9-11</sup> The reaction of **1** with dichloromethane to give **2** is rather unusual. It is interesting that the activation of the C-X bond in dichloromethane or other halocarbons by other similar square-planar  $d^8$  metal systems usually proceeds *via* an oxidative addition pathway.<sup>5,12</sup> It is likely that formation of **2** involves the attack of the nucleophilic sulfido bridges of **1** on dichloromethane which acts as the electrophile. A side product identified as  $[\text{Pt}(\text{dppy})_2\text{Cl}_2]$  has also been isolated in the reaction mixture [eqn. (1)].




The electronic absorption spectra of both **1** and **2** show low-energy absorption bands in the 350–425 nm region. Excitation of solid samples of both **1** and **2** at  $\lambda > 350$  nm shows long-lived luminescence at *ca.* 630 nm at 77 K. The photophysical data are collected in Table 1. An emission lifetime of *ca.* 10  $\mu\text{s}$  has been measured for a solid sample of **1** at room temperature. The long excited-state lifetime observed suggests that the emission is most likely associated with a spin-forbidden transition. Since both the low-energy absorption bands at *ca.* 350–425 nm and the emission at *ca.* 630 nm are only observable in the platinum sulfido complexes **1** and **2** and not in  $[\text{Pt}(\text{dppy})_2\text{Cl}_2]$ , we suggest that the origin of these low-energy transitions is not derived from the dppy intraligand (IL) or the metal-to-ligand charge



**Fig. 2** Perspective drawing of compound **2** with the atomic numbering scheme. Of the hydrogen atoms, only the ones attached to the methylene carbon atom are shown; the others have been omitted for clarity. Thermal ellipsoids are shown at the 30% probability levels. Selected bond lengths ( $\text{\AA}$ ) and angles ( $^\circ$ ): Pt-S(1) 2.314(1), Pt-S(2) 2.310(1), Pt-P(1) 2.279(1), Pt-P(2) 2.277(1), S(1)-C(1) 1.814(7), S(2)-C(1) 1.831(7), S(1)-Pt-S(2) 76.23(5), S(1)-Pt-P(1) 90.72(6), S(1)-Pt-P(2) 168.02(5), S(2)-Pt-P(1) 166.94(6), S(2)-Pt-P(2) 92.38(5), P(1)-Pt-P(2) 100.65(5), Pt-S(1)-C(1) 90.3(2), Pt-S(2)-C(1) 90.0(3), S(1)-C(1)-S(2) 103.0(3).

transfer  $[\text{d}(\text{Pt}) \rightarrow \pi^*(\text{dppy}) \text{ MLCT}]$  transitions. With the good  $\sigma$ -donating capability of the sulfido moiety, it is likely that the low-energy emitting state should arise from states bearing a



**Fig. 3** Diagram showing the  $^1\text{H}$  (a),  $^{13}\text{C}\{^1\text{H}\}$  (b), and  $^{13}\text{C}$  DEPT135 (c) NMR spectra of compound **2** in  $\text{CD}_2\text{Cl}_2$

Table 1 Photophysical data for compounds 1 and 2

Compound	Medium (T/K)	$\lambda_{em}/nm$	$\tau_o/\mu s$
1	Solid (298)	430	0.30 ± 0.04
		645	10.0 ± 1.0
	Solid (77) THF (298)	635	
		430	< 0.1
		550	0.06 ± 0.01
2	Solid (298)	460	
	Solid (77)	440, 630	
	CHCl <sub>3</sub> glass (77)	440(sh), 600	
	CH <sub>2</sub> Cl <sub>2</sub> (298)	420	< 0.1

large amount of ligand-to-metal charge transfer (LMCT) character. Similar assignments have been suggested in a tetranuclear copper(I) sulfur cluster.<sup>2</sup>

It is envisaged that the present platinum sulfido system should show interesting thermal and photochemical reactivities with electrophiles. Work is in progress to investigate the reactivity of this class of compounds with various halocarbons and electrophiles such as Lewis acids and to explore its potential as a metalloligand in the synthesis of polynuclear heterometallic complexes.

V. W. W. Y. acknowledges financial support from the Research Grants Council and The University of Hong Kong. P. K. Y. Y. acknowledges the receipt of a postgraduate studentship, administered by The University of Hong Kong.

Received, 26th September 1994; Com. 4/05849F

### Footnotes

† 1. UV-VIS (THF):  $\lambda/nm$  ( $\epsilon/dm^3 mol^{-1} cm^{-1}$ ) 350 (8000), 425 (3000); positive FAB-MS: ion cluster at  $m/z$  1506 {M}<sup>+</sup>.

‡ Crystal data: 1, triclinic, space group  $P\bar{1}$ ,  $a = 9.496(4)$ ,  $b = 12.726(2)$ ,  $c = 14.236(2)$  Å,  $\alpha = 64.93(1)$ ,  $\beta = 81.09(2)$ ,  $\gamma = 74.11(2)^\circ$ ,  $V = 1497.3(1.0)$  Å<sup>3</sup>,  $Z = 1$ ,  $D_c = 1.672$  g cm<sup>-3</sup>,  $\mu(Mo-K\alpha) = 49.3$  cm<sup>-1</sup>,  $F(000) = 740$ ;  $R = 0.026$  for 3373 reflections with  $I \geq 3.0\sigma(I)$ . 2, triclinic, space group  $P\bar{1}$ ,  $a = 10.303(1)$ ,  $b = 12.882(1)$ ,  $c = 14.113(2)$  Å,  $\alpha = 101.33(2)$ ,  $\beta = 102.15(2)$ ,  $\gamma = 112.56(2)^\circ$ ,  $V = 1609.3(1.0)$  Å<sup>3</sup>,  $Z = 2$ ,  $D_c = 1.650$  g cm<sup>-3</sup>,  $\mu(Mo-K\alpha) = 46.5$  cm<sup>-1</sup>,  $F(000) = 788$ ;  $R = 0.027$  for 4504 reflections with  $I \geq 3.0\sigma(I)$ . For both structures, Enraf-Nonius CAD-4 diffractometer with graphite-monochromated Mo-K $\alpha$  radiation ( $\lambda = 0.71073$  Å);  $\omega$ -2 $\theta$  scans. The structures were solved by Patterson and Fourier methods and subsequent refinement by full-matrix least squares analysis using the Enraf-Nonius SDP Programs on a Micro Vax II computer. Atomic coordinates, bond lengths and angles, and thermal parameters have been deposited at the Cambridge Crystallographic Data Centre. See Information for Authors, Issue No. 1.

§ 2. UV-VIS (CH<sub>2</sub>Cl<sub>2</sub>):  $\lambda/nm$  ( $\epsilon/dm^3 mol^{-1} cm^{-1}$ ) 373 (405); positive FAB-MS: ion cluster at  $m/z$  817 {M + H<sub>2</sub>O}<sup>+</sup>, 799 {M}<sup>+</sup>; <sup>1</sup>H NMR (270 MHz, CD<sub>2</sub>Cl<sub>2</sub>, 298 K):  $\delta$  5.54 [t, 2H, <sup>4</sup>J(P-H) 1.8, <sup>3</sup>J(Pt-H) 42.2 Hz, -CH<sub>2</sub>], 7.0-8.5 (m, 28H, dppy); <sup>13</sup>C{<sup>1</sup>H} NMR (67.9 MHz, CD<sub>2</sub>Cl<sub>2</sub>, 298 K):  $\delta$  38.0 (-CH<sub>2</sub>, negative in the DEPT spectrum), 124-150 (dppy, positive in the DEPT spectrum); <sup>31</sup>P NMR (109.4 MHz, CD<sub>2</sub>Cl<sub>2</sub>, 298 K):  $\delta$  20.8 [s, J(P-Pt) 2910 Hz].

### References

- 1 T. Shibahara, *Coord. Chem. Rev.*, 1993, **123**, 73; A. Müller and B. Krebs, *Sulfur: Its Significance for Chemistry, for the Geo-, Bio- and Cosmosphere and Technology*, Elsevier, Amsterdam, 1984; R. H. Holm, S. Ciurli and J. A. Weigel, *Prog. Inorg. Chem.*, 1991, **38**, 1; A. Müller and E. Diemann, *Comprehensive Coordination Chemistry*, ed. G. Wilkinson, Pergamon, Oxford, 1987, vol. 2, pp. 515-550.
- 2 V. W. W. Yam, W. K. Lee and T. F. Lai, *J. Chem. Soc., Chem. Commun.*, 1993, 1571.
- 3 V. W. W. Yam, L. P. Chan and T. F. Lai, *Organometallics*, 1993, **12**, 2197; V. W. W. Yam, L. P. Chan and T. F. Lai, *J. Chem. Soc., Dalton Trans.*, 1993, 2075.
- 4 V. W. W. Yam, W. K. Lee and T. F. Lai, *Organometallics*, 1993, **12**, 2383; V. W. W. Yam and W. K. Lee, *J. Chem. Soc., Dalton Trans.*, 1993, 2097; V. W. W. Yam, T. F. Lai and C. M. Che, *J. Chem. Soc., Dalton Trans.*, 1990, 3747; V. W. W. Yam and S. W. K. Choi, *J. Chem. Soc., Dalton Trans.*, 1994, 2057.
- 5 See for example, J. Chatt and D. M. P. Mingos, *J. Chem. Soc. A*, 1970, 1243; R. Ugo, G. La Monica, S. Cenini, A. Segre and F. Conti, *J. Chem. Soc. A*, 1971, 522; C. E. Briant, T. S. A. Hor, N. D. Howells and D. M. P. Mingos, *J. Chem. Soc., Chem. Commun.*, 1983, 1118; C. E. Briant, C. J. Gardner, T. S. A. Hor, N. D. Howells and D. M. P. Mingos, *J. Chem. Soc., Dalton Trans.*, 1984, 2645; C. E. Briant, D. I. Gilmour, M. A. Luke and D. M. P. Mingos, *J. Chem. Soc., Dalton Trans.*, 1985, 851; W. Bos, J. J. Bour, P. P. J. Schlebos, P. Hageman, W. P. Bosman, J. M. M. Smits, J. A. C. van Wietmarschen and P. T. Beurskens, *Inorg. Chim. Acta*, 1986, **119**, 141; D. I. Gilmour, M. A. Luke and D. M. P. Mingos, *J. Chem. Soc., Dalton Trans.*, 1987, 335; M. Zhou, Y. Xu, L. L. Koh, K. F. Mok, P. H. Leung and T. S. A. Hor, *Inorg. Chem.*, 1993, **32**, 1875; M. Zhou, Y. Xu, C. F. Lam, P. H. Leung, L. L. Koh, K. F. Mok and T. S. A. Hor, *Inorg. Chem.*, 1994, **33**, 1572; R. H. Summerville and R. Hoffmann, *J. Am. Chem. Soc.*, 1976, **98**, 7240.
- 6 J. P. Collman, L. S. Hegeudus, J. R. Norton and R. G. Finke, *Principles and Applications of Organotransition Metal Chemistry*, University Science Books, Mill Valley, 1987.
- 7 D. Seyferth, R. S. Henderson and L. C. Song, *Organometallics*, 1982, **1**, 125.
- 8 J. A. Evans, M. J. Hacker, R. D. W. Kemmitt, D. R. Russell and J. Stocks, *J. Chem. Soc., Chem. Commun.*, 1972, 72.
- 9 See for example, P. Haake and P. C. Turley, *J. Am. Chem. Soc.*, 1967, **89**, 4611; M. Schmidt and G. G. Hoffmann, *J. Organomet. Chem.*, 1977, **124**, C5; *Chem. Ber.*, 1979, **112**, 2190.
- 10 H. Werner and W. Bertleff, *Chem. Ber.*, 1980, **113**, 267; *Inorg. Chim. Acta*, 1980, **43**, 199.
- 11 W. McFarlane and R. F. M. White, *Chem. Commun.*, 1969, 439; P. C. Turley and P. Haake, *J. Am. Chem. Soc.*, 1967, **89**, 4617.
- 12 M. P. Brown, R. J. Puddephatt and C. E. E. Upton, *J. Chem. Soc., Dalton Trans.*, 1976, 2490.