

## A Novel Receptor for Amino Acid Derivatives

Glen J. Pernía,<sup>a</sup> Jeremy D. Kilburn<sup>\*a</sup> and Michael Rowley<sup>b</sup>

<sup>a</sup> Department of Chemistry, University of Southampton, Southampton, UK SO17 1BJ

<sup>b</sup> Department of Medicinal Chemistry, Merck Sharp and Dohme, Terlings Park, Harlow, Essex, UK CM20 2QR

A novel macrobicyclic featuring a thiourea as a carboxylate binding site, and amide functionality to provide further hydrogen-bonding interactions with suitable guests is prepared; the ability of this novel macrobicyclic to bind simple amino acid derivatives is investigated.

Considerable progress has been made in recent years in the design and synthesis of novel receptors for a range of organic substrates.<sup>1</sup> As part of our own efforts to develop novel receptors,<sup>2</sup> we set out to prepare a range of macrobicyclics that feature a specific binding site for carboxylate functionality at the base of the cavity (represented schematically in Fig. 1). By incorporating additional amide functionality around the rim of such a macrobicyclic structure, we hoped to provide further hydrogen-bonding sites, suitably preorganised to interact with guests, such as amino acid derivatives, bound within the macrobicyclic cavity. In this paper, we describe the straightforward synthesis of such a macrobicyclic **6**, and report on preliminary binding studies of **6** with simple amino acid derivatives.

The carboxylate binding site in **6** is provided by a thiourea moiety, which has been shown to provide a strong binding site for tetraalkylammonium carboxylates, even in relatively competitive solvents such as Me<sub>2</sub>SO.<sup>3</sup> The thiourea also has the potential to be converted into a guanidinium, which could again serve as a carboxylate binding site. A biaryl methane unit forms the rigidifying part of the rim of the macrobicyclic, and chirality and amide functionality are introduced *via* two lysine derivatives.

The synthesis of receptor **6** proved to be relatively straightforward (Scheme 1). A palladium-mediated Suzuki coupling<sup>4</sup> of bromide **1** with boronic acid **2** gave the nitrile ester **3**. Reduction of **3** with borane–methyl sulfide complex gave the corresponding amine which was coupled to *N*<sup>α</sup>-*tert*-butoxycarbonyl-*N*<sup>ε</sup>-benzyloxycarbonyl lysine. Selective removal of the benzyloxycarbonyl protecting group then gave amine **4**, which was converted in two steps to the thiourea **5**, *via* the isothiocyanate. Hydrolysis of the methyl esters of **5**, conversion to the corresponding bis(pentafluorophenyl) ester, removal of the *tert*-butoxycarbonyl protecting groups and, finally, cyclisation by slow addition of the bis(TFA salt) to a refluxing solution of diisopropylethylamine in acetonitrile, gave receptor **6**† in *ca.* 30% yield from the diester **5**. The synthesis is short and reasonably efficient, and provides considerable scope for variation of the building blocks, which should allow access to a range of related macrobicyclics.

Binding studies with receptor **6** were carried out with various acylated amino acids<sup>5</sup> as the tetrabutylammonium salts (see Table 1). Initially, we tried to determine binding constants by conventional NMR titration experiments, with deuteriochloroform as solvent.<sup>6</sup> This approach was hindered by the fact that the various NH signals we tried to monitor during the titration shifted into the aromatic region of the NMR spectrum.‡ Instead,





Fig. 1 CBS = carboxylate binding site  
RS = rigid spacer

Fig. 1 CBS = carboxylate binding site, RS = rigid spacer

we were able to determine binding constants by extracting the guests from aqueous solution into a chloroform solution of the receptor, by analogy with the picrate extraction method developed by Cram.<sup>7</sup> Thus, the tetrabutylammonium salts were partitioned between water and chloroform, both in the absence of receptor **6**, and with a known quantity of receptor **6** in the chloroform layer. Aliquots of the aqueous and chloroform layers were taken and the quantities of the tetrabutylammonium salts present in each aliquot were determined by integration of signals for the tetrabutylammonium salts in the <sup>1</sup>H NMR relative to a known quantity of dioxane. Partition and extraction



**Scheme 1** Reagents and conditions: i, Pd(PPh<sub>3</sub>)<sub>4</sub>, Na<sub>2</sub>CO<sub>3</sub>, DME; ii, BH<sub>3</sub>·SMe<sub>2</sub>, THF; iii, *N*<sup>α</sup>-*tert*-butoxycarbonyl-*N*<sup>ε</sup>-benzyloxycarbonyl-L-lysine, 1-hydroxybenzotriazole hydrate, DCC, THF; iv, H<sub>2</sub>, Pd, C; v, thiophosgene, K<sub>2</sub>CO<sub>3</sub>, CHCl<sub>3</sub>, reflux; vi, **4**, pyridine, reflux; vii, LiOH, H<sub>2</sub>O, dioxane; viii, C<sub>6</sub>F<sub>5</sub>OH, 1-hydroxybenzotriazole hydrate, DCC, cat. DMAP, THF; ix, TFA, CH<sub>2</sub>Cl<sub>2</sub>, (1 : 1, v/v); x, syringe pump addition of the bis(TFA salt) to a refluxing solution of diisopropylethylamine (3.0 equiv.) in MeCN, 7 mmol dm<sup>-3</sup> final concentration

**Table 1** Association constants for receptor **6** with various tetrabutylammonium carboxylates in CDCl<sub>3</sub>

Substrate (tetrabutylammonium salt)	K <sub>d</sub> /mol <sup>-1</sup>
<i>N</i> -Ac-Glycine	68 600
<i>N</i> -Ac-L-Alanine	16 900
<i>N</i> -Ac-D-Alanine	14 600
<i>N</i> -Ac-L-Phenylalanine	22 000
<i>N</i> -Ac-D-Phenylalanine	13 300
<i>N</i> -Ac-L-Glutamine	11 100
<i>N</i> -Ac-L-Asparagine	9 600
<i>N</i> <sup>α</sup> -Ac-L-Histidine	5 800
<i>N</i> <sup>α</sup> -Ac-L-Lysine	130 000

coefficients for each guest could thus be determined allowing calculation of association constants following the analysis described by Cram<sup>7</sup> and elsewhere.<sup>8</sup> This analysis assumes 1 : 1 complexation, which was confirmed by a Job plot<sup>8</sup> for the L-phenylalanine derivative, and assumes that none of the receptor **6** is extracted from the chloroform layer into the aqueous layer. None of receptor **6** could be detected in the <sup>1</sup>H NMR spectra of the aqueous layers. The partition and extraction experiments were each carried out five times to obtain consistent results, and dilution experiments confirmed that neither the receptor **6**, nor any of the guests, were dimerising or aggregating to a significant extent at the concentrations used in the binding experiments.

From the results (Table 1) it is clear that the receptor binds simple acylated amino acids, and binding is dominated by the carboxylate–thiourea interaction. The glycine derivative is strongly bound, but binding is reduced for derivatives of alanine, phenylalanine, asparagine, histidine and glutamine suggesting that the amino acid sidechains are not well tolerated, presumably for steric reasons. The lysine derivative, however, is particularly strongly bound, suggesting a favourable interaction between the free amine residue and the macrobicycle, which overrides any steric problems. Inspection of the <sup>1</sup>H NMR spectra of 1 : 1 mixtures (10 mmol dm<sup>-3</sup> in CDCl<sub>3</sub>) of receptor **6** and the various amino acid derivatives, revealed significant shifts of several proton signals with all of the three NH signals moving downfield by > 1 ppm, and in the case of the thiourea NH, by as much as 2.4 ppm. While there is no significant binding selectivity for the two alanine or phenylalanine enantiomers, the formation of diastereomeric complexes with these substrates is evidenced by significant differences in the <sup>1</sup>H NMR of 1 : 1 complexes of receptor **6** and substrate, depending on the amino acid enantiomer being complexed. For the 1 : 1 complex of **6** with L-alanine, for example, the alanine methyl sidechain gives a signal at δ -0.01 (1.43 unbound), the signal for the alanine α proton appears at δ 1.91 (4.44 unbound) and that for the alanine NH at δ 5.15 (δ 7.10 unbound). Such significant upfield shifts, relative to those of the unbound substrate, clearly indicate that the L-alanine substrate is bound within the cavity as anticipated, with the α proton and methyl sidechain considerably shielded by the aromatic side walls of the receptor. For the 1 : 1 complex of **6** with D-alanine the alanine methyl sidechain gives a signal at δ 1.12 the signal for the alanine α proton appears at δ 4.21 and that for the alanine NH at δ 6.40. Clearly the bound D-alanine substrate is in a substantially different environment compared to the bound L-alanine substrate. Similarly, the 1 : 1 complex of **6** with the L-phenylalanine substrate showed dramatic upfield shifts of the signals for the benzylic, α and NH protons, as well as for the acetoxy methyl protons, of the L-phenylalanine substrate, while the D-phenylalanine substrate did not. In addition, intermolecular NOEs, from the α and benzylic protons of the L-phenylalanine substrate to the aromatic protons of the biaryl-methane unit of the macrobicycle, place this substrate within the cavity of the macrobicyclic receptor. While the clearly different modes of binding of L and D substrates do not result in significantly enantioselective binding, it suggests that a fuller understanding of the interactions involved in the diastereomeric complex formation should allow the rational design of an enantioselective receptor, by incorporating features into the receptor which promote or block one of the binding modes.

In conclusion, we have developed a straightforward synthesis of a macrobicyclic receptor, which should allow for consider-

able structural variation in the future. Preliminary binding results indicate that the receptor can bind carboxylate salts as anticipated, with moderate selectivity for some amino acid derivatives.

We thank Merck, Sharp and Dohme and the SERC for a CASE award to G. J. P., and we thank Dr Russell Mortishire-Smith (MSD) for acquisition of 500 MHz NMR spectra and helpful suggestions regarding assignments of the 1 : 1 complexes described above.

Received, 2nd November 1994; Com. 4/06707J

## Footnotes

† Selected data for **6**: δ<sub>H</sub> (CDCl<sub>3</sub>, 360 MHz) 7.60 (4H, d, J 8 Hz, ArH), 7.11 (4H, d, J 8 Hz, ArH), 7.05 (8H, d, J 8 Hz, ArH), 6.91 (2H, bs, NH), 6.56 (2H, bs, NHCS), 6.43 (2H, d, J 7 Hz, NH), 4.77 (2H, dd, J 8 and 14 Hz), 4.48 (2H, d, J 6 and 14 Hz), 4.0–3.8 (8H, m), 3.05 (2H, m), 2.26 (2H, m), 1.80 (2H, m), 1.37 (4H, m), 1.05–0.95 (4H, m); δ<sub>C</sub> (CDCl<sub>3</sub>) 183.0 (0), 171.6 (0), 168.5 (0), 146.3 (0), 140.1 (0), 136.7 (0), 131.7 (0), 129.6 (1), 129.2 (1), 128.9 (1), 127.8 (1), 54.4 (1), 43.6 (2), 42.9 (2), 41.6 (2), 30.5 (2), 30.1 (2), 20.8 (2); m/z (FAB) (M + H)<sup>+</sup> 745 (Found: M<sup>+</sup> 744.3477. C<sub>43</sub>H<sub>48</sub>N<sub>6</sub>O<sub>4</sub>S requires 744.34578).

‡ It was possible to monitor various CH signals during the titration, but chemical shift changes (Δδ ca. 0.2 ppm) were smaller than for the NH signals, and resolution of the multiplets was troublesome. Resolution problems also meant that the lower limit for the concentration of the host in the NMR experiments was of the order of 5 × 10<sup>-3</sup> mol dm<sup>-3</sup>. Because of this titration curves approached the 'infinite' limit with Weber's *p* value,<sup>6,9</sup> assuming *K*<sub>a</sub> values of > 10<sup>4</sup> mol<sup>-1</sup> (see Table 1), always being > 0.8, and no useful data could be extracted from these NMR experiments.

## References

- For recent reviews on molecular recognition see: J. D. Kilburn and H. K. Patel, *Contemp. Org. Synth.*, 1994, **1**, 259; T. H. Webb and C. S. Wilcox, *Chem. Soc. Rev.*, 1993, 383; C. Seel and F. Vogtle, *Angew. Chem., Int. Ed. Engl.*, 1992, **31**, 528; J. Rebek, Jr., *Acc. Chem. Res.*, 1990, **23**, 399.
- S. S. Flack, J.-L. Chaumette, J. D. Kilburn, G. J. Langley and M. Webster, *J. Chem. Soc., Chem. Commun.*, 1993, 399; H. K. Patel, J. D. Kilburn, G. J. Langley, P. Edwards, T. Mitchell and R. Southgate, *Tetrahedron Lett.*, 1994, **35**, 481.
- E. Fan, S. A. Van Arman, S. Kincaid and A. D. Hamilton, *J. Am. Chem. Soc.*, 1993, **115**, 369; P. J. Smith, M. V. Reddington and C. S. Wilcox, *Tetrahedron Lett.*, 1992, **33**, 6085.
- A. Suzuki, *Acc. Chem. Res.*, 1982, **15**, 178; A. R. Martin and Y. Yang, *Acta Chem. Scand.*, 1993, **47**, 221.
- For other approaches to the binding of amino acid and peptide derivatives see: ref. 1; S. S. Yoon and W. C. Still, *Tetrahedron Lett.*, 1994, **35**, 2117; A. Borchardt and W. C. Still, *J. Am. Chem. Soc.*, 1994, **116**, 373; M. F. Cristofaro and A. R. Chamberlin, *J. Am. Chem. Soc.*, 1994, **116**, 5089 and refs therein; H.-J. Schneider, *Angew. Chem., Int. Ed. Engl.*, 1993, **32**, 848 and refs therein.
- For a discussion of the consequences for measuring binding constants by NMR see C. S. Wilcox, in *Frontiers in Supramolecular Organic Chemistry and Photochemistry*, ed. H.-J. Schneider and H. Durr, VCH, Weinheim, 1991.
- E. P. Kyba, R. C. Helgeson, K. Madan, G. W. Gokel, T. L. Tarnowski, S. S. Moore and D. J. Cram, *J. Am. Chem. Soc.*, 1977, **99**, 2564.
- Binding Constants, The Measurement of Molecular Complex Stability*, K. A. Connors, Wiley, 1987.
- G. Weber, in *Molecular Biophysics*, ed. B. Pullman and M. Weissbluth, Academic, 1965.