

A Novel Dinuclear Ruthenium–1,8-Naphthyridine Catalyst for the Oxidation of Alcohols and the Epoxidation of Alkenes

Alexandra E. M. Boelrijk,^a Maaïke M. van Velzen,^a Thomas X. Neenan,^{†a} Jan Reedijk,^{*a} Huub Kooijman^b and Anthony L. Spek^{*b}

^a Leiden Institute of Chemistry, Gorlaeus Laboratories, Leiden University, P. O. Box 9502, 2300 RA Leiden, The Netherlands

^b Bijvoet Center for Biomolecular Research, Crystal and Structural Chemistry, Utrecht University, Padualaan 8, 3584 CH Utrecht, The Netherlands

A novel ruthenium–1,8-naphthyridine complex is synthesized, characterized by X-ray analysis, and proven to be an active and stable catalyst in the oxidation of several alcoholic substrates and in the epoxidation of *trans*- and *cis*-stilbene and cyclooctene.

When an alcoholic group is part of a polyfunctional molecule or of a molecule that is sensitive towards acidic or alkaline reagents, the choice of effective and selective oxidants is rapidly narrowed.¹ Accordingly, research leading to new mild and selective oxidation and epoxidation systems is still an important area. Several catalytic epoxidation systems are known using RuCl₃ with bipyridyl or substituted phenanthrolines.² In addition to mononuclear ruthenium polypyridyl complexes,³ dinuclear complexes are also of interest as catalytic oxidants.⁴ Oxo-bridged ruthenium dimers with 2,2'-bipyridyl ligands have received considerable attention as oxidants, due to their enhanced thermodynamic oxidation strength, which leads to accelerated rates of oxidation of organic compounds, or to oxidations that are otherwise unobservable with monomeric analogues.⁴ However, due to their instability they have not been effective under catalytic conditions.⁵

In this study 1,8-naphthyridine (napy), a potentially dinucleating ligand of the carboxylate type,⁶ is used as a ligand, bridging the two ruthenium centres thereby stabilizing the bridge already formed by chloride, oxygen or hydroxide.

The crystal structure of a novel dinuclear ruthenium(III) complex, [Ru₂(napy)₂(H₂O)₄Cl(OH)]ClO₄ **1**, and its use as a catalyst for the oxidation of alcohols and for the epoxidation of alkenes is presented. The complex is highly soluble and catalytically active in the non-toxic solvent water, which provides for its facile recovery from organic reaction products by simple phase separation.

Compound **1** was synthesized from its chloride analogue,[‡] [Ru₂(napy)₂Cl₄] **2**, by chloride replacement using an excess of AgClO₄ in acetone–water. [Ru₂(napy)₂Cl₄] was in turn synthesized from commercial hydrated 'RuCl₃' and naphthyridine in methanol.

X-Ray structure analysis§ (Fig. 1) showed that the complex is a dinuclear Ru^{III}/Ru^{III} complex with the two naphthyridine ligands in a *trans* configuration. The bridging moieties consist of Cl[−] and OH[−]⋯H₂O ions, disordered over the crystallographic inversion centre in ratio 1 : 1.

The catalytic results are summarized in Tables 1 and 2.¶ It is evident that the dinuclear complex is an active and stable catalyst for the oxidation of primary alcohols by NaBrO₃ (reaching from 520 for octanol up to 980 turnovers for *n*-butanol in 15 h) and for the oxidation of 1,2-cyclohexanediol yielding adipic acid (1000 turnovers in 15 h). Also secondary alcohols such as cyclohexanol and cyclobutanol are easily oxidized. Cyclobutanol is oxidized to cyclobutanone and propanoic acid, suggesting that both a two- and a one-electron oxidation step are present.⁷ The mononuclear complex *cis*-[Ru(bpy)₂Cl₂]⁸ (which can be regarded as a mononuclear analogue for the dinuclear ruthenium complex) appears to be much less active under the same reaction conditions yielding only a small amount of cyclohexanemethanoic acid from cyclohexylmethanol; cyclohexanol proved to be not reactive at all, clearly showing the advantage of the dinuclear complex over the mononuclear complex. It is quite difficult to compare these results with alcohol oxidation data obtained with other complexes in the

literature, because of the difference in substrates, co-oxidants, solvents and reaction conditions. However, some relevant literature examples are added to the data in Table 1. Complex **1** was also tested for its catalytic properties in the epoxidation of *trans*- and *cis*-stilbene and cyclooctene. Compared to the previously reported highly active catalyst [RuO₂(bpy)-{IO₃(OH)₃]}(H₂O)_{1.5},⁹ our complex is less selective when NaIO₄ is used as a co-oxidant. However, when dioxygen/butyraldehyde, a cheaper and a more environmentally acceptable co-oxidant, is used, much better reaction selectivities (up to 100% for 1,2-epoxycyclooctane), albeit with lower activities for the epoxidation reaction, were obtained. The mechanism of epoxidation is generally believed to proceed then by the autoxidation of butyraldehyde to peracids and alkylhydroperoxides, which can then be used as oxygen-transfer agents in the epoxidation of alkenes.¹⁰

The absorption spectrum of **1** in aqueous solution is characterized by a visible band at 394 nm which is characteristic of metal-to-ligand charge transfer (MLCT) d_π–p_π* transitions. A spectrophotometric titration|| of **1** with Ce^{IV} in 0.1 mol dm^{−3}




Fig. 1 Molecular structure of the dinuclear ruthenium complex **1**. Only one disorder component is shown. Hydrogen atoms and non-coordinated water molecules and perchlorate anions have been omitted for clarity. Selected bond distances (Å) and angles (°): Ru(1)–Ru(1') 2.4978(13), Ru(1)–O(1) 2.048(7), Ru(1)–O(2) 2.068(8), Ru(1)–O(3) 1.81(2), Ru(1)–Cl(3) 2.292(7), Ru(1)–N(11) 2.071(9), Cl(3)–Ru(1)–N(11) 87.7(3), O(3)–Ru(1)–N(11) 93.1(8), O(1)–Ru(1)–N(11) 88.2(3), O(2)–Ru(1)–N(11) 94.3(3), O(1)–Ru(1)–O(2) 83.5(3), O(3)–Ru(1)–Cl(3) 105.1(8). (Primed atoms related by symmetry operation $-x, -y, -z$.)

Table 1 Oxidation of alcoholic substrates, catalysed by $[\text{Ru}_2(\text{napy})_2(\text{OH})\text{Cl}(\text{H}_2\text{O})_4][\text{ClO}_4]_4$ **1** and some reference compounds from the literature

Substrate	Reaction conditions	Turnovers (turnover h^{-1})	Product(s) (% Yield)
Cyclohexanol ^a	15 h (60 °C)	410 (27)	cyclohexanone (41)
Cyclohexanol ^b	2 h (room temp.)	62 (31)	cyclohexanone (75) ⁹
Cyclohexanol ^c	15 h (60 °C)	0	
Octanol ^a	15 h (60 °C)	520 (35)	octanoic acid (52)
Cyclohexylmethanol ^a	15 h (60 °C)	950 (63)	cyclohexanemethanoic acid (95)
Cyclohexylmethanol ^c	15 h (60 °C)	210 (14)	cyclohexanemethanoic acid (21)
<i>n</i> -Butanol ^a	15 h (60 °C)	980 (65)	<i>n</i> -butanoic acid (74) propanoic acid (24)
1,2-Cyclohexanediol ^a	15 h (60 °C)	1000 (67)	adipic acid (100)
Benzylic alcohol ^a	4 h (60 °C)	910 (227)	benzoic acid (85) benzaldehyde (6)
Benzylic alcohol ^d	4 h (room temp.)	49 (12)	benzoic acid (72) ¹⁰
Cyclobutanol ^a	15 h (room temp.)	990 (66)	cyclobutanone (44) propanoic acid (45)
Cyclobutanol ^e	4 h (room temp.)	81 (20)	cyclobutanone (100) ¹¹

Reaction conditions: ^a 0.3 mmol of substrate, 0.3×10^{-3} mmol catalyst and 0.6 mmol of NaBrO_3 as co-oxidant were mixed in 5 ml of water. ^b Catalyst $\text{trans-}[\text{RuO}_2(\text{HIO}_6)_2]^{6-}$ and NaIO_4 as co-oxidant.¹¹ ^c Catalyst $\text{cis-}[\text{Ru}(\text{bpy})_2\text{Cl}_2]$ and NaBrO_3 as co-oxidant. ^d Catalyst $[\text{RuO}_2(\text{bpy})\text{-}[\text{IO}_3(\text{OH})_3]] \cdot 1.5\text{H}_2\text{O}$ and NaIO_4 as co-oxidant.⁹ ^e Catalyst $[\text{Ru}(\text{dcbpy})_2(\text{H}_2\text{O})_2]^{2+}$ ¹² (electrochemically); dcbpy = 6,6'-dichlorobipyridine.

Table 2 Epoxidation of alkenes, catalysed by $[\text{Ru}_2(\text{napy})_2(\text{OH})\text{Cl}(\text{H}_2\text{O})_4][\text{ClO}_4]_4$ **1** and some reference compounds from the literature

Substrate	Reaction conditions	Co-oxidant	Turnovers	Product(s) (% yield)
<i>trans</i> -Stilbene ^a	24 h (room temp.)	NaIO_4	760	benzaldehyde (62) benzoic acid (10) <i>trans</i> -stilbene oxide (4)
<i>trans</i> -Stilbene ^{b,c}	72 h (40 °C)	$\text{O}_2/\text{isobutyraldehyde}$	980	benzaldehyde (15) <i>trans</i> -stilbene oxide (83)
<i>cis</i> -Stilbene ^{b,c}	72 h (40 °C)	$\text{O}_2/\text{isobutyraldehyde}$	660	benzaldehyde (13) <i>cis</i> -stilbene oxide (53)
<i>trans</i> -Stilbene ^d	15 h (room temp.)	NaIO_4	249	<i>trans</i> -stilbene oxide (99) ¹⁰
<i>cis</i> -Stilbene ^d	15 h (room temp.)	NaIO_4	189	<i>cis</i> -stilbene oxide (75) ¹⁰
Cyclooctene ^{b,c}	48 h (40 °C)	$\text{O}_2/\text{isobutyraldehyde}$	840	1,2-epoxycyclooctane (84)

Reaction conditions: ^a 0.15 mmol of *trans*-stilbene, 0.15×10^{-3} mmol catalyst and 0.3 mmol of NaBrO_3 or NaIO_4 were dissolved in a mixture of 2 ml of water and 2 ml of 1,2-dichloroethane. ^b 0.15 mmol of *trans*-stilbene, 54 μl isobutyraldehyde and 0.15×10^{-3} mmol catalyst were dissolved in a vigorously stirred aerated mixture of 2 ml of water and 2 ml of 1,2-dichloroethane. ^c Blank experiments without the ruthenium catalyst yielded no more than 10% conversion under the same reaction conditions. ^d Catalyst $[\text{RuO}_2(\text{bpy})\text{-}[\text{IO}_3(\text{OH})_3]] \cdot 1.5\text{H}_2\text{O}$.⁹

$\text{CF}_3\text{SO}_3\text{H}$ shows a two-electron oxidation of $[\text{Ru}^{\text{III}}\text{Ru}^{\text{III}}(\text{napy})_2(\text{H}_2\text{O})_4\text{Cl}(\text{OH})]^{4+}$ to the $\text{Ru}^{\text{IV}}\text{Ru}^{\text{IV}}$ analogue, with an isobestic point at 390 nm. A plot of the absorbance of **1** at 384 nm (also observed at other wavelengths) vs. the $\text{Ce} : [\text{Ru}_2]$ mol ratio is linear ($R^2 = 0.987$), with an apparent end point at $\text{Ce} : [\text{Ru}_2]$ mol ratio of 2 ± 0.2 . Addition of an excess of NaBrO_3 to the $\text{Ru}^{\text{III}}\text{Ru}^{\text{III}}$ species gives a similar absorbance spectrum (as of the high-valent complex obtained with cerium), albeit at a slower rate. The spectroscopic and oxidation results suggest that the $\text{Ru}^{\text{III}}\text{Ru}^{\text{III}}$ complex is oxidized to $\text{Ru}^{\text{IV}}\text{Ru}^{\text{IV}}$ by NaBrO_3 , after which this high-valent species can react either by two subsequent one-electron steps or by a two-electron step back to $\text{Ru}^{\text{III}}\text{Ru}^{\text{III}}$, thereby oxidizing alcoholic substrates. However, more detailed (electrochemical) experiments are needed to support this mechanism. In addition, we are currently investigating the catalytic potential of ruthenium complexes with 1,8-naphthyridine derivatives.

This work has been sponsored by The Netherlands Ministry of Agriculture, through its special Programme on Carbohydrate Oxidation, The Netherlands Organisation for Scientific Research (T. X. N.) and in part (A. L. S) by the Netherlands Foundation of Chemical Research (SON) with financial aid from the Netherlands Organisation for Scientific Research

(NWO). Professor Dr W. P. Griffith is acknowledged for useful suggestions. A generous loan of RuCl_3 by Johnson and Matthey is gratefully acknowledged.

Received, 18th July 1995; Com. 5/04713G

Footnotes

† On leave from AT & T Bell Laboratories, USA.

‡ Selected data for **1**: ^1H NMR (300 MHz, 0.1 mol dm^{-3} $\text{CF}_3\text{SO}_3\text{D}$ vs. 4,4-dimethyl-4-silapentanesulfonic acid), δ 7.83 (dd, 4H, $\text{H}^{3,6}$), 8.57 (dd, 4H, $\text{H}^{4,5}$), 8.90 (dd, 4H, $\text{H}^{2,7}$). For **2**: ^1H NMR [300 MHz, $(\text{CD}_3)_2\text{SO}$ vs. SiMe_4], δ 7.76 (m, 4H, $\text{H}^{3,6}$), 8.48 (dd, 4H, $\text{H}^{4,5}$), 9.53 (dd, 4H, $\text{H}^{2,7}$). Elemental analysis. Calc. for $\text{C}_{16}\text{H}_{12}\text{Cl}_4\text{N}_4\text{Ru}_2 \cdot 2\text{H}_2\text{O}$: C, 30.02; H, 2.52; Cl, 22.15; N, 8.75. Found: C, 29.94; H, 2.53; Cl, 22.91; N, 7.95%.

§ Crystal data for compound **1**, $\text{C}_{16}\text{H}_{12}\text{Cl}_4\text{N}_4\text{O}_{21}\text{Ru}_2 \cdot 3\text{H}_2\text{O}$, $M_r = 1038.81$, green, needle-shaped crystal ($0.05 \times 0.10 \times 0.60$ mm), triclinic, space group $P\bar{1}$ (no. 2) with $a = 8.7920(9)$, $b = 10.2100(12)$, $c = 11.0160(12)$ Å, $\alpha = 111.089(11)$, $\beta = 109.759(9)$, $\gamma = 100.608(9)^\circ$, $U = 813.98(17)$ Å³, $Z = 1$, $D_c = 2.119$ g cm^{-3} , $F(000) = 516$, $\mu(\text{Mo-K}\alpha) = 14.4$ cm^{-1} , 5356 reflections measured, 3737 independent, $R_{\text{int}} = 0.100$, ($2.21^\circ < \theta < 27.49^\circ$), ω scan, $T = 150$ K, Mo-K α radiation, graphite monochromator, $\lambda = 0.71073$ Å) on an Enraf-Nonius CAD4 Turbo diffractometer on

rotating anode. Data were corrected for Lp effects and for a linear decay of 4% of the reference reflections, but not for absorption. The structure was solved by automated Patterson methods (SHELXS86). Refinement on F^2 was carried out by full-matrix least-squares techniques (SHELXL-93); no observance criterion was applied during refinement. Refinement converged at a final $wR2$ value of 0.191, $R1 = 0.082$ [for 2298 reflections with $F_0 > 4\sigma(F_0)$], $S = 1.01$, for 266 parameters. A final difference Fourier map showed no residual density outside -1.32 and $1.48 \text{ e } \text{Å}^{-3}$ (near Ru). Atomic coordinates, bond lengths and angles and thermal parameters have been deposited at the Cambridge Crystallographic Data Centre. See Information for Authors, Issue No. 1.

¶ All catalytic reactions were conducted under oxygen. Specific blank experiments did not show any activity towards the substrate unless stated otherwise. All reaction substrates, except adipic acid, were analysed by gas chromatography and satisfactory compared to commercial samples. Adipic acid was analysed by ^1H NMR.

|| Spectrophotometric redox titrations were carried out by adding aliquots of a $1.8 \times 10^{-3} \text{ mol dm}^{-3}$ solution of Ce^{IV} in $0.1 \text{ mol dm}^{-3} \text{ CF}_3\text{SO}_3\text{H}$ to aliquots (0.5 ml) of a $3.1 \times 10^{-4} \text{ mol dm}^{-3}$ solution of **1** in $0.1 \text{ mol dm}^{-3} \text{ CF}_3\text{SO}_3\text{H}$. The volume was adjusted to 3 ml by addition of $0.1 \text{ mol dm}^{-3} \text{ CF}_3\text{SO}_3\text{H}$, and the changes were monitored in the range 200–600 nm. The Ce: [Ru₂] mol ratio was varied from 0 to 6. Ce^{IV} solutions in $0.1 \text{ mol dm}^{-3} \text{ CF}_3\text{SO}_3\text{H}$ were prepared from $[\text{NH}_4]_2[\text{Ce}(\text{NO}_3)_6]$.

References

- 1 T. V. Lee in *Comprehensive Organic Synthesis*, ed. B. M. Trost, I. Fleming and S. V. Ley, Pergamon, Oxford, 1991, ch. 7, 291; *Fieser and Fieser's Reagents for Organic Synthesis*, ed. M. Fieser and J. C. Smith, Wiley, 1988, vol. 13, index 'oxidation'.
- 2 G. Balavoine, C. Eskenazi, F. Meunier and H. Riviere, *Tetrahedron Lett.*, 1984, **25**, 3187.
- 3 See for example: W. P. Griffith, *Chem. Soc. Rev.*, 1991, **21**, 179; L. Roecker and T. J. Meyer, *J. Am. Chem. Soc.*, 1987, **109**, 746; M. S. Thompson and T. J. Meyer, *J. Am. Chem. Soc.*, 1982, **104**, 4106; L. Roecker and T. J. Meyer, *J. Am. Chem. Soc.*, 1986, **108**, 4067; S. J. Raven and T. J. Meyer, *Inorg. Chem.*, 1988, **27**, 4478. C. M. Che, W. T. Tang, K. Y. Lee, K. Y. Wong and T. C. Lau, *J. Chem. Soc., Dalton Trans.*, 1992, 1551.
- 4 J. A. Gilbert, D. S. Eggleston, W. R. Murphy, D. A. Gesolowitz, S. W. Gersten, D. W. Hodgson and T. J. Meyer, *J. Am. Chem. Soc.*, 1985, **107**, 3855; K. Honda and A. J. Frank, *J. Chem. Soc., Chem. Commun.*, 1984, 1635; J. P. Collin and J. P. Sauvage, *Inorg. Chem.*, 1986, **25**, 135; J. K. Hurst, J. Zhou and Y. Lei, *Inorg. Chem.*, 1992, **31**, 1010; C. D. Ellis, J. A. Gilbert, W. R. Murphy and T. J. Meyer, *J. Am. Chem. Soc.*, 1983, **105**, 4843; S. J. Raven and T. J. Meyer, *Inorg. Chem.*, 1988, **27**, 4478.
- 5 A. Gerli and J. Reedijk, *J. Mol. Catal.*, 1994, **89**, 101; W. Kutner, J. A. Gilbert, A. Tomaszewsky, T. J. Meyer and R. W. Murray, *J. Electroanal. Chem.*, 1986, **205**, 185; L. Roecker, W. Kutner, J. A. Gilbert, M. Simmons, R. W. Murray and T. J. Meyer, *Inorg. Chem.*, 1985, **24**, 3784.
- 6 R. W. Brookes and R. L. Martin, *Inorg. Chem.*, 1975, **14**, 528; L. Sacconi, C. Mealli and D. Gatteschi, *Inorg. Chem.*, 1974, **13**, 1985; D. Gatteschi, C. Mealli and L. Sacconi, *Inorg. Chem.*, 1976, **15**, 2774; K. Emerson, A. Emad, R. W. Brookes and R. L. Martin, *Inorg. Chem.*, 1973, **12**, 978; C. Mealli and F. Zanobini, *J. Chem. Soc., Chem. Commun.*, 1982, 97; T. Tsuda, S. Ohba, M. Takahashi and M. Ito, *Acta Crystallogr., Sect. C.*, 1989, **45**, 887.
- 7 J. Roček and A. E. Radkowski, *J. Am. Chem. Soc.*, 1973, **95**, 7123.
- 8 B. P. Sullivan, D. J. Salmon and T. J. Meyer, *Inorg. Chem.*, 1978, **17**, 3334.
- 9 A. J. Bailey, W. P. Griffith, A. J. P. White and D. J. Williams, *J. Chem. Soc., Chem. Commun.*, 1994, 1833.
- 10 T. Mukaiyama, T. Yamada, T. Nagata and K. Imagawa, *Chem. Lett.*, 1993, 327; T. Yamada, K. Imagawa, T. Nagata and T. Mukaiyama, *Chem. Lett.*, 1992, 2231; G. A. Barf and R. A. Sheldon, *J. Mol. Catal. A*, 1995, **102**, 23.
- 11 A. M. El-Hendawy, W. P. Griffith, B. Piggott and D. J. Williams, *J. Chem. Soc., Dalton Trans.*, 1988, 1983.
- 12 K. Y. Wong, W. O. Lee and C. M. Che, *J. Electroanal. Chem.*, 1991, **319**, 207.