


# 1,12-Hydrogen atom abstraction reactions of $\alpha$ -keto esters

George A. Kraus,\* Weijiang Zhang and Yusheng Wu


Department of Chemistry, Iowa State University, Ames, IA 50011, USA

## Eleven-membered ring lactones are made by 1,12-hydrogen atom abstraction.


Photochemical hydrogen atom abstraction reactions proceeding via six-atom transition states are well known in organic chemistry. Such reactions include the Norrish Type II reaction and the photoenolization reaction.<sup>1</sup> Hydrogen atom abstraction reactions proceeding via many-atom transition states have also been extensively studied. Examples include the remote functionalization reactions developed by Breslow and others.<sup>2</sup> In contrast, hydrogen atom abstraction reactions which proceed by medium-ring transition states are comparatively rare. Wagner recently reported a number of examples.<sup>3</sup>



Scheme 1



Scheme 2




Scheme 3 Reagents: i,  $\text{BrCH}_2\text{COBr}$ ,  $\text{Pr}_2\text{NEt}$ ,  $\text{PhCOCO}_2\text{H}$ ,  $\text{K}_2\text{CO}_3$ , DMF; ii, LiOH

In the course of examining photochemical strategies for the initiation of radical cyclizations, we recently discovered several novel 1,9-hydrogen atom abstraction reactions.<sup>4</sup> An example is shown in Scheme 1. Our hypothesis is that the more stable *syn* conformation of the ester is the key control element that favours the 1,9-hydrogen atom abstraction, leading to the eight-membered ring lactone.

The hydroxy lactones produced in this reaction are potential intermediates for the preparation of natural products containing the oxocene ring system.<sup>5</sup> In order for our photochemical methodology to be more directly applicable, we decided to prepare aldehyde 1b. When diol 2 was subjected to anhydrous periodate,<sup>6</sup> only hex-3-en-1-ol was isolated (Scheme 2).

We next devised a photochemical method for the production of 1b. Binkley has shown that the photolysis of  $\alpha$ -keto esters affords labile aldehydes.<sup>7</sup> The plan was to produce 1b under conditions where it could be irradiated without isolation. Alcohol 3 was acylated with bromoacetyl bromide and diisopropylethylamine in THF at 0 °C and was converted into diester 4 by displacement of the bromide with 1.5 equiv. of the potassium salt of phenylglyoxylic acid in DMF at 25 °C (Scheme 3). Irradiation in benzene using a Rayonet apparatus (350 nm bulbs) for 18 h afforded products 5 and 6 in 24 and 37% yield, respectively. The structure of 5 was assigned based on double irradiation experiments and infrared absorptions at 1735 and 1752  $\text{cm}^{-1}$ .<sup>†</sup> Hydrolysis of 5 with LiOH at 25 °C afforded acid 7.  $\beta$ -Lactone 6 was characterized by its infrared spectrum showing absorptions at 1846 and 1735  $\text{cm}^{-1}$  and its NMR spectra. Compound 5 could be derived from a 1,12-hydrogen atom abstraction followed by cyclization. Interestingly, only one ring size was produced, even though an allylic radical was involved. Although a few 1,12-hydrogen atom abstraction reactions have been observed, these reactions involve  $\omega$ -aminophthalimides, where charge transfer reactions can be operative.<sup>8</sup>

Additionally, 8 was prepared by the same pathway and was irradiated in the Rayonet reactor (Scheme 4). Lactone 9 was obtained in 28% yield.<sup>‡</sup>



Scheme 4

## Footnotes

† Selected data for **5**: 400 MHz  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  0.75–0.80 (m, 3 H), 2.00–2.20 (m, 1 H), 2.60–2.75 (m, 1 H), 3.41–3.55 (m, 1 H), 3.69 (s, 1 H), 4.10–4.18 (m, 2 H), 4.25 (d,  $J$  15 Hz, 1 H), 5.07 (d,  $J$  15 Hz, 1 H), 5.40–5.50 (m, 1 H), 5.75 (t,  $J$  9 Hz, 1 H), 7.25–7.40 (m, 3 H), 7.60–7.72 (m, 2 H); IR (KBr): 3522, 3016, 2926, 1752, 1735, 1286, 752  $\text{cm}^{-1}$ . HRMS for  $\text{C}_{16}\text{H}_{18}\text{O}_5$ : found, 290.1152. Calc., 290.1154.

‡ Selected data for **9**: 300 MHz  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  3.72 (br s, 1 H), 5.07 (s, 1 H), 5.12 (s, 2 H), 5.42 (dd,  $J$  5, 10 Hz, 2 H), 6.98 (d,  $J$  6 Hz, 2 H), 7.26–7.60 (m, 12 H); IR (KBr) 3446, 3063, 1734, 1719, 1246, 754  $\text{cm}^{-1}$ . HRMS for  $\text{C}_{24}\text{H}_{20}\text{O}_6$ : found, 404.1251. Calc., 290.1259.

## References

- 1 P. J. Wagner and B.-S. Park, *Org. Photochem.*, 1991, **11**, 227.
- 2 R. Breslow, *Comprehensive Organic Synthesis*, ed. B. M. Trost, Pergamon Press, NY, 1991, pp. 39–52.
- 3 B. Zhou and P. J. Wagner, *J. Am. Chem. Soc.*, 1989, **111**, 6796.
- 4 G. A. Kraus and Y. Wu, *J. Am. Chem. Soc.*, 1992, **114**, 8705.
- 5 D. Faulkner, *Nat. Prod. Rep.*, 1986, **3**, 1.
- 6 T. R. Kelly, T. E. Schmidt and J. G. Haggerty, *Synthesis*, 1972, 544.
- 7 R. W. Binkley, *Synth. Commun.*, 1976, **6**, 281.
- 8 T. Hasegawa, K. Miyata, T. Ogawa, N. Yoshihara and M. Yoshioka, *J. Chem. Soc., Chem. Commun.*, 1985, 363.

Received, 2nd July 1996; Com. 6/04666E