

Synthesis and structure of the first 2*H*-1,3,2-diazaphosphole complex

Hendrik Wilkens, Jörg Jeske, Peter G. Jones and Rainer Streubel*

Institut für Anorganische und Analytische Chemie der Technischen Universität Braunschweig, Postfach 3329, D-38023 Braunschweig, Germany


Thermal decomposition of the 2*H*-azaphosphirene tungsten complex **1** in benzonitrile and in the presence of dimethylcyanamide yields the 2*H*-1,3,2-diazaphosphole tungsten complex **4**, containing a novel five-membered heterocyclic ligand; the overall reaction may be described as a 1,3-dipolar cycloaddition of an *in situ* generated nitrilium phosphane ylide complex to benzonitrile.

Nitrilium ylides, like nitrile oxides,¹ sulfides² and nitrilimines,^{1,3} are versatile building blocks in heterocyclic chemistry. Very recently, we obtained the first evidence of a nitrilium phosphane ylide complex as a reactive intermediate in the thermolysis of 2*H*-azaphosphirene tungsten complexes in the presence of dimethylacetylenedicarboxylate (DMAD) in toluene.⁴ The formation of the 2*H*-1,2-azaphosphole tungsten complexes, which were obtained as by-products, was explained as a 1,3-dipolar cycloaddition of a nitrilium phosphane ylide complex to the trapping reagent.⁴

We now report the first synthesis of a 2*H*-1,3,2-diazaphosphole tungsten complex **4**, which has been obtained in good yield by thermal decomposition of the 2*H*-azaphosphirene tungsten complex **1** in benzonitrile and in the presence of 2 equiv. of dimethylcyanamide. Under these reaction conditions the 1,3-dipolar cycloaddition is stereoselective and the exclusive formation of this regioisomer was observed.

This reaction is interpreted as a further example of a 1,3-dipolar cycloaddition of the transiently formed nitrilium phosphane ylide tungsten complex **2b** to benzonitrile (Scheme 1, pathway (c)). Pathways (a) and (b) seem less plausible, especially, if taken into account that under these reaction conditions the formation of the diphenyl-substituted 2*H*-1,3,2-diazaphosphole tungsten complex **3** was not observed [pathway (a)].

The composition of 2*H*-1,3,2-diazaphosphole tungsten complex **4** is confirmed by elemental analysis and mass spectrometry;† the structural formulation is based on the characteristic



Scheme 1 Reagents and conditions: 1 mmol **1** was treated with 2 mmol dimethylcyanamide in benzonitrile at 75 °C for 1.5 h. Work-up by column chromatography at low temperature afforded **4** in good yield (67%, mp 90 °C).




Fig. 1 Molecular structure of complex **4** in the crystal. Radii are arbitrary. Selected bond lengths (Å) and angles (°): P–W 2.479(11), P–N(1) 1.705(3), P–N(2) 1.705(3), N(1)–C(7) 1.291(5), N(2)–C(8) 1.284(5), C(7)–C(8) 1.488(5); N(1)–P–N(2) 96.3(2), P–N(1)–C(7) 108.2(3), N(1)–C(7)–C(8) 114.0(3), N(2)–C(8)–C(7) 113.4(4), P–N(2)–C(8) 107.8(3).

NMR spectral data† in solution and was confirmed for the solid state by a X-ray crystal structure analysis.‡ In comparison to related 2*H*-1,2-azaphosphole tungsten complexes,⁴ the 2*H*-1,3,2-diazaphosphole tungsten complex **4** displays a low-field shifted resonance signal of the phosphorus nucleus at δ 149.8 (*cf.* δ 102–105⁴) and a markedly increased phosphorus tungsten coupling constant of 257.1 Hz (*cf.* $^1J_{WP}$ 236–238 Hz⁴). The phosphorus–carbon coupling constants of the quaternary ring carbon atoms are 2–10 Hz, which are characteristically small for both five-membered heterocycles.

Remarkable features of the molecular structure of **4** and 2*H*-1,2-azaphosphole complexes⁴ are the planar five-membered ring system and the localised double bonds; the latter being in contrast to the situation of aromatic heterocycles like 2*H*-1,2,3-diazaphospholes^{5,6} and 2*H*-1,2,3-triazoles.⁷ The coordination sphere of phosphorus is distorted tetrahedral and the phosphorus–tungsten bond length is 2.479(11) Å.

We are currently investigating the synthetic potential of nitrilium phosphane ylide complexes in 1,3-dipolar cycloaddition reactions.

This work was supported by the Fonds der Chemischen Industrie and by the Deutsche Forschungsgemeinschaft.

Footnotes and References

* E-mail: streubel@mvs.anchem.nat.tu-bs.de

† Satisfactory elemental analysis was obtained for complex **4**. NMR data were recorded at room temp. in CDCl₃ solution at 50.3 MHz (¹³C) and 81.0

(³¹P), using TMS and 85% H₃PO₄ as standard references; *J*/Hz. Selected spectroscopic data for **4**: ¹³C NMR, δ 162.4 (d, Σ*J*_{PC} 9.4, P–N=C), 165.0 (d, Σ*J*_{PC} 2.5, P–N=C); ³¹P NMR, δ 149.8 (5, ¹*J*_{WP} 257.1); MS (EI): M⁺ at *m/z* 687.

‡ Crystal data for **4**: C₂₂H₃₀N₃O₅PSi₂W, *M* = 687.49; triclinic, space group *P*1̄: *a* = 9.284(3), *b* = 10.359(3), *c* = 16.321(4) Å, α = 75.07(2), β = 88.21(2), γ = 67.67(2)°; *U* = 1398.9(6) Å³; *Z* = 2, *D*_c = 1.632 Mg m⁻³, *T* = 143 K, a crystal (0.7 × 0.7 × 0.6 mm) was mounted in inert oil. A total of 4946 absorption corrected intensities were measured to 2θ = 50° using Mo-Kα radiation on a STOE Stadi-4 diffractometer; 4915 were unique (*R*_{int} = 0.0546) and used for all calculations (program SHELXL-93). The structure was solved by direct methods and refined anisotropically on *F*². The final *wR*₂ was 0.0686, with conventional *R*(*F*) 0.0262 for 316 parameters and 151 restraints; max. Δρ 2.48 e Å⁻³. CCDC 182/666.

- 1 P. Caramella and P. Grünanger, in *1,3 Dipolar Cycloaddition Chemistry*, ed. A. Padwa, Wiley, New York, 1984, ch. 3.
- 2 R. M. Paton, *Chem. Soc. Rev.*, 1989, **18**, 33.
- 3 G. Bertrand and C. Wentrup, *Angew. Chem., Int. Ed. Engl.*, 1994, **33**, 527.
- 4 R. Streubel, H. Wilkens, A. Ostrowski, C. Neumann, F. Ruthe and P. G. Jones, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 1492.
- 5 J. H. Weinmaier, H. Tautz, A. Schmidpeter and S. Pohl, *J. Organomet. Chem.*, 1980, **185**, 53.
- 6 K. Karaghiosoff, W. S. Sheldrick and A. Schmidpeter, *Chem. Ber.*, 1986, **119**, 3213.
- 7 H. Dehne, in Houben-Weyl, *Meth. Org. Chem.*, Bd. E 8d, 1994, p. 305.

Received in Basel, Switzerland, 18th June 1997; 7/04285J