

Chemoselective isomerization of amide-substituted oxetanes with Lewis acid to give oxazine derivatives or bicyclic amide acetals

Tomonari Nishimura,^a Shigeyoshi Kanoh,^{*a†} Hitoshi Senda,^a Toshiyuki Tanaka,^b Kohji Ando,^a Hiroshi Ogawa^a and Masatoshi Motoi^{*a}

^a Department of Industrial Chemistry, Faculty of Engineering, Kanazawa University, Kodatsuno, Kanazawa 920-8667, Japan

^b Department of Pharmacognosy, Gifu Pharmaceutical University, Mitahora-higashi, Gifu 502-8585, Japan

The Lewis-acid catalyzed isomerization of secondary and tertiary amide-substituted oxetanes takes place chemoselectively, giving 5-hydroxymethyl-5,6-dihydro-4*H*-1,3-oxazines and reactive amide acetals consisting of a bicyclo[2.2.2]octane skeleton, respectively.


Oxetane and its various derivatives are well known to be polymerizable cationically *via* conventional ring-opening processes.^{1,2} As an exception, certain oxetanes carrying a carbonyl-containing substituent, however, can undergo Lewis-acid catalyzed intramolecular nucleophilic attack of the unsaturated oxygen atom, for example, ester-substituted oxetanes **1** give the bridged orthoesters **2**,³ and cyclic imide-substituted oxetanes **3** give bicyclic acetals with a lactam ring **4**⁴ (Scheme 1). Here we report a novel chemoselective isomerization of oxetanes having an amide group at their 3-positions. The isomerization of the

amide-substituted oxetanes **5** with Lewis acid gives two quite different types of products, depending on the structure of amide group. One type is 5-hydroxymethyl-5,6-dihydro-4*H*-1,3-oxazines **6** produced from the secondary amides of **5** by a new mode of rearrangement involving the oxetanyl group. The other type is bicyclic amide acetals, *i.e.* 2,6-dioxa-7-azabicyclo[2.2.2]octanes **7**, which are produced from the tertiary amides of **5** by a similar mode to those in the already reported cases of **1** and **3**.

Starting oxetane amides **5a–e** were prepared *via* acylation of oxetane amine **8** derived from **3** (phthalimide), and the resulting secondary amides were *N*-alkylated under phase-transfer catalyzed conditions⁵ to give the tertiary amides **5f–k**.[‡]

The isomerization of **5** was successfully carried out in anhydrous chlorobenzene or CH₂Cl₂ using Lewis acids such as BF₃·OEt₂ and AlMe₃ (Table 1).[§] Like **1** and **3** cycloacetalization occurred in the reactions of **5f–k**. The products (**7f–k**) obtained are a new series of bicyclic amide acetals.[¶] Analogous amide acetals of the bicyclo[*n*.3.0] type (*n* = 3 or 4) have already been synthesized by intermolecular cycloaddition or transacetalization, and they have been reported to possess high susceptibility to nucleophiles.⁶ Similarly compounds **7** were so sensitive to moisture that hydrolytic fission proceeded rapidly under normal conditions.^{||} When the above isomerization was run for 24 h at 100 °C and above, most of the **7** produced was consumed. The final products were oligomers having a polyether backbone.

In sharp contrast, the isomerization of **5a** unexpectedly afforded 5-hydroxymethyl-5-methyl-2-phenyl-5,6-dihydro-4*H*-1,3-oxazine **6a** instead of a bicyclic amide acetal, and no polymerization took place.[¶] The molecular structure of **6a** was established by X-ray analysis (Fig. 1).^{**} The isomerization of the other secondary amides (**5b–e**) always gave oxazine derivatives (**6b–e**) in good yields (Table 1).[¶] Compounds **6** were stable during distillation or recrystallization, but chromatographic manipulation over alumina often caused hydrolysis to propanediols with amide groups.



Scheme 1

Table 1 Lewis-acid catalyzed isomerization of oxetane amides **5a**

Amide	(R ¹ , R ²)	Lewis acid	Reaction conditions			Product	
			(mol%)	T/°C	t/h	Yield (%) ^b	
5a ^c	(Ph, H)	BF ₃ ·OEt ₂	(25)	35	96	6a	89
5b	(H, H)	AlMe ₃	(50)	120	24	6b	50 ^d
5c	(Me, H)	AlMe ₃	(5)	120	24	6c	92
5d	(Pr ⁿ , H)	AlMe ₃	(5)	120	24	6d	89
5e	(PhCH ₂ , H)	AlMe ₃	(5)	120	24	6e	71
5f	(Ph, Et)	BF ₃ ·OEt ₂	(5)	120	1	7f	59
5g	(Ph, PhCH ₂)	BF ₃ ·OEt ₂	(5)	130	1	7g	61
5h	(H, PhCH ₂)	BF ₃ ·OEt ₂	(35)	70	96	7h	63
5i	(Me, Et)	BF ₃ ·OEt ₂	(5)	130	1	7i	55
5j	(Me, PhCH ₂)	BF ₃ ·OEt ₂	(5)	130	1	7j	65
5k	(Pr ⁱ , Et)	BF ₃ ·OEt ₂	(5)	130	1.5	7k	48

^a In PhCl. ^b Isolated yield. ^c In CH₂Cl₂. ^d The reaction was quenched with Et₃N followed by MeOH containing a small amount of dilute NaOH.





Fig. 1 ORTEP view of **6a** (30% probability thermal ellipsoids)

Although the secondary and tertiary amides of **5** are isomerized giving structurally different products, these reactions can be explained by the following mechanisms passing through a common intermediate **B** (Scheme 2). Both isomerizations begin with the coordination of Lewis acid E to oxetanyl oxygen atom of **5**, and then oxonium **A** undergoes intramolecular nucleophilic attack of the carbonyl oxygen atom by neighboring group participation. The fate of the resulting **B** determines the consequent chemoselection. For tertiary amides, subsequent ring closure of **B** gives an equilibrium mixture of **C** and **C'**. On the other hand, **B** is interconvertible to **B'** when R^2 is a hydrogen atom. The iminium and ammonium salts, **B'** and **C'**, are probably thermodynamically more stable than **B** and **C**, and the catalytic turnover of Lewis acid would decrease as the isomerization progresses. This was strongly supported by the observation that the isomerization rates of **5** steeply slowed down at increased yields of basic **6** and **7**. In order to force these reactions to completion, it was necessary to employ either high temperature or a very large amount of Lewis acid, or both (Table 1). Under such conditions, however, the yields of **7** began to decrease after reaching maxima at relatively early stages, because of inevitable polymerization as a competing reaction.

In the chemoselective isomerization of **5**, moderate to fairly good yields of **6** and **7** were obtained using simple operations. Like other bicyclic amide acetals,⁶ compounds **7** are also interesting in a variety of chemical transformations not only as reactive amino ethers but also as ring-opening polymerizable monomers.^{6,7} Numerous methods of preparing dihydro-1,3-oxazines appear in the literature.⁸ The advantages (as well as differences) in the present reaction may be summarized as follows: (a) the preparation of dihydro-1,3-oxazines having a hydroxymethyl group, which is a useful functional group for further modification. (b) A general method for obtaining not




Scheme 2

only the phenyl- or alkyl-substituted oxazines at the 2-positions but also the unsubstituted oxazine.

Footnotes and References

† E-mail: kanoh@kenroku.ipc.kanazawa-u.ac.jp

‡



Reagents and conditions: i, 3-methyl-3-(phthalimidomethyl)oxetane⁴ (1.0 equiv.), hydrazine hydrate (5 equiv.), in EtOH, 60 °C, 1.5 h, and then a catalytic amount of Raney Ni (W-2), vigorous reflux, 81% (GLPC purity 96%), bp 90–95 °C/100 mmHg; ii, acylating reagents: (R¹CO)₂O, HCO₂COMe, or R¹CO₂H (DCC coupling), 76–92%; iii, **5** (1.0 equiv.), R²Br (1.1 equiv.), K₂CO₃ (1.0 equiv.), NaOH (2.5 equiv.), Buⁿ₄NHSO₄ (0.10 equiv.), in benzene, 40 °C to reflux, 3–15 h, 55–78%.

§ *Typical experimental procedure:* To an anhydrous PhCl (1.0 ml) solution of **5c** (150 mg, 1.1 mmol) in a tube was added 1.0 mol l⁻¹ AlMe₃ in hexane (0.05 mmol) under nitrogen. The resulting solution was allowed to react at 120 °C for 24 h. After the reaction was quenched by addition of Et₃N (0.1 ml) followed by MeOH (3.0 ml), the solvents were replaced by CH₂Cl₂ (10 ml). Distillation of the CH₂Cl₂-soluble part afforded **6c** in 92% yield, bp 100–110 °C/0.1 mmHg. Similarly, the isomerization of **5f** was carried out in a small distillation flask at 120 °C for 1 h. After the addition of anhydrous Et₃N (0.1 ml) followed by CaH₂ (ca. 50 mg), distillation of the resulting mixture afforded **7f** (0.12 g, 0.52 mmol) in 59% yield, bp 110–120 °C/0.1 mmHg. The product isolated was stable under nitrogen.

¶ All compounds were characterized by ¹H NMR and IR spectroscopy, although the IR spectra of **7** were always observed as a mixture of **7** and its hydrolysis product. *Selected data for 6a:* mp 135–136 °C (CH₂Cl₂–hexane); ν_{max}(KBr)/cm⁻¹ 3400 (OH), 1650 (C=N); the atomic numbering system refers to that used in Fig. 1; δ_H(CDCl₃, J/Hz) 7.89 (dd-like, J 6.9, 1.5, 2 H, ArH_o), 7.44–7.33 (m, 3 H, ArH_{m,p}), 4.23 [dd, J 10.7, 2.4, 1 H, H(6b)], 3.92 [d-like, J 10.7, 1 H, H(6a)], 3.57, 3.47 [each d, J 10.7, 2 H, H(14a) and H(14b)], 3.48 [dd, J 16.9, 2.2, 1 H, H(4b)], 3.27 [d-like, J 16.6, 1 H, H(4a)], 2.02 (br s, 1 H, OH), 1.02 [s, 3 H, C(13) H₃]; δ_C (100 MHz, CDCl₃) 155.0 [C(2)], 133.3 [C(7)], 130.5 [C(10)], 128.0 [C(9), C(11)], 127.0 [C(8), C(12)], 70.1 [C(6)], 65.8 [C(14)], 51.1 [C(4)], 32.7 [C(5)], 19.0 [C(13)]. HRMS: *m/z* 205.1115, calc. for C₁₂H₁₅NO₂: 205.1104. Analysis for C₁₂H₁₅NO₂, found (calc.): C, 69.96 (70.22); H, 7.33 (7.37); N, 6.76 (6.82%). For **7f**: δ_H(anhydrous CDCl₃ under nitrogen, J/Hz) 7.60 (dd, J 6.8, 3.2, 2 H, ArH_o), 7.38–7.29 (m, 3 H, ArH_{m,p}), 3.98 (s, 4 H, OCH₂), 2.99 (s, 2 H, NCH₂), 2.36 (q, J 7.2, 2 H, CH₂CH₃), 0.94 (t, J 7.3, 3 H, CH₂CH₃), 0.88 (s, 3 H, CCH₃). HRMS: *m/z* 233.1439, calc. for C₁₄H₁₉NO₂: 233.1417.

|| The hydrolysis product is 2-ethylaminomethyl-2-hydroxymethylpropyl benzoate. The initially formed esters except for the benzoates and isobutyrate were spontaneously converted to the corresponding amide-substituted propanediols by acyl exchange.

** **6a:** Mp 135–136 °C (CH₂Cl₂–hexane). *Crystal data for 6a:* space group P2₁/c, Z = 4, a = 11.044(1), b = 9.3865(8), c = 11.817(1) Å, β = 116.336(8)°, D_c = 1.242 g cm⁻³. 2816 measured, 2685 independent reflections, of which 1813 were considered as observed [I > 3.00 σ(I)]. R = 0.039, R_w = 0.035. CCDC 182/639.

- M. P. Dreyfuss and P. Dreyfuss, in *Encyclopedia of Polymer Science and Engineering*, Wiley, New York, 1987, vol. 10, pp. 653–670.
- H. Ogawa, T. Hosomi, T. Kosaka, S. Kanoh, A. Ueyama and M. Motoi, *Bull. Chem. Soc. Jpn.*, 1997, **70**, 175; M. Motoi, S. Sekizawa, K. Asakura and S. Kanoh, *Polym. J.*, 1993, **25**, 1283 and references cited therein.
- E. J. Corey and N. Raju, *Tetrahedron Lett.*, 1983, **24**, 5571; P. Ducray, H. Lamotte and B. Rousseau, *Synthesis*, 1997, 404.
- S. Kanoh, T. Hashiba, K. Ando, H. Ogawa and M. Motoi, *Synthesis*, 1997, in the press.
- T. Gajda and A. Zwierzak, *Synthesis*, 1981, 1005.
- R. Feinauer, *Synthesis*, 1971, 16.
- C. Lu and G. Odian, *J. Polym. Sci., Part A: Polym. Chem.*, 1994, **32**, 2283.
- W. Seeliger, E. Aufderhaar, W. Diepers, R. Feinauer, R. Nehring, W. Thier and H. Hellmann, *Angew. Chem., Int. Ed. Engl.*, 1966, **5**, 875 and references cited therein; Y. Ito, Y. Inubushi, M. Zenbayashi, S. Tomita and T. Saegusa, *J. Am. Chem. Soc.*, 1973, **95**, 4447.

Received in Cambridge, UK, 5th August 1997; 7/05679F