

The synthesis of α -amino- γ -substituted adipic acids: stereoelectronic effects during the 1,4-addition of organocuprates to methyl *N*-*tert*-butoxy-6-oxo-1,2,3,6-tetrahydropyridine-2-carboxylate

Mireille Muller, Angèle Schoenfelder, Bruno Didier, André Mann* and Camille-Georges Wermuth

Laboratoire de Pharmacochimie de la Communication Cellulaire, ERS 655, Faculté de Pharmacie, 74, route du Rhin, BP 24, F-67401 Illkirch Cedex, France. E-mail: andre.mann@pharma.u-strasbg.fr

Received (in Liverpool, UK) 26th November 1998, Accepted 4th March 1999


The α,β -unsaturated lactam **3** was submitted to 1,4-addition of organocuprate reagents R_2CuLiI_2 ($R = Me, Bu$ or Ph) to provide the 4-substituted compounds **5a–c**, **6a–c** as a mixture of diastereomers; these intermediates were then transformed into the corresponding α -amino- γ -substituted adipic acids **7a–c**, **8a–c**.

The synthesis of modified and unnatural amino acids is a source of constant inspiration in organic chemistry.^{1–3} In recent years glutamic acid, a major neurotransmitter in the mammalian brain, has received considerable attention from chemists involved in neurosciences.^{4,5} During earlier studies on modified glutamates, we and others have found that the γ -substituted analogues of structure **1** are invaluable tools for the pharmaco-

logical exploration of excitatory amino acid receptors.^{6,7} Interestingly it was found that the biological activity of such derivatives was largely confined to the 2*S*,4*S* diastereomers. For structure–activity studies devoted to this class of amino acids, we decided to synthesize compounds of type **2** in which the distal carboxylate is moved an additional methylene unit away from the α -amino acid group compared with **1**, generating γ -substituted α -amino adipic acids (Adi). As earlier observations have already demonstrated that Adi itself possesses good affinity for the glutamate receptors,⁸ the introduction of different substituents on the Adi backbone may contribute to a better knowledge of steric demands at the receptor level.⁹ Our goal was to employ an enantiospecific and diastereoselective synthesis which would provide analogues **2** in which the C-2 and C-4 substituents have the correct configuration for high biological activity. Since we have previously prepared modified glutamic acids using a conjugate addition of organocopper reagents,¹⁰ we therefore proposed to submit compound **3** to a similar sequence, in the expectation that the presence of the C-2 carboxylate would provide facial selectivity.^{11,12} We report herein our observations on the diastereoselectivity obtained in the 1,4-addition of organocuprate reagents to the α,β -unsaturated lactam **3**.

The starting compound **3** was obtained from (*S*)-lysine in enantiomerically pure form *via* piperidone **4**, following literature reports.^{13–15} The unsaturation was introduced by a sequence involving selenation and oxidative elimination, as described for the preparation of the corresponding benzyl ester (Scheme 1).¹⁵ Since lactam **3** is potentially prone to nucleophilic attack, we were pleased to find that the reaction of this substrate in Et_2O with different organocuprates gave the expected adducts as mixtures of *cis/trans* diastereomers (**5a–c** and **6a–c**) in acceptable yields, and in moderate to good stereoselectivity.[†] Examination of the ¹H NMR (200 MHz) spectra of the crude mixtures revealed that each diastereomer

has a well resolved signal (dd) between δ 4.5–4.8 for the hydrogen attached at C-2. Surprisingly, in analyzing these signals for the **5a/6a**, **5b/6b** and **5c/6c** mixtures, we observed that the intensity for the upfield signal (*trans* diastereomer) increases at the expense of the corresponding downfield one (*cis* diastereomer). These results indicate a change in the *cis:trans* ratio related to the structure of the transferred substituent. After purification, the pattern of the H-5ax proton was identified in each diastereomer as a doublet of doublets (B part of an ABX spin system) in which largest *J* value (around 17 Hz), corresponds to the geminal coupling H-5ax/H5eq. The smaller *J* value (around 11 Hz) arises from the coupling between H-5ax and H-4, and these were of comparable value in all the compounds (**5a–c** and **6a–c**), consistent with an axial disposition of the proton at C-4. As a consequence the Me, Bu or Ph substituents attached at C-4 have an equatorial disposition in each diastereomer. For the protons attached at C-2 in **5a** and **6a** two sets of coupling constants were measured: *J* = 10 and 6 Hz, and *J* = 4 and 2 Hz, respectively. From these data it was clear that in the analogues **5a–c** the carboxylate adopts an equatorial disposition (large *J* values = axial orientation of the proton at C-2) whilst in contrast for derivatives **6a–c** the carboxylate has



Scheme 1 Reagents and conditions: i, LiHMDS (1.1 equiv.), PhSeCl (1.2 equiv.), THF, $-78^\circ C$, 2 h (78%); ii, MCPBA (2 equiv.), CH_2Cl_2 , $25^\circ C$, 12 h (72%); iii, for **5a/6a**: MeLi (2 equiv.), CuI (1 equiv.), Et_2O , $-78^\circ C$, 2 h, **5a** (68%) and **6a** (9%); for **5b/6b**: BuLi (2 equiv.), CuI (1 equiv.), $-78^\circ C$, 2 h, **5b** (35%) and **6b** (33%); for **5c/6c**: PhLi (2 equiv.), CuI (1 equiv.), Et_2O , $-78^\circ C$, 2 h, **6c** (48%); HCl in AcOH, reflux, 3 h (from **45** to 80%).

an axial orientation (small J values = equatorial orientation of the proton at C-2). These data are consistent with a chair conformation for all of the isolated compounds, in which the products **5a–c** have the substituents in a 1,3-*cis* arrangement, and **6a–c** in a 1,3-*trans* arrangement. In order to explain these findings, we performed calculations on lactam **3** (MOPAC out of SYBYL 6.3) which demonstrated that the conformer having the carboxylate at C-2 in an axial disposition is, by far, more stable (5 kcal mol⁻¹) than the corresponding equatorial conformer[‡] (Scheme 1). This result is the obvious consequence of a strong A^{1,3} strain between the carboxylate at C-2 and the Boc group on the nitrogen atom.¹⁶ Based on these calculations and several literature reports on the stereoelectronic effects during conjugated addition in related α,β -unsaturated cyclohexanones or 2,3-dihydro-4-pyridones,^{17,18} a strong axial bias was expected to operate during the addition of nucleophilic reagents to lactam **3**. Therefore one should observe the preferential formation of the *cis* adducts (**5a–c**) at the expense of the corresponding *trans* epimers (**6a–c**). Indeed for the smallest substituent (R = Me) an appreciable diastereoselectivity in favor of the *cis* adduct was observed (**5a** : **6a** = 5 : 1). However when the steric demand of the organocuprate was increased (R = Bu), the facial selectivity was lost (**5b** : **6b** = 1 : 1) or, for still bulkier groups (R = Ph) the facial stereoselectivity was reversed, and the *trans* diastereomer was the only one detected (**5c** : **6c** = 1 : 30). Allinger has discussed a closely related example involving the 1,4-addition of Grignard reagents to 5-methylcyclohex-2-enone,¹⁹ in which the importance of the conformation of the conjugated enolate (chair or boat) produced by parallel or anti-parallel attack was considered. As the carboxylate function in **3** is assumed to be axial, the major pathway for the reaction with Me₂CuLi₂I was *via* an anti-parallel attack to give the transient enolate of **5a**. This intermediate possesses a low energy chair-like conformation (TS chair), in spite of the 1,3-diaxial interaction between the carboxylate and the incoming methyl group. When the nucleophile is larger, such as in Bu₂CuLi₂I (**5b/6b**) or Ph₂CuLi₂I (**5c/6c**), the reaction pathway *via* parallel attack takes place preferentially due to a competing

strong 1,3-diaxial interaction between the carboxylate and the bulky incoming nucleophile. This gives a transient, boat-like (TS boat) enolate, which collapses to the *trans* chair conformer. The final transformation of **5a–c** and **6a–c** into the desired substituted adipic acids was realized under hydrolytic conditions and the amino acids **7a,b** and **8a–c** were obtained as crystalline hydrochlorides.[§]

In conclusion, the diastereoselectivity observed for the conjugate addition of organocuprates to cyclic enamide **3** could be controlled by an appropriate choice of the transferable nucleophilic reagent. The significant A^{1,3} strain inherent to the ring system and the size of the nucleophile are probably the key factors responsible for the diastereoselectivity observed in this conjugate addition reaction. Our findings demonstrate that α,β -unsaturated species **3** is an excellent substrate for the preparation of enantiomerically pure (2*S*,4*S*)- or (2*S*,4*R*)-2-amino-4-substituted adipic acids.


We thank Dr Jacques Royer (Gif sur Yvette, France) and Professor Maurizio Taddei (Sassari, Italy) for valuable discussions.

Notes and references

† Selected data for **5a**: mp 60 °C; R_f 0.30 (hexanes–Et₂O = 5 : 5); $[\alpha]_D$ –66.2 (*c* 5, CHCl₃); δ_H (200 MHz, 30 °C) 4.51 (dd, J 10.2 and 6.2, 1 H), 3.74 (s, 3 H), 2.62 (dt, J 17 and 2.4, 1 H), 2.30 (m, 1 H), 2.10 (dd, J 17 and 10, 1 H), 2.15 (m, 1 H), 1.53 (m, 1 H), 1.48 (s, 9 H), 1.02 (d, J 6, 3 H); δ_C (50 MHz) 172.1, 170.1, 151.9, 83.6, 58.6, 52.3, 42.7, 34.1, 28.3, 27.7, 26.3, 20.7. (Calc. for C₁₃H₂₁NO₅: C, 57.54; H, 7.80; N, 5.16. Found: C, 57.3; H, 7.9; N, 5.2%). For **6a**: oil; R_f 0.35 (hexanes–Et₂O = 6 : 5); $[\alpha]_D$ +20.1 (*c* 2, CHCl₃, 30 °C); δ_H (200 MHz) 4.71 (dd, J 4 and 1.8, 1 H), 3.80 (s, 3 H), 2.66 (ddd, J 16.3, 3.2 and 1.4, 1 H), 2.21 (dd, J 9.1, 4 and 1.4, 1 H), 2.09 (dd, J 16.3 and 9.6, 1 H), 1.96 (m, 1 H), 1.73 (m, 1 H), 1.51 (s, 9 H), 1.01 (d, J 4.2, 3 H); δ_C (50 MHz) 172.2, 169.9, 152.2, 83.6, 58.3, 52.6, 42.8, 33.5, 29.7, 27.9, 25.1, 21.1. (Calc. for C₁₃H₂₁NO₅: C, 57.54; H, 7.80; N, 5.16. Found: C, 57.6; H, 7.6; N, 5.1%).

‡ In compound **3**, the recorded coupling constant (dd, J 4.5) is consistent with an axial orientation of the carboxylate.

§ Preparation of **7a**: Lactam **5a** (96 mg, 0.35 mmol) was heated at reflux for 2 h in a mixture of concentrated HCl (2 ml) and AcOH (4 ml). The mixture was concentrated *in vacuo* and triturated with Et₂O, to provide a solid which was filtered to yield **7a** (60 mg, 81%) as a hygroscopic solid; mp 178 °C; $[\alpha]_D$ –79 (*c* 1, MeOH); m/z (FAB) 176 (M + H⁺). For **8a**: mp 196 °C; $[\alpha]_D$ +26 (*c* = 1.8, MeOH); m/z (FAB) 176 (M + H⁺).



Scheme 2 Possible transition states and recorded values for coupling constants.

- F. J. Sardina and H. Rapoport, *Chem. Rev.*, 1996, **96**, 1825.
- R. O. Duthaler, *Tetrahedron*, 1994, **50**, 1539.
- J. Jurczak and A. Golebiowski, *Chem. Rev.*, 1989, **89**, 149.
- M. G. Moloney, *Nat. Prod. Rep.*, 1998, 205.
- T. Knöpfel, R. Kuhn and A. Allgeier, *J. Med. Chem.*, 1995, **38**, 1417.
- C. G. Wermuth, A. Mann, A. Schoenfelder, R. A. Wright, B. G. Johnson, J. P. Burnett, N.G. Mayne and D. D. Schoepp, *J. Med. Chem.*, 1996, **39**, 814.
- Z.-Q. Gu, X.-F. Lin and D. P. Hesson, *Bioorg. Med. Chem. Lett.*, 1995, **5**, 1973.
- R. H. Evans, A. A. Francis and J. C. Watkins, *Brain Res.*, 1978, **148**, 536.
- J. Ezquerra, C. Pedregal, A. Escribano, M. C. Carreno and J. L. G. Ruano, *Tetrahedron Lett.*, 1995, **36**, 3247.
- I. Jako, P. Uiber, A. Mann, C. G. Wermuth, T. Boulanger, B. Norberg, G. Evrard and F. Durant, *J. Org. Chem.*, 1991, **56**, 5729.
- N. Krause and A. Gerold, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 186.
- P. Perlmutter, in *Conjugate Addition Reactions in Organic Synthesis*, Pergamon, Oxford, 1992, pp. 137–197 and references cited therein.
- S. A. Hermitage and M. G. Moloney, *Tetrahedron: Asymmetry*, 1994, **5**, 1463.
- E. Davies, T. D. Heightman, S. A. Hermitage and M. G. Moloney, *Synth. Commun.*, 1996, **26**, 687.
- P. J. Murray and I. D. Starkey, *Tetrahedron Lett.*, 1996, **37**, 1875.
- R. W. Hoffmann, *Chem. Rev.*, 1989, **89**, 1841.
- P. Deslongchamps, in *Stereoelectronic Effects in Organic Chemistry*, Pergamon, Oxford, 1983, p. 209.
- J. D. Brown, M. A. Foley and D. L. Comins, *J. Am. Chem. Soc.*, 1988, **110**, 7445.
- N. L. Allinger and C. K. Riew, *Tetrahedron Lett.*, 1966, **7**, 1269.