

# A highly efficient synthesis of triisopropylsilyldifluorobromopropyne yields a versatile *gem*-difluoromethylene building block

ZhiGang Wang and Gerald B. Hammond\*

Department of Chemistry and Biochemistry, The University of Massachusetts Dartmouth, North Dartmouth, Massachusetts 02747, USA. E-mail: ghammond@umassd.edu

Received (in Corvallis, OR, USA) 23rd September 1999, Accepted 8th November 1999

**Triisopropylsilyldifluorobromopropyne, readily prepared in excellent yield from the reaction of lithium triisopropylsilylacetylide with  $\text{CF}_2\text{Br}_2$ , provides a convenient entry into a functionalized  $\text{CF}_2$  synthon.**

The *gem*-difluoromethylene unit is a key structural motif in many fluorine containing compounds of biological and pharmaceutical significance. For this reason, it is an important synthetic target.<sup>1</sup> Two complementary approaches to such important unit exist. These are (i) substitution of a carbonyl or an active methylene group by fluorine;<sup>2</sup> and (ii) use of small *gem*-difluoromethylene-containing building blocks.<sup>3</sup> When it comes to fluorinate complex molecules, the latter approach is preferred because of the reactivity, thermal instability, hazards and cost associated with electrophilic and nucleophilic fluorinating agents. The two most frequently used *gem*-difluoromethylene synthons,  $(\text{EtO})_2\text{P}(\text{O})\text{CF}_2\text{Br}$  and  $\text{EtOC}(\text{O})\text{CF}_2\text{Br}$ , were developed in the late 70s and early 80s by Burton<sup>4</sup> and Fried,<sup>5</sup> respectively. Stimulated by our earlier work in the synthesis of fluorinated phosphonates,<sup>6</sup> we sought a new generation of difluorinated building blocks from inexpensive industrial fluorine feedstock such as  $\text{CF}_2\text{Br}_2$  (not included in the list of CFCs to be phased out). Our initial target, triisopropylsilyldifluoropropyne **1** (Fig 1), is a highly functionalized three-carbon backbone that contains a propargyl silane moiety. This feature should facilitate multiple synthetic conversions containing the *gem*-difluoromethylene unit.

A literature search revealed that difluoropropargyl substrates, without exception, have been prepared in disappointingly low yields.<sup>7</sup> The low yields obtained have probably contributed to the lack of use of the difluoropropargyl building block in the literature. To our satisfaction, **1** was very efficiently assembled in one step (92% GC-MS, 81% isolated) by the reaction of  $\text{CF}_2\text{Br}_2$  with lithium triisopropylsilylacetylide. This reaction has been carried out on 5, 16 and 50 g scales and the results have shown excellent reproducibility.<sup>8</sup> We attributed the success of this reaction to the presence of the TIPS group, which possesses remarkably different properties compared to other alkylsilyl groups.<sup>9</sup> Chiefly, the presence of TIPS enhances the stability of the triisopropylsilyldifluoropropyne anion intermediate.<sup>10</sup> Although the reaction mechanism is still unclear, our experimental observation of a typical 5–10 min induction period supports the

difluorocarbene based ionic chain path proposed by Wakselman and co-workers.<sup>7a</sup> With a highly efficient preparation of **1** in hand, we explored the synthesis of various *gem*-difluoromethylene-containing compounds. Preliminary results, summarized in Scheme 1,<sup>11</sup> unveiled a highly versatile building block.

Reduction with  $\text{LiAlH}_4$  gave allene **2**, whereas nucleophilic substitution with  $\text{MP}(\text{O})(\text{OEt})_2$  afforded difluoropropyne **3** and difluoropropargyl phosphonate **4**, although the latter was obtained in low yield. Because compound **4** is a potential precursor for isosteric and isoelectronic phosphate mimics of enzyme inhibitors,<sup>12</sup> an optimization of this reaction is in progress. Using ultrasound, **1** reacted rapidly with zinc dust yielding the alkynyl organozinc reagent **5** *in situ*. This organozinc intermediate is a useful synthetic building block as can be seen in the Reformatsky-type reactions shown at the bottom of Scheme 1. Addition of Zn to **1** produced dimer **6** in nearly quantitative yield. Compound **6** is a potential intermediate in the synthesis of  $\text{CF}_2\text{CF}_2$ -containing bioactive molecules.<sup>13</sup> If **5** is quenched with powdered iodine, it produces difluoroiodopropyne **7**, another important *gem*-difluoromethylene synthon.<sup>14</sup> Addition of *trans*-cinnamaldehyde to **5** afforded difluoro alcohol **8** in 70% yield. When needed, the TIPS-protecting group can be easily removed, as demonstrated by the conversion of **8** to **9** under mild conditions, and in excellent yield. The latter result will allow an easy entry to the preparation of propargylic, and possibly allylic,  $\alpha,\alpha$ -difluorocarbonyl compounds, after oxidation of the alcohol and hydrogenation of the triple bond.




Fig. 1 Building block potential of **1**.

**Scheme 1** Reagents and conditions: i, BuLi,  $\text{CF}_2\text{Br}_2$ , THF,  $-20^\circ\text{C}$  to room temp.; ii,  $\text{LiAlH}_4$  (0.6 equiv.), THF,  $-80^\circ\text{C}$ ; iii, NaN (TMS)<sub>2</sub>,  $\text{HP}(\text{O})(\text{OEt})_2$  (1.0 equiv.), THF,  $-10^\circ\text{C}$ ; iv,  $\text{HP}(\text{O})(\text{OEt})_2$ , BuLi (1.5 equiv.),  $-10^\circ\text{C}$ ; v, Zn (1.2 equiv.), THF, room temp., ultrasound; vi,  $\text{I}_2$  (1 equiv.),  $0^\circ\text{C}$ ; vii, *trans*-cinnamaldehyde, 20 h, room temp.; viii, **1** (1.0 equiv.), Zn (2.0 equiv.), room temp.; ix, TBAF (1.0 equiv., 1 M in THF),  $-80^\circ\text{C}$ .

Other synthetic modifications of **1** and the preparation of a monofluorinated counterpart to **1** are under investigation.

The generous support of the National Science Foundation (CHE-9711062), Petroleum Research Fund (PRF#32595-B1) and the Camille and Henry Dreyfus Foundation (TH-96-012) is gratefully appreciated.

## Notes and references

- 1 For a recent leading reference see: K. Uneyama, G. Mizutani, K. Maeda and T. Kato, *J. Org. Chem.*, 1999, **64**, 6717. For a recent review see: M. J. Tozer and T. F. Herpin, *Tetrahedron*, 1996, **52**, 8619.
- 2 *Organo-Fluorine Compounds*, ed. B. Baasner, H. Hagemann and J. C. Tatlow, Thieme, Stuttgart, 1999, vol. E 10 a.
- 3 For an updated compilation of references see: J. M. Percy, *Top. Curr. Chem.*, 1997, **193**, 131.
- 4 D. J. Burton and R. M. Flynn, *J. Fluorine Chem.*, 1977, **10**, 329. For recent reviews, see: D. J. Burton and L. Lu, *Top. Curr. Chem.*, 1997, **193**, 45; D. J. Burton, Z. Y. Yang and W. M. Qiu, *Chem. Rev.*, 1996, **96**, 1641.
- 5 E. A. Hallinan and J. Fried, *Tetrahedron Lett.*, 1984, **25**, 2301. Also: T. Taguchi, O. Kitagawa, Y. Suda, S. Ohkawa, A. Hashimoto, Y. Iitaka and Y. Kobayashi, *Tetrahedron Lett.*, 1988, **29**, 5291 and references therein; S. Mcharek, S. Sibille, J. Y. Nedelec and J. Perichon, *J. Organomet. Chem.*, 1991, **401**, 211; J. M. Altenburger and D. Schirlin, *Tetrahedron Lett.*, 1991, **32**, 7255; J. M. Andres, M. A. Martinez, R. Pedrosa and A. Perez-Encabo, *Synthesis*, 1996, 1070.
- 6 F. Benayoud, D. J. deMendonca, C. A. Digits, G. A. Moniz, T. C. Sanders and G. B. Hammond, *J. Org. Chem.*, 1996, **61**, 5159; F. Benayoud and G. B. Hammond, *Chem. Commun.*, 1996, 1447.
- 7 (a) I. Rico, D. Cantauzene and C. Wakselman, *J. Chem. Soc., Perkin. Trans. 2*, 1982, 1063; (b) P. Y. Kwok, F. W. Muellner, C. K. Chen and J. Fried, *J. Am. Chem. Soc.*, 1987, **109**, 3684; (c) Y. Hanzawa, K. Inazawa, A. Kon, H. Aoki and Y. Kobayashi, *Tetrahedron Lett.*, 1987, **28**, 659.
- 8 Under Ar atmosphere, a solution of triisopropylsilylacetylene (36 g, 0.2 mol) in THF (300 ml) was cooled to  $-20^{\circ}\text{C}$ . BuLi (125 ml, 1.6 M solution in hexane) was added *via* syringe and the resulting solution was stirred for 30 min before  $\text{CF}_2\text{Br}_2$  (62 g, 1.5 equiv.) was added. The solution was allowed to warm to room temperature and stirred for 9 h. After solvent removal,  $\text{NH}_4\text{Cl}$  (120 ml) was added and the oily layer was extracted with  $\text{Et}_2\text{O}$  ( $3 \times 100$  ml). Standard work-up gave the crude product as a viscous orange oil (92% yield by GC-MS) which was purified by distillation (bp  $51\text{--}52^{\circ}\text{C}/0.1$  mmHg) to yield **1** (49.9 g, 81%);  $\delta_{\text{H}}$ (300 MHz,  $\text{CDCl}_3$ ) 1.11 (s, TIPS protons);  $\delta_{\text{C}}$ (75 MHz,  $\text{CDCl}_3$ ) 100.61 (t,  $^1J_{\text{CF}}$  290), 97.16 (t,  $^2J_{\text{CF}}$  36.5), 95.1 (t,  $^3J_{\text{CF}}$  4.7), 18.01, 11.28;  $\delta_{\text{F}}$   $-32.60$  (s, 2F);  $m/z$  (GC-MS) 312 ( $\text{M}^+ + 1, 4$ ), 310 ( $\text{M}^+ - 1, 4$ ), 269 (14), 267 (14), 143 (27), 77 (100) (calc. for  $\text{C}_{12}\text{H}_{21}\text{SiF}_2\text{Br}$ : C, 46.33; H, 6.75. Found: C, 46.91; H, 6.83%). The triisopropylsilylacetylene starting material, purchased from GFS Co., contained diisopropylpropenylsilylacetylene (11%) and diisopropylpropylsilylacetylene (14%). Fractional distillation did not remove these impurities. The GC-MS analysis of **1** shows all three components were alkylated and their ratios perfectly match that of the starting material. The calculated C% and H% were based on the formula of triisopropylsilyldifluorobromopropyne. If the two impurities are removed from the calculation, the values found for C% and H% are 46.84 and 6.67%, respectively.
- 9 For a recent review of the TIPS group in organic chemistry see: C. Rücker, *Chem. Rev.*, 1995, **95**, 1009.
- 10 Similarly, whereas ynolates lack convenient methods for their generation a lithium silyl ynolate has been used in ketylation reactions: H. Kai, K. Iwamoto, N. Chatani and S. Murai, *J. Am. Chem. Soc.*, 1996, **118**, 7634.
- 11 Selected data for **2**:  $\delta_{\text{H}}$ (300 MHz,  $\text{CDCl}_3$ ) 1.06 (s, TIPS protons), 4.81 (t, 1H,  $^4J_{\text{HH}}$  7.23), 4.30 (d, 2H,  $^4J_{\text{HH}}$  7.23);  $m/z$  (GC-MS) 196 ( $\text{M}^+$ ), 157, 153, 125, 97, 83, 67. For **3**:  $\delta_{\text{H}}$ (300 MHz,  $\text{CDCl}_3$ ) 1.09 (s, TIPS protons), 6.17 (t,  $^2J_{\text{HF}}$  54.8);  $\delta_{\text{F}}$   $-105.49$  (s);  $m/z$  (GC-MS) 232 ( $\text{M}^+$ ), 189, 161, 133, 105, 81, 77. For **4**:  $\delta_{\text{H}}$ (300 MHz,  $\text{CDCl}_3$ ) 1.11 (s, 21H, TIPS protons), 1.39 (t, 6H,  $^3J_{\text{HH}}$  7.00), 4.32 (m, 4H);  $\delta_{\text{F}}$   $-96.79$  (d,  $^2J_{\text{FF}}$  109.01);  $\delta_{\text{P}}$  4.23 (t,  $^2J_{\text{PF}}$  109.10);  $m/z$  (GC-MS) 325 ( $\text{M}^+ - 43$ ), 297, 269, 153, 109, 81. For **6**:  $\delta_{\text{H}}$ (300 MHz,  $\text{CDCl}_3$ ) 1.10 (s, TIPS protons);  $\delta_{\text{F}}$   $-99.01$  (s, 4F);  $m/z$  (GC-MS) 462 ( $\text{M}^+$ ), 377, 307, 239, 183, 115, 77. For **7**:  $\delta_{\text{H}}$ (300 MHz,  $\text{CDCl}_3$ ) 1.11 (s, TIPS protons);  $\delta_{\text{F}}$   $-28.13$  (s);  $m/z$  (GC-MS) 315 ( $\text{M}^+ - 43$ ), 265, 237, 189, 165, 119, 105, 77. For **8**:  $\delta_{\text{H}}$ (300 MHz,  $\text{CDCl}_3$ ) 7.41–7.26 (m, 5H), 6.82 (d, 1H,  $^3J_{\text{HH}}$  15.9), 6.26 (dd, 1H,  $^3J_{\text{HH}}$  15.9,  $^3J_{\text{HH}}$  6.2), 4.55 (m, 1H, CHOH), 2.05 (s, 1H, OH), 1.06 (s, 21 H, TIPS protons);  $\delta_{\text{F}}$   $-94.3$  (d, 1F,  $^2J_{\text{FF}}$  274),  $-96.7$  (d, 1F,  $^2J_{\text{FF}}$  274). For **9**:  $\delta_{\text{H}}$ (300 MHz,  $\text{CDCl}_3$ ) 7.45–7.30 (m, 5H), 6.85 (dd, 1H,  $^4J_{\text{HH}}$  0.9,  $^3J_{\text{HH}}$  15.9), 6.24 (dd, 1H,  $^3J_{\text{HH}}$  15.9,  $^3J_{\text{HH}}$  6.3), 4.55 (m, 1H, CHOH), 2.85 (t, 1H,  $^3J_{\text{HF}}$  5.0), 2.04 (s, 1H);  $\delta_{\text{F}}$   $-95.9$  (d, 1F,  $^2J_{\text{FF}}$  277),  $-96.9$  (d, 1F,  $^2J_{\text{FF}}$  277);  $m/z$  (GC-MS) 206 ( $\text{M}^+ + 5$ ), 170 (3), 159 (2), 133 (100), 115 (30), 77 (26), 55 (46).
- 12 The major by-product is **3**, an indication that, compared to the bromide, the triisopropylsilyldifluoropropyne anion is a better leaving group. For a leading reference on fluorinated phosphate mimics, see: D. O'Hagan and H. S. Rzepa, *Chem. Commun.*, 1997, 645.
- 13 W. C. Sun, C. S. Ng and G. D. Prestwich, *J. Org. Chem.*, 1992, **57**, 132.
- 14 We have also prepared **8** in 73% yield by the reaction of lithium triisopropylsilylacetylide with  $\text{CF}_2\text{I}_2$ . However, the prohibitive cost of commercial  $\text{CF}_2\text{I}_2$  will most likely limit its use in large-scale synthesis.

Communication 9/07784G