

The effect of solvent on the α -effect: the MeCN–H₂O solvent system†

Ik-Hwan Um,^a Young-Min Park^a and Erwin Buncel*^b

^a Department of Chemistry, Ewha Womans University, Seoul 120-750, Korea

^b Department of Chemistry, Queen's University, Kingston, Ontario K7L 3N6, Canada.

E-mail: buncel@chem.queensu.ca; Tel: 613-533-2635; Fax: 613-533-6669


Received (in Corvallis, OR, USA) 10th July 2000, Accepted 23rd August 2000

First published as an Advance Article on the web 18th September 2000

The increasing α -effect observed in MeCN–H₂O solvent mixtures, which contrasts with the previously found bell-shaped dependence on solvent composition in the DMSO–H₂O system, is attributed to the differential solvent effect on basicities of the α - and normal nucleophiles in the former case.

Among the possible causes of the α -effect, that is the enhanced reactivity of nucleophiles having an unshared pair of electrons adjacent to the nucleophilic centre,¹ the effect of solvent has been particularly controversial.^{2–9} As one possible approach to the problem, in 1986 we examined the reactivities of an α -nucleophile, butane-2,3-dione monoximate (Ox[−]), and a normal nucleophile, *p*-chlorophenoxide (ClPhO[−]) with *p*-nitrophenyl acetate (PNPA) in dimethyl sulfoxide (DMSO)–H₂O mixtures,^{7a} due to the unique properties of this solvent system.¹⁰ Unexpectedly, we observed a bell-shaped profile for the dependence of the α -effect on solvent composition, with a maximum at ca. 50 mol % DMSO.^{7a} More recently, we observed a similar bell-shaped α -effect trend for the corresponding reactions of *p*-nitrophenyl diphenyl phosphinate (PNPDP) .^{7b} However, a contrary result was reported for the reaction of PNPA with *O*-iodosylbenzoate (IBO[−]) and ClPhO[−]: the α -effect showed no maximum but decreased steadily as the DMSO content in the reaction medium increased.⁹ Clearly, more work is called for in order to understand the effect of solvent on the α -effect.

We have now extended our study to a different solvent system, acetonitrile (MeCN)–H₂O mixtures, and have found a new type of solvent effect on the α -effect, namely an increasing α -effect. A plausible cause of this trend for the reaction of PNPA with Ox[−] and ClPhO[−] (eqn. 1) in MeCN–H₂O mixtures is herein presented.



Kinetic studies were performed spectrophotometrically under pseudo-first-order conditions with the nucleophile in excess. Pseudo-first-order rate constants (k_{obs}) were obtained from linear plots of $\ln(A_\infty - A_t)$ vs. t . Second-order rate constants (k^{Ox^-} and k^{ClPhO^-}) were calculated from the slope of the linear plot of k_{obs} vs. nucleophile concentration. As shown in Fig. 1, the magnitude of the α -effect ($k^{\text{Ox}^-}/k^{\text{ClPhO}^-}$) increases as the mol % MeCN in the medium increases, from ca. 100 in H₂O to 500 in 90 mol % MeCN. Recently, we found a decreasing α -effect trend for the reaction of PNPA with hydroxamates in MeCN–H₂O mixtures, and a shift in equilibrium from an OH acid to an NH acid upon addition of MeCN to H₂O was attributed as responsible for the observed α -effect trend.^{8b}

Transition-state stabilization is one possible cause of the α -effect¹¹ and recently the bell-shaped α -effect trend observed for

the reaction of PNPA in DMSO–H₂O mixtures has been explained through a dissection of ground-state and transition-state contributions.^{7c} In that system the basicities of the α -Nu (Ox[−]) and normal-Nu (ClPhO[−]) exhibit a parallel dependence on the DMSO–H₂O composition.⁷

The addition of MeCN to H₂O could influence not only the reaction rate but also the basicity of the anionic nucleophiles. Although the pK_a values for some phenols, carboxylic acids and amines in pure MeCN are available,^{12a,b} very few pK_a data have been reported for MeCN–H₂O mixtures.^{8b} Therefore, we have measured the relative basicity of Ox[−] and ClPhO[−] in MeCN–H₂O mixtures using piperazine as a reference base. One can define ΔpK_a as the pK_a difference between the conjugate acids of the nucleophile (Ox[−] or ClPhO[−]) and piperazine, *i.e.* $\Delta pK_a = pK_a$ of the conjugate acid of the nucleophile (Ox[−] or ClPhO[−]) – pK_a of the conjugate acid of the reference base (piperazine). Hence, the magnitude of ΔpK_a represents the relative basicities of these nucleophiles. We determined ΔpK_a values spectrophotometrically using the relationship: $\Delta pK_a = \log [\text{HA}]_{\text{eq}}[\text{B}]_{\text{eq}}/[\text{A}^-]_{\text{eq}}^2$, in which $[\text{HA}]_{\text{eq}}$ represents the equilibrium concentration of the conjugate acid of the nucleophile, and $[\text{B}]_{\text{eq}}$ and $[\text{A}^-]_{\text{eq}}$ represent the equilibrium concentration of the reference base (piperazine) and the nucleophile, respectively. The ΔpK_a values thus determined in MeCN–H₂O mixtures are summarized in Table 1. As shown in the table, the ΔpK_a values of Ox[−] and ClPhO[−] in H₂O are −0.38 and −0.44, respectively. The pK_a values of the conjugate acids of Ox[−], ClPhO[−] and piperazine in H₂O at 25 °C, have been reported to be 9.44,^{7b} 9.38^{12c} and 9.82,^{12c} respectively, *i.e.* the ΔpK_a data determined in H₂O in the present study are identical to the literature values. It is also noted that Ox[−] and ClPhO[−] are less basic than piperazine in H₂O but appear to be more basic upon addition of MeCN. This is consistent with the report that the increase in pK_a values upon changing medium from H₂O to MeCN is more significant for phenols than for amines: the pK_a enhancements were reported to be 12–17 and 7–8 in pK_a units for phenols and for the conjugate acids of alicyclic secondary




Fig. 1 Plots showing the effect of solvent on the α -effect ($k^{\alpha\text{-Nu}}/k^{\text{normal-Nu}}$) for the reaction of PNPA at 25.0 °C: $k^{\text{Ox}^-}/k^{\text{ClPhO}^-}$ in MeCN–H₂O (●); $k^{\text{Ox}^-}/k^{\text{ClPhO}^-}$ in DMSO–H₂O (○);^{7b} $k^{\text{IBO}^-}/k^{\text{ClPhO}^-}$ in DMSO–H₂O (□).⁹

† Electronic supplementary information (ESI) is available: second-order rate constants for the reaction of *p*-nitrophenyl acetate with butane-2,3-dione monoximate and *p*-chlorophenoxide in MeCN–H₂O mixtures at 25 °C. See <http://www.rsc.org/suppdata/cc/b0/b005610n/>

Table 1 Summary of the relative basicity of the nucleophile ($\Delta pK_a = pK_a$ of the conjugate acid of the nucleophile— pK_a of the conjugate acid of the reference base, piperazine) in MeCN–H₂O mixtures of varying compositions at 25.0 ± 0.1 °C^a

Mol % MeCN	ΔpK_a (Ox ⁻)	ΔpK_a (ClPhO ⁻)	$\Delta\Delta pK_a^b$
0	-0.38	-0.44	0.06
10	0.58	0.43	0.15
20	1.17	0.90	0.27
30	1.80	1.44	0.36
40	2.32	1.89	0.43
50	2.69	2.19	0.50
60	3.43	2.84	0.59
70	3.90	3.20	0.70
80	4.96	4.18	0.78
90	5.85	4.90	0.95

^a The uncertainty in ΔpK_a values is estimated to be less than ±0.03 pK_a units. ^b $\Delta\Delta pK_a = \Delta pK_a$ (Ox⁻) - ΔpK_a (ClPhO⁻).

amines (e.g. piperidine and morpholine), respectively, upon solvent change from H₂O to MeCN.^{12a,b}

The difference in the relative basicity between Ox⁻ and ClPhO⁻ can be expressed as $\Delta\Delta pK_a$, i.e. $\Delta\Delta pK_a = \Delta pK_a$ (Ox⁻) - ΔpK_a (ClPhO⁻). As shown in Table 1, $\Delta\Delta pK_a$ increases as the mol % MeCN in the medium increases, i.e. Ox⁻ is more basic than ClPhO⁻ by 0.06, 0.50 and 0.95 pK_a units in H₂O, 50 and 90 mol % MeCN, respectively, which contrasts with the situation, noted above, for DMSO–H₂O mixtures. We




Fig. 2 Plot of $\log k^{Ox^-}/k^{ClPhO^-}$ vs. $\Delta\Delta pK_a$ for the reaction of PNPA with Ox⁻ and ClPhO⁻ in MeCN–H₂O mixtures at 25.0 ± 0.1 °C.

propose therefore, that the differential solvent effect on the basicity of these nucleophiles is the cause of the contrasting α -effect trends observed for the reaction of PNPA with Ox⁻ and ClPhO⁻ in DMSO–H₂O and in MeCN–H₂O mixtures.

Furthermore, it is found that a plot of $\log k^{Ox^-}/k^{ClPhO^-}$ against $\Delta\Delta pK_a$ is linear with a slope of 0.85 (Fig. 2). This provides evidence that the increase in the $\Delta\Delta pK_a$ values is almost fully reflected in the increase in the α -effect for the reaction of PNPA with Ox⁻ and ClPhO⁻ upon addition of MeCN into the reaction medium.

I. H. U. is grateful for financial support from KOSEF of Korea (1999-2-123-003-5). E. B. thanks NSERC of Canada for a research grant.

Notes and references

- J. O. Edwards and R. G. Pearson, *J. Am. Chem. Soc.*, 1962, **84**, 16.
- Reviews: (a) E. Buncl and S. Hoz, *Isr. J. Chem.*, 1985, **26**, 313; (b) A. P. Grekov and V. Y. Veselov, *Usp. Khim.*, 1978, **47**, 1200; (c) N. J. Fina and J. O. Edwards, *Int. Chem. Kinet.*, 1973, **5**, 1.
- (a) C. F. Bernasconi and C. Murray, *J. Am. Chem. Soc.*, 1986, **108**, 5251; (b) D. J. Palling and W. P. Jencks, *J. Am. Chem. Soc.*, 1984, **106**, 4869.
- (a) G. Moutiers, E. Guevel, L. Villien and F. Terrier, *J. Chem. Soc., Perkin Trans. 2*, 1997, 710; (b) F. Terrier, G. Moutiers, L. Xiao, E. Guevel and F. Guir, *J. Org. Chem.*, 1995, **60**, 1748.
- (a) I. H. Um, J. S. Lee and S. M. Yuk, *J. Org. Chem.*, 1998, **63**, 9152; (b) K. R. Fountain, T. W. Dunkin and K. D. Patal, *J. Org. Chem.*, 1997, **62**, 2738; (c) E. Buncl, C. Chuaqui and H. Wilson, *J. Am. Chem. Soc.*, 1982, **104**, 4896.
- W. L. Jorgensen, *J. Am. Chem. Soc.*, 1987, **109**, 2349.
- (a) E. Buncl and I. H. Um, *J. Chem. Soc., Chem. Commun.*, 1986, 595; (b) R. M. Tarkka and E. Buncl, *J. Am. Chem. Soc.*, 1995, **117**, 1503; (c) I. H. Um and E. Buncl, *J. Org. Chem.*, 2000, **65**, 577.
- (a) I. H. Um, D. S. Kwon and G. J. Lee, *Bull. Korean Chem. Soc.*, 1989, **10**, 620; (b) I. H. Um, H. W. Yoon, J. S. Lee and H. J. Moon, *J. Org. Chem.*, 1997, **62**, 5939.
- R. A. Moss, S. Swarup and S. Ganguli, *J. Chem. Soc., Chem. Commun.*, 1987, 860.
- E. Buncl and H. Wilson, *Adv. Phys. Org. Chem.*, 1977, **14**, 133.
- (a) E. Buncl and S. Hoz, *Tetrahedron Lett.*, 1983, **24**, 4777; (b) S. Hoz and E. Buncl, *Tetrahedron Lett.*, 1984, **25**, 3411; (c) K. R. Fountain, D. B. Tad-y, T. W. Paul and M. V. Golynskiy, *J. Org. Chem.*, 1999, **64**, 6547; (d) K. R. Fountain and K. D. Patel, *J. Org. Chem.*, 1997, **62**, 4795.
- (a) M. K. Chantooni and I. M. Kolthoff, *J. Phys. Chem.*, 1976, **80**, 1306; (b) J. F. Coetzee and G. R. Padmanabhan, *J. Am. Chem. Soc.*, 1965, **87**, 5005; (c) W. P. Jencks and J. Regenstein, in *Handbook of Biochemistry. Selected Data for Molecular Biology*, ed. H. A. Sober, The Chemical Rubber Co., Cleveland, OH, 1968.