

An unusual oxidative rearrangement of azabicyclo[2.2.1]heptenes, providing a stereoselective route to 2'- and 3'-hydroxycyclopentylglycines

Patrick D. Bailey,^{*a} I. M. McDonald,^{*b} Georgina M. Rosair^a and David Taylor^a

^a Department of Chemistry, Heriot-Watt University, Riccarton, Edinburgh, UK EH14 4AS.
E-mail: P.D.Bailey@hw.ac.uk

^b James Black Foundation, 68 Half Moon Lane, Dulwich, London, UK SE24 9JE

Received (in Liverpool, UK) 11th August 2000, Accepted 30th October 2000


First published as an Advance Article on the web

Treatment of the azabicyclo[2.2.1]heptene derivative **4** with *m*CPBA for 3–5 seconds generates the oxazabicyclo[3.2.1]octene derivative **7** (X-ray structure) via rapid Meisenheimer rearrangement of the *N*-oxide **5**, whilst heating **7** in MeCN for 4–6 h leads to further rearrangement to the more thermodynamically stable oxazabicyclo[3.3.0]octene isomer **6**; **6** and **7** can be readily reduced to enantio-pure hydroxylated cyclopentylglycine derivatives.

Pharmacologically active peptoids can be generated by attaching the side-chains of a parent peptide to carrier moieties such as sugars, steroids, or porphyrins.¹ We have been exploring the use of azabicyclo[2.2.1]heptanes as rigid templates for such systems which, if they can be derivatised appropriately, offer potential as 3D structures on which to construct peptoid libraries. We were also attracted to such systems because we had developed an extremely short, efficient, and highly stereoselective route to these systems some years ago;² the azadiels–Alder reaction of the imine **3** with cyclopentadiene gives the azabicyclo[2.2.1]heptene **4** as a single stereoisomer, and we wished to explore the derivatisation of this adduct.³ This communication, however, reports some unexpected results that have instead provided a short, efficient, stereo- and regio-specific route to hydroxylated cyclopentylglycines.

As outlined in Scheme 1, the adduct **4** could be efficiently prepared using ethyl glyoxylate generated using the procedure developed by Roberts' group.⁴ After formation of the imine, the cycloaddition reaction was found to be most reliable when conducted in trifluoroethanol in the presence of TFA (1 equiv.). As reported by ourselves⁴ and others,⁵ this reaction proceeds both with excellent *exo* diastereo-control (in contrast to the *endo* selectivity observed for acyclic dienes²), and with extremely high asymmetric induction resulting from the presence of the α -methylbenzyl auxiliary (72% isolated yield, 95:5 *exo:endo*, 90% asymmetric induction). After chromatography, **4** was obtained as a single stereoisomer, from which we hoped an epoxidation/nucleophilic ring-opening sequence would allow derivatisation at site **C**, whilst further pharmacophores could be introduced at sites **A** and **B**. However, treatment with *m*CPBA failed to give the desired epoxide (even under acidic conditions whereby the nitrogen would be expected to be protonated), but instead yielded an isomeric oxygenated product which we initially assigned as the *N*-oxide **5**. Small amounts of a by-product were also generated, which was identified as the oxazabicyclo[3.3.0]octene **6** from NMR data.[†]

We had expected that an X-ray crystal structure of the '*N*-oxide' would confirm its identity and allow us to assign its stereochemistry at the chiral nitrogen, but this surprisingly revealed that the isomeric oxazabicyclo[3.2.1]octene **7** had been formed instead (Fig. 1[‡]). Optimisation of the reaction conditions established that the highest yield of **7** was obtained when the cyclo-adduct **4** was reacted at room temperature with *m*CPBA for only 3–5 seconds and, that clean conversion to **6** could be achieved by refluxing **7** in acetonitrile for 24 h. Presumably **7** is formed via a Meisenheimer rearrangement,^{6–9} but the rapidity and selectivity of the initial reaction, in which



Scheme 1 Formation of the oxazabicyclo[3.3.0]octane system **6**. The azadiels–Alder reaction (**3** → **4**) was in TFE, TFA (1 eq.), 0 °C, 3.5 h (72% yield, 95% *exo*, 90% asymmetric induction). See text and Scheme 2 for yields and identification of the products **5** and **6** from *m*CPBA oxidation of **4**. [R* = (*S*)-1-phenylethyl].






Fig. 1 X-Ray crystal structure of **7** with crystallographic numbering.[‡]



Scheme 2 Possible mechanism for the formation of **6** and **7**; heterolytic bond cleavage of **7** would also be consistent with its thermolysis to **6**. [$R^* = (S)$ -1-phenylethyl].



Scheme 3 Conversion of the oxazabicyclooctanes **6** and **7** into hydroxylated cyclopentylglycines such as **9** and **10**. Conditions: (i) Zn–AcOH; (ii) H_2 (50 psi)–Pd(OH) $_2$ /C; (iii) MeCN, reflux, 24 h. [$R^* = (S)$ -1-phenylethyl].

the di-radical must be retained within a solvent cage, is extraordinary. Unprecedented, as far as we are aware, is the establishment of an equilibrium with the Meisenheimer intermediate (or possibly *via* heterolytic bond cleavage of **7**), providing a route by which the thermodynamically more stable adduct **6** can be generated (see Scheme 2).

With an efficient route to two isomeric oxazabicyclooctanes achieved, we wished to explore whether they could be converted into hydroxy-substituted cyclopentylglycines. The key transformations are shown in Scheme 3. In particular, Zn–AcOH treatment of the [3.2.1] isomer **7** allowed selective cleavage of the N–O bond, yielding the 4'-hydroxycyclopent-2'-enyl glycine derivative **8**. Hydrogenation of the double bond (Pearlman's catalyst, 50 psi) resulted in concomitant removal of the α -methylbenzyl auxiliary to afford the 3'-hydroxycyclopentyl glycine **9** as a single (2*R*,1'*R*,3'*S*) stereoisomer; the same hydrogenation conditions also allowed direct conversion of **7** to **9** (90% yield). On the other hand, these hydrogenation conditions enabled us to reduce the [3.3.0] isomer **6** directly to the 2'-hydroxycyclopentyl glycine **10**, also as a single (2*R*,1'*R*,2'*R*) stereoisomer. Closely related analogues of **8** have been used in the synthesis of cyclopentylglycine,¹¹ and of carbocyclic analogues of nikkomycin¹² and polyoxin C.¹³

In summary, the asymmetric aza-Diels–Alder adduct **4** reacts with *m*CPBA to generate the Meisenheimer rearranged product

6, which can be converted into the [3.3.0] system **7** by simple heating. These two isomers can be separately reduced, to provide access to a wide range of cyclopentylglycine derivatives, with complete regio-, diastereo- and enantio-control.

We thank Dr A. S. F. Boyd and Dr R. Ferguson for NMR and mass spectra, the EPSRC for financial support, and we acknowledge the use of the EPSRC's Chemical Database Service at Daresbury.

Notes and references

† Selected data for **6**: $C_{17}H_{21}NO_3$; δ_H (CDCl $_3$) 1.1 (3H, t, *J* 7.2), 1.5 (3H, d, *J* 6.6), 2.4 (2H, m), 3.05 (1H, q, *J* 6.6), 3.55 (1H, m), 3.7 (1H, d, *J* 9.1), 3.95 (2H, q, *J* 7.2), 5.3 (1H, dd, *J* 7.4, 0.7), 5.8 (1H, dd, *J* 3.5, 1.7), 5.95 (1H, dd, *J* 1.7, 0.7), 7.2–7.4 (5H, m); δ_C (CDCl $_3$) 14.15 (CH $_3$), 20.47 (CH $_3$), 34.16 (CH $_2$), 47.20 (CH), 60.52 (CH $_2$), 65.68 (CH), 69.85 (CH), 87.35 (CH), 127.66 (CH), 128.25 (2 \times CH), 128.30 (2 \times CH), 129.49 (CH), 135.17 (CH), 141.91 (C), 169.90 (C).

‡ *Crystal data* for **7**. A single crystal of **7** was grown by slow evaporation of 40–60 pet. ether at rt and coated in Nujol and mounted on a glass fibre covered with vacuum grease for data collection with a Bruker P4 diffractometer¹⁴ at 160 K. Colourless plate, 0.12 \times 0.72 \times 0.34 mm, $C_{17}H_{21}NO_3$, $M = 287.35$, monoclinic, space group $P2_1$, $a = 6.0734$ (4), $b = 13.0445$ (9), $c = 10.1045$ (7) Å, $\beta = 98.280$ (5) $^\circ$, $U = 792.18$ (13) Å 3 , $Z = 2$, μ (Mo–K α) = 0.082 mm $^{-1}$. Data measured = 3180, unique data = 1593, $R = 0.0778$ on all data ($wR_2 = 0.0306$). Absolute structure parameter = 3.5(19), so the absolute structure could not be confirmed from the X-ray data. Crystallographic computing was performed using the SHELXTL¹⁵ system version 5.1. CCDC 182/1836. See <http://www.rsc.org/suppdata/b0/b006683o/> for crystallographic data in .cif format.

- R. Hirschmann, *Angew. Chem., Int. Ed. Engl.*, 1991, **30**, 1278.
- P. D. Bailey, R. D. Wilson and G. R. Brown, *J. Chem. Soc., Perkin Trans. 1*, 1991, 1337; P. D. Bailey, G. R. Brown, F. Korber, A. Reid and R. D. Wilson, *Tetrahedron: Asymmetry*, 1991, **2**, 1263.
- Andersson's group has made extensive use of the azabicyclo-[2.2.1]heptene **4**, as exemplified by P. Pinho and P. G. Andersson, *Chem. Commun.*, 1999, 597, and reference 5 therein; the Malpass group has also developed the chemistry of these azabicyclic systems, as exemplified by J. R. Malpass and C. D. Cox, *Tetrahedron Lett.*, 1999, **40**, 1419.
- M. B. Hursthouse, K. M. A. Malik, D. E. Hibbs, S. M. Roberts, A. J. H. Seago, V. Sik and R. Storer, *J. Chem. Soc., Perkin Trans 1*, 1995, 2419.
- L. Stella, H. Abraham, J. Feneu-Dupont, B. Tinant and J. P. Declercq, *Tetrahedron Lett.*, 1990, **31**, 2603; H. Abraham and L. Stella, *Tetrahedron*, 1992, **48**, 9707.
- For a useful discussion of the Meisenheimer rearrangement, see V. Rautenstrauch, *Helv. Chim. Acta*, 1973, **56**, 2492, and references 8–18 therein.
- For a study of the 2,3-Meisenheimer rearrangement of allyl *N*-oxides, see Y. Yamamoto, J. Oda and Y. Inouye, *J. Org. Chem.*, 1976, **41**, 303, and references therein; it might be noted that this process is believed to be pericyclic rather than radical, and that **6** is (formally) the product of the 2,3-Meisenheimer rearrangement of **5**.
- For the synthesis of 7-, 8- and 9-membered 1,2-oxaza ring systems *via* the Meisenheimer rearrangement, see J. B. Bremner, E. J. Browne, P. E. Davies and L. van Thuc, *Aust. J. Chem.*, 1980, **33**, 833; J. B. Bremner, E. J. Browne, P. E. Davies, C. L. Raston and A. H. White, *Aust. J. Chem.*, 1980, **33**, 1323; J. B. Bremner, E. J. Browne and P. E. Davies, *Aust. J. Chem.*, 1980, **33**, 1335.
- Spontaneous Meisenheimer rearrangements are rare, but has been observed in the ring expansion of a highly strained azetidene *N*-oxide: T. Kurihara, Y. Sakamoto, K. Tsukamoto, H. Ohishi, S. Harusawa and R. Yoneda, *J. Chem. Soc., Perkin Trans 1*, 1993, 81.
- O. Mitsunobu, *Synthesis*, 1981, 1.
- N. Katagiri, M. Okada, Y. Morishita and C. Kaneko, *Chem. Commun.*, 1996, 2137.
- S. E. Ward, A. B. Holmes and R. McCague, *Chem. Commun.*, 1997, 2085.
- D. Zhang and M. J. Miller, *J. Org. Chem.*, 1998, **63**, 755; V. K. Aggarwal and N. Monteiro, *J. Chem. Soc., Perkin Trans. 1*, 1997, 2531.
- XSCANS X-ray Single Crystal Analysis System. Version 2.2. Bruker AXS Inc., Madison, Wisconsin, USA, 1994.
- G. M. Sheldrick, 1999. SHELXTL Version 5.1, Bruker AXS Inc., Madison, Wisconsin, USA.