

A novel radical cyclization of 2-bromoindoles. Synthesis of hexahydropyrrolo[3,4-*b*]indoles


Gordon W. Gribble,* Heidi L. Fraser and Jeanese C. Badenock

Department of Chemistry, Dartmouth College, Hanover, NH 03755-3564, USA.
 E-mail: grib@dartmouth.edu

Received (in Corvallis, OR, USA) 16th February 2001, Accepted 14th March 2001
 First published as an Advance Article on the web 17th April 2001

Hexahydropyrrolo[3,4-*b*]indoles **6**, **10**, and **13** are obtained from 2-bromo-3-carboxamides **5**, **9**, and **12**, respectively, by a 1,5-radical translocation process followed by 5-*endo-trig* cyclization to the indole C-2 position.


We wish to describe a new synthesis of hexahydropyrrolo[3,4-*b*]indoles from 2-bromoindole-3-carboxamides involving sequential indole C-2 radical generation, 1,5-hydrogen atom abstraction, and 5-*endo-trig* cyclization to the indole C-2 position. In connection with our interest in pyrrolo[3,4-*b*]indoles¹ and indolo[2,3-*a*]quinolizidines,² we envisioned the free radical sequence shown in Scheme 1 as an attractive route to these ring systems.




Scheme 1

We herein report the viability of this sequence for the synthesis of hexahydropyrrolo[3,4-*b*]indoles. The radical precursor 2-bromoindole **5** was prepared as shown in Scheme 2. Indole (**1**) was converted to amide **3** in two steps as previously described.³ Subsequent *N*-methylation and C-2 bromination using the conditions described by Bergman⁴ for *N*-carboxyindole afforded **5**† in excellent overall yield. An X-ray crystal structure determination confirmed the structure of **5**.⁵

The radical cyclization was performed by the slow addition over 36 h of a degassed solution of tri-*n*-butyltin hydride and catalytic AIBN in toluene to a refluxing solution of bromoindole **5** in toluene. This resulted in the formation of the desired




Scheme 2 Reagents and conditions: i, (CF₃CO)₂O, Et₂O, 0 °C (87%); ii, BuLi, piperidine, THF, 0 °C (97%); iii, NaOH, EtOH, MeI, acetone, rt (99%); iv, Bu^tLi, THF, BrCl₂CCl₂Br, -78 °C to rt (88%).



Scheme 3 Reagents and conditions: i, Bu₃SnH, AIBN, toluene, reflux, 36 h (**6**, 54%; **4**, 42%).

dihydroindole **6**† (54%), along with the reduction product **4** (42%) (Scheme 3). The structure of **6** is fully supported by spectral and analytical data, including an X-ray crystal structure determination.⁵

We believe that this reaction involves the sequence (1) generation of the expected C-2 radical, (2) 1,5-H atom abstraction to give the α-amidoyl radical, (3) 5-*endo-trig* cyclization to the indole double bond, and (4) hydrogen abstraction to give indoline **6**. The first two steps in this process have been termed ‘radical translocation’ by Snieckus and Curran.⁶ Attempts to improve the yield of **6** relative to reduction product **4** using other radical generation methods have not been successful. Treatment of indoline **6** with DDQ (CH₂Cl₂, rt) gave indole **7** in 50% yield (Scheme 4).




Scheme 4 Reagents and conditions: DDQ, CH₂Cl₂, rt, 18 h (50%).

As summarized in Scheme 5 we have applied this radical cyclization to several other substrates (**9**, **12**, **15**) and two of these afforded the expected cyclized pyrrolo[3,4-*b*]indoles **10** and **13**. However, only reduction product **14** was obtained from **15**.

Amides **8**,† **11**,† and **14**,† were synthesized by *N*-methylation of the corresponding indole-3-carboxamides³ in 97, 77, and 80% yields, respectively, as illustrated in Scheme 2 for **4**. The usual bromination procedure afforded the 2-bromoindoles **9**,† **12**,† and **15**† in excellent yields. The structures of **9**, **12**, and **15** were confirmed by X-ray crystallography.⁵

Although radical cyclizations to the indole C-2 position are precedented⁷ and the generation of indole C-2 radicals has been described by Jones,^{8,9} our work is the first example of a 1,5-hydrogen atom transfer reaction of a 2-bromoindole-3-carboxamide and subsequent 5-*endo-trig* cyclization to the indole double bond. Following the completion of our initial work, Jones reported⁹ a similar 1,5-H atom abstraction from the radical derived from 2-bromo-3-formyl-*N*-(4-phenylbutyl)indole and subsequent radical translocation and 5-*exo-trig* cyclization to the indole C-2 position. Jones has also recently described¹⁰ a radical translocation sequence leading to indole C-3 5-*exo-trig* cyclization. Some other hydrogen atom abstrac-



Scheme 5 Reagents and conditions: i, BuLi, THF, BrCl₂CCl₂Br, -78 °C to rt (86%); ii, Bu₃SnH, AIBN, toluene, reflux, 48 h (**10**, 51%; **8**, 22%); iii, BuLi, THF, BrCl₂CCl₂Br, -78 °C to rt (90%); iv, Bu₃SnH, AIBN, toluene, reflux, 72 h (**13**, 51%; **11**, 30%); v, BuLi, THF, BrCl₂CCl₂Br, -78 °C to rt (94%); vi, Bu₃SnH, AIBN, toluene, reflux, 72 h.

tion–translocation of α -amidoyl radical schemes not involving indoles have been reported.^{6,11} While we favor the direct 5-*endo-trig* radical cyclization pathway shown in Scheme 1, of which there are precedents,^{12,13} a 4-*exo-trig* cyclization to a spiro β -lactam intermediate followed by a 1,2-alkyl migration (ring expansion), as suggested by a referee, cannot be ruled out. However, such 1,2-alkyl shifts for radicals are rare¹⁴ and the migration terminus is a nucleophilic carbon-centered radical. Moreover, based on the work of Ikeda,¹³ we would have expected to isolate spiro β -lactams if a 4-*exo-trig* radical cyclization pathway was operating.

We thank the National Institutes of Health (GM58601) for support and Drs Mukund Sibi, Jack Li, and Larry Yet for helpful discussion.

Notes and references

† Selected physical and spectroscopic data: **4**: mp 103–104 °C; δ_{H} (300 MHz, CDCl₃) 7.59–7.63 (m, 2H), 7.33–7.36 (m, 1H), 6.98–7.13 (m, 2H), 3.78 (s, 3H), 3.53–3.57 (t, 4H, $J = 5.4$ Hz), 1.38–1.61 (m, 6H); δ_{C} (500 MHz, CDCl₃, -50 °C) 166.3, 135.7, 131.6, 125.3, 121.9, 120.4, 120.2, 110.2, 109.7, 48.8, 43.2, 33.3, 26.6, 25.4, 24.4; IR $\nu(\text{KBr})/\text{cm}^{-1}$ 2934, 2850, 1611; m/z 242 (M⁺), 228, 158 (100%), 131, 103, 77 (Calc. for C₁₅H₁₈N₂O: C, 74.35; H, 7.49; N, 11.56. Found: C, 74.23; H, 7.51; N, 11.50%). **5**: mp 105–107 °C; δ_{H} (300 MHz, CDCl₃) 7.46–7.49 (m, 1H), 7.07–7.26 (m, 3H), 3.73 (br s, 5H), 1.59 (m, 8H); δ_{C} (500 MHz, CDCl₃, -50 °C) 164.5, 135.6, 125.0, 122.1, 120.6, 118.6, 113.9, 110.7, 109.5, 48.2, 42.7, 31.4, 26.6, 25.4, 24.1; IR $\nu(\text{KBr})/\text{cm}^{-1}$ 2934, 2845, 1622, 1522, 1428; m/z 320 (M⁺), 236 (100%), 209, 158, 129; HRMS: calc. m/z 320.0524, found m/z 306.0517. **6**: mp 120–121 °C; δ_{H} (300 MHz, CDCl₃) 7.40 (d, 1H, $J = 7.3$ Hz), 7.13 (t, 1H, $J = 8.0$ Hz), 6.70 (td, 1H, $J' = 1.0$ Hz, $J'' = 7.3$ Hz), 6.42 (d, 1H, $J = 7.7$ Hz), 4.14 (dd, 2H, $J' = 4.5$ Hz, $J'' = 13.2$ Hz), 4.08 (d, 1H, $J = 9.4$ Hz), 3.84 (dd, 1H, $J' = 2.1$ Hz, $J'' = 9.4$ Hz), 2.86 (s, 3H), 2.64 (td, 1H, $J' = 3.4$ Hz, $J'' = 12.8$ Hz), 1.95 (m, 2H), 1.48 (m, 3H); δ_{C} (CDCl₃) 170.6, 151.0, 128.6, 125.1, 124.9, 118.0, 106.1, 69.2, 62.9, 48.9, 40.6, 33.4, 26.7, 24.6, 23.9; IR $\nu(\text{film})/\text{cm}^{-1}$ 2922, 2856, 2344, 1678, 1606, 1489; m/z 242 (M⁺), 158, 131 (100%), 103, 77; HRMS: calc. m/z 242.1420, found m/z 242.1421; (Calc. for C₁₅H₁₈N₂O: C, 74.34; H, 7.49; N, 11.57. Found: C, 74.12; H, 7.41; N, 11.54%). **8**: mp 165–166 °C; δ_{H} (300 MHz, CDCl₃) 7.68–7.71 (dd,

1H, $J' = 1.39$ Hz, $J'' = 7.31$ Hz), 7.46 (s, 1H), 7.19–7.35 (m, 3H), 3.81 (s, 3H), 3.06 (s, 3H), 1.57 (s, 9H); δ_{C} (CDCl₃) 169.4, 136.8, 132.4, 126.1, 122.4, 121.2, 121.0, 114.4, 109.9, 56.4, 35.5, 33.4, 28.1; IR $\nu(\text{KBr})/\text{cm}^{-1}$ 3458, 3113, 2975, 1627 (Calc. for C₁₅H₂₀N₂O: C, 73.74; H, 8.25; N, 11.47. Found: C, 73.67; H, 8.30; N, 11.44%). **9**: mp 105–106 °C; δ_{H} (300 MHz, CDCl₃) 7.57–7.71 (dt, 1H, $J' = 1.2$ Hz, $J'' = 8.0$ Hz), 7.14–7.34 (m, 3H), 3.76 (s, 3H), 2.96 (s, 3H), 1.57 (s, 9H); δ_{C} (CDCl₃) 167.1, 136.4, 125.8, 122.5, 121.0, 119.6, 114.7, 114.2, 109.5, 56.5, 34.2, 31.5, 28.0; IR $\nu(\text{film})/\text{cm}^{-1}$ 3456, 3052, 2979, 1627 (Calc. for C₁₅H₁₉BrN₂O: C, 55.74; H, 5.92; N, 8.67; Br, 24.72. Found: C, 56.13; H, 5.94; N, 8.64; Br, 24.42%). **11**: mp 111–112 °C; δ_{H} (300 MHz, CDCl₃) 7.80–7.83 (dt, 1H, $J' = 1.1$ Hz, $J'' = 7.0$ Hz), 7.13–7.28 (m, 4H), 3.70 (s, 3H), 3.63–3.67 (t, 4H, $J = 5.9$ Hz), 1.57–1.74 (m, 8H); δ_{C} (500 MHz, CDCl₃, -45 °C) 166.9, 135.7, 129.8, 126.2, 121.8, 120.8, 120.1, 110.1, 109.3, 49.0, 45.6, 33.0, 29.3, 28.2, 27.2, 26.0; IR $\nu(\text{film})/\text{cm}^{-1}$ 3053, 2932, 1605 (Calc. for C₁₆H₂₀N₂O: C, 74.97; H, 7.86; N, 10.93. Found: C, 74.90; H, 7.93; N, 10.79%). **12**: mp 125–126 °C; δ_{H} (300 MHz, CDCl₃) 7.46 (d, 1H, $J' = 7.1$ Hz), 7.10–7.27 (m, 4H), 3.73 (s, 3H), 3.40–3.70 (m, 4H), 1.52–1.87 (m, 8H); δ_{C} (CDCl₃) 166.0, 136.3, 125.6, 122.5, 120.8, 119.0, 113.0, 112.9, 109.5, 49.4, 45.9, 31.4, 29.5, 27.9, 27.4, 26.4; IR $\nu(\text{film})/\text{cm}^{-1}$ 3053, 2933, 1618; m/z 334 (M⁺), 255, 236 (100%), 130, 103, 77 (Calc. for C₁₆H₁₉BrN₂O: C, 57.32; H, 5.71; N, 8.36; Br, 23.83. Found: C, 57.57; H, 5.69; N, 8.35; Br, 23.54%). **14**: mp 110–113 °C; δ_{H} (300 MHz, CDCl₃) 8.15 (d, 1H, $J' = 7.5$ Hz), 7.38 (s, 1H), 7.34–7.21 (m, 3H), 3.81 (s, 3H), 3.69 (m, 2H), 1.95 (m, 2H); δ_{C} (500 MHz, CDCl₃, -40 °C) 165.0, 135.9, 130.7, 127.0, 122.3, 121.8, 120.8, 110.2, 109.2, 48.9, 46.3, 33.3, 26.4, 24.2; IR $\nu(\text{film})/\text{cm}^{-1}$ 2942, 2872, 1590; m/z 228 (M⁺), 158 (100%), 130, 103, 77. **15**: mp 100–105 °C; δ_{H} (300 MHz, CDCl₃) 7.54 (d, 1H, $J = 8.1$ Hz), 7.29 (d, 1H, $J = 8.3$ Hz), 7.23 (td, 1H, $J' = 7.6$ Hz, $J'' = 1.0$ Hz), 7.15 (td, 1H, $J' = 7.6$ Hz, $J'' = 1.2$ Hz), 3.76 (s, 3H), 3.74 (t, 2H, $J = 6.6$ Hz), 3.41 (t, 2H, $J = 6.6$ Hz), 2.00 (m, 2H), 1.87 (m, 2H); δ_{C} (CDCl₃) 164.7, 136.2, 125.1, 122.5, 121.0, 120.8, 119.7, 119.4, 109.6, 48.3, 45.8, 31.5, 25.9, 24.6; IR $\nu(\text{KBr})/\text{cm}^{-1}$ 3456, 2922, 2856, 1733, 1622; m/z 308 (M⁺), 262, 236, 192 (100%), 158, 129; HRMS: calc. m/z 306.0368, found m/z 306.0367.

- E. T. Pelkey and G. W. Gribble, *Chem. Commun.*, 1997, 1873; G. W. Gribble, E. T. Pelkey and F. L. Switzer, *Synlett*, 1988, 1061; G. W. Gribble, E. T. Pelkey, W. M. Simon and H. A. Trujillo, *Tetrahedron*, 2000, **56**, 10133.
- G. W. Gribble, *J. Org. Chem.*, 1972, **37**, 1833; G. W. Gribble, J. L. Johnson and M. G. Saulnier, *Heterocycles*, 1981, **16**, 2109.
- H. L. Hassinger, R. M. Soll and G. W. Gribble, *Tetrahedron Lett.*, 1998, **39**, 3095.
- J. Bergman and L. Venemalm, *J. Org. Chem.*, 1992, **57**, 2495.
- J. P. Jasinski, J. C. Badenock, H. L. Fraser and G. W. Gribble, unpublished results.
- V. Snieckus, J.-C. Cuevas, C. P. Sloan, H. Liu and D. P. Curran, *J. Am. Chem. Soc.*, 1990, **112**, 896.
- For recent examples, see G. A. Kraus and H. Kim, *Synth. Commun.*, 1993, **23**, 55; S. Caddick, K. Aboutayab, K. Jenkins and R. I. West, *J. Chem. Soc., Perkin Trans. 1*, 1996, 675; F. E. Ziegler and M. Belema, *J. Org. Chem.*, 1997, **62**, 1083; C. J. Moody and C. L. Norton, *J. Chem. Soc., Perkin Trans. 1*, 1997, 2639; W. Zhang and G. Pugh, *Tetrahedron Lett.*, 1999, **40**, 7591; S.-F. Wang, C.-P. Chuang and W.-H. Lee, *Tetrahedron*, 1999, **55**, 6109; L. D. Miranda, R. Cruz-Almanza, M. Pavón, Y. Romero and J. M. Muchowski, *Tetrahedron Lett.*, 2000, **41**, 10181.
- A. P. Dobbs, K. Jones and K. T. Veal, *Tetrahedron*, 1998, **54**, 2149.
- A. Fiumana and K. Jones, *Tetrahedron Lett.*, 2000, **41**, 4209.
- S. T. Hilton, T. C. T. Ho, G. Pljevaljcic and K. Jones, *Org. Lett.*, 2000, **2**, 2639; S. T. Hilton, T. C. T. Ho, G. Pljevaljcic, M. Schulte and K. Jones, *Chem. Commun.*, 2001, 209.
- A. L. J. Beckwith and J. M. D. Storey, *J. Chem. Soc., Chem. Commun.*, 1995, 977; T. Sato, Y. Kugo, E. Nakaumi, H. Ishibashi and M. Ikeda, *J. Chem. Soc., Perkin Trans. 1*, 1995, 1801; M. Ikeda, Y. Kugo and T. Sato, *J. Chem. Soc., Perkin Trans. 1*, 1996, 1819; J. Rancourt, V. Gorys and E. Jolicoeur, *Tetrahedron Lett.*, 1998, **39**, 5339; J. M. D. Storey, *Tetrahedron Lett.*, 2000, **41**, 8173.
- For leading references to 5-*endo-trig* radical cyclizations, see S. Bogen, M. Gulea, L. Fensterbank and M. Malacria, *J. Org. Chem.*, 1999, **64**, 4920; J. Cassayre, B. Quiclet-Sire, J.-P. Saunier and S. Z. Zard, *Tetrahedron Lett.*, 1998, **39**, 8995.
- H. Ishibashi, N. Nakamura, K. Ikeda, M. Okada, H. Ishibashi and M. Ikeda, *J. Chem. Soc., Perkin Trans. 1*, 1992, 2399; T. Sato, N. Machigashira, H. Ishibashi and M. Ikeda, *Heterocycles*, 1992, **33**, 139.
- B. Giese, *Radicals in Organic Synthesis: Formation of Carbon–Carbon Bonds*, Pergamon, Oxford, 1986, pp. 19–20; J. Fossey, D. Lefort and J. Sorba, *Free Radicals in Organic Chemistry*, Wiley, Chichester, 1995, pp. 161–165.