

# Real-time monitoring of microwave-promoted organometallic ligand-substitution reactions using *in situ* Raman spectroscopy†

Thomas M. Barnard<sup>a</sup> and Nicholas E. Leadbeater<sup>\*b</sup>

Received (in Berkeley, CA, USA) 20th June 2006, Accepted 12th July 2006

First published as an Advance Article on the web 1st August 2006

DOI: 10.1039/b608793k

**An apparatus has been developed for real-time monitoring of organometallic reactions under microwave irradiation using *in situ* Raman spectroscopy and its application for monitoring ligand substitution reactions of Mo(CO)<sub>6</sub> demonstrated.**

The use of microwave irradiation as a tool for synthetic chemistry is an area of increasing interest in both academic and industrial laboratories. It can enhance the rate of reactions and, in many cases, improve product yields.<sup>1</sup> Many of the problems associated with using domestic microwave apparatus, such as reproducibility, accurate temperature measurement and safety, have been overcome with the development of dedicated scientific microwave apparatus. With these it is possible to perform reactions either in sealed tubes or open vessels, monitoring reaction temperature and pressure as well as microwave power input.


A major problem with performing reactions using microwave apparatus is that monitoring its progress generally requires stopping it, allowing the reaction mixture to cool and then using standard analysis techniques such as IR and NMR spectroscopy. Therefore, optimization of reaction conditions such as time and temperature is often a matter of trial and error. This can be a particular problem when trying to prepare organometallic complexes. For example, in ligand exchange reactions, stopping a reaction after a desired number of substitutions is difficult if the reaction cannot be continually monitored. With conventional heating the reaction can be slow and aliquots removed and analyzed over time, but with microwave heating the reaction may be complete within a matter of minutes and accessing a sealed vessel during a reaction is not possible. This may be one reason why there have been very few reports of the application of microwave heating to preparative organometallic chemistry.<sup>2</sup>

There have been some attempts to monitor reactions under microwave irradiation. Neutron and X-ray scattering have been used for studying the preparation of inorganic materials.<sup>3–6</sup> Near IR spectroscopy has been used for the study of a C=O band shift during the reaction between 3-pentanone and triethylorthoformate to form 3,3-diethoxypentane and ethylformate.<sup>7</sup> Pivonka and Empfield have reported the use of Raman spectroscopy as a tool for monitoring organic transformations.<sup>8</sup> They studied an imine

formation reaction and a Knoevenagel condensation. Raman spectroscopy is a particularly useful technique for *in situ* spectroscopy because it relies on light scattering and hence no mechanical interaction with the sample. In addition, glass vessels typically give a low background signal in Raman scattering. Given the extensive use of time-resolved vibrational spectroscopy as a tool for probing reactive organometallic intermediates,<sup>9,10</sup> we wanted to develop an apparatus for the monitoring of organometallic reactions under microwave irradiation using *in situ* Raman spectroscopy.

Construction of the apparatus involved interfacing a scientific monomode microwave apparatus with a commercially available Raman module (excitation source 785 nm; spectral coverage 2400 cm<sup>-1</sup>–200 cm<sup>-1</sup>). This was achieved by drilling a hole (0.8 cm id) in the microwave cavity then attachment of an RF stub to the outer cavity wall (to prevent microwave leakage) and an extender (2.16 cm id) through to the outer casing of the microwave unit. A fiber-optic probe attached to the Raman module was introduced into the microwave cavity, the laser being focused *via* a quartz light tube positioned 5 mm from the reaction vessel. To test the applicability of the apparatus, we decided to study ligand substitution reactions of Mo(CO)<sub>6</sub>. Heating a solution of Mo(CO)<sub>6</sub> in pyridine (py) from room temperature to 180 °C over a period of 90 s using a microwave power of 150 W, we monitored the Raman spectrum in the region 2400–1550 cm<sup>-1</sup>. Scans were taken at 6 s intervals. In pyridine, Mo(CO)<sub>6</sub> shows two bands in the CO region of the Raman spectrum at 2119 cm<sup>-1</sup> (*E<sub>g</sub>*) and 2015 cm<sup>-1</sup> (*A<sub>1g</sub>*) as shown in Fig. 1 (the signal observed at 1580 cm<sup>-1</sup> is from pyridine).

Upon microwave irradiation, within 20 s these bands were seen to decrease in intensity and four new bands grew in (Fig. 2). By comparison with spectra from authentic samples, the bands at 2071 cm<sup>-1</sup> and 1981 cm<sup>-1</sup> were attributed to Mo(CO)<sub>5</sub>(py), that at 1892 cm<sup>-1</sup> to Mo(CO)<sub>4</sub>(py)<sub>2</sub> and that at 1600 cm<sup>-1</sup> to




**Fig. 1** Raman spectrum of Mo(CO)<sub>6</sub> in pyridine (in the region 2400–1500 cm<sup>-1</sup>).


<sup>a</sup>CEM Microwave Technology, 3100 Smith Farm Road, Matthews, NC 28104, USA. E-mail: mike.barnard@cem.com; Fax: +1 704 821-7894; Tel: +1 704 821-7015

<sup>b</sup>Department of Chemistry, University of Connecticut, 55 North Eagleville Road, Storrs, CT 06269-3060, USA. E-mail: nicholas.leadbeater@uconn.edu; Fax: +1 860 486 2981; Tel: +1 860 486 5076

† Electronic supplementary information (ESI) available: Apparatus and experimental details. See DOI: 10.1039/b608793k



**Fig. 2** Raman spectra of  $\text{Mo}(\text{CO})_6$  in pyridine as a function of time upon microwave irradiation (in the region  $2400\text{--}1500\text{ cm}^{-1}$ ).




**Fig. 3** Raman spectra of  $\text{Mo}(\text{CO})_6$  in octylamine as a function of time upon microwave irradiation (in the region  $2400\text{--}1500\text{ cm}^{-1}$ ).

$\text{Mo}(\text{CO})_3(\text{py})_3$ . Holding the reaction mixture at  $180\text{ }^\circ\text{C}$  for a further 1 min resulted in further conversion to  $\text{Mo}(\text{CO})_3(\text{py})_3$ .

The results of a similar study using octylamine (oct) are shown in Fig. 3. Formation of  $\text{Mo}(\text{CO})_5(\text{oct})$  ( $2072\text{ cm}^{-1}$  and  $1976\text{ cm}^{-1}$ ) and some  $\text{Mo}(\text{CO})_4(\text{oct})_2$  ( $1899\text{ cm}^{-1}$ ) was observed after 20 s microwave heating. Heating for 1 min leads to the formation of more  $\text{Mo}(\text{CO})_4(\text{oct})_2$  but prolonging the microwave irradiation to 3 min leads to extensive decomposition.

We also performed the reaction using triphenylphosphine as a ligand. The reaction mixture was heated from  $100\text{ }^\circ\text{C}$  to  $160\text{ }^\circ\text{C}$  over a period of 60 s using a microwave power of 150 W and monitoring the Raman spectrum in the region  $2400\text{--}1500\text{ cm}^{-1}$  (Fig. 4). The bands from the parent hexacarbonyl decreased in intensity and three new bands grew in at  $2073\text{ cm}^{-1}$ ,  $1988\text{ cm}^{-1}$  and  $1943\text{ cm}^{-1}$  (the band at  $1584\text{ cm}^{-1}$  originates from triphenylphosphine). By comparison with literature data,<sup>11</sup> the three bands can be attributed to the formation of  $\text{Mo}(\text{CO})_3(\text{PPh}_3)_3$ . Holding the reaction mixture at  $160\text{ }^\circ\text{C}$  for a further 1 min has little effect.



**Fig. 4** Raman spectra of  $\text{Mo}(\text{CO})_6$  in  $\text{PPh}_3$  as a function of time upon microwave irradiation (in the region  $2400\text{--}1500\text{ cm}^{-1}$ ).

Our results show the potential for using *in situ* Raman spectroscopy as a tool for monitoring the progress of organometallic reactions performed using microwave heating. Work is now underway to broaden the scope of the method for organometallic reactions as well as expand it to organic and metal-mediated catalytic transformations.

Enwave Optonics is thanked for supplying the Raman spectroscopy module. The University of Connecticut is acknowledged for funding.

## Notes and references

- For recent reviews on the general area of microwave-promoted organic synthesis see: (a) C. O. Kappe, *Angew. Chem., Int. Ed.*, 2004, **43**, 6250; (b) M. Larhed, C. Moberg and A. Hallberg, *Acc. Chem. Res.*, 2002, **35**, 717; (c) A. Lew, P. O. Krutzik, M. E. Hart and A. R. Chamberlain, *J. Comb. Chem.*, 2002, **4**, 95.
- For examples see: (a) S. L. VanAtta, B. A. Duclos and D. B. Green, *Organometallics*, 2000, **19**, 2397; (b) M. Ardon, G. Hogarth and D. T. W. Oscrift, *J. Organomet. Chem.*, 2004, **689**, 2429; (c) D. R. Baghurst, D. M. P. Mingos and J. Watson, *J. Organomet. Chem.*, 1989, **368**, C43; (d) D. R. Baghurst, S. R. Cooper, D. L. Greene, D. M. P. Mingos and S. M. Reynolds, *Polyhedron*, 1990, **9**, 893; (e) N. Kuhnert and T. N. Danks, *J. Chem. Res.*, 2002, 66.
- For a review see: G. A. Tompsett, W. C. Conner and K. S. Yngvesson, *ChemPhysChem*, 2006, **7**, 296.
- G. R. Robb, A. Harrison and A. G. Whittaker, *Phys. Chem. Commun.*, 2002, 135.
- W. H. Dokter, T. P. M. Beelen, H. F. Vangarderen, R. A. Vansanten, W. Bras, G. E. Derbyshire and G. R. Mant, *J. Appl. Crystallogr.*, 1994, **27**, 901.
- S. Y. Yang and A. Navrotsky, *Chem. Mater.*, 2004, **16**, 3682.
- S. Hocd , C. Pledel-Boussard, D. Le Coq, G. Fonteneau and J. Lucas, *Proc. SPIE-Int. Soc. Opt. Eng.*, 1999, **3849**, 50.
- D. E. Pivonka and J. R. Empfield, *Appl. Spectrosc.*, 2004, **58**, 41.
- For reviews see: (a) M. W. George and J. J. Turner, *Coord. Chem. Rev.*, 1998, **177**, 201; (b) J. R. Schoonover and G. F. Strouse, *Chem. Rev.*, 1998, **98**, 1335.
- For recent examples see: (a) A. Gabrielsson, F. Hartl, H. Zhang, J. R. Lindsay Smith, M. Towrie, A. Vlcek and R. N. Perutz, *J. Am. Chem. Soc.*, 2006, **128**, 4253; (b) I. V. Rubtsov, Y. K. Kang, N. P. Redmore, R. M. Allen, J. Zheng, D. N. Beratan and M. J. Therien, *J. Am. Chem. Soc.*, 2004, **126**, 5022; (c) M. K. Kuimova, W. Z. Alsindi, J. Dyer, D. C. Grills, O. S. Jina, P. Matousek, A. W. Parker, P. Portius, X. Z. Sun, M. Towrie, C. Wilson, J. X. Yang and M. W. George, *Dalton Trans.*, 2003, 3996.
- T. N. Day, P. J. Hendra, A. J. Rest and A. J. Rowlands, *Spectrochim. Acta, Part A*, 1991, **47A**, 1251.