## Enantioselective nitroaldol (Henry) reaction using copper(II) complexes of (-)-sparteine

H. Maheswaran, K. Leon Prasanth, G. Gopi Krishna, K. Ravikumar, B. Sridhar and M. Lakshmi Kantam *Chem. Commun.*, 2006, 4066–4068 (DOI: 10.1039/b610203d)

The graphic in Table 1 is incorrect. A nitro group is missing from the product. The correct graphic is shown below.

CHO
$$\begin{array}{c} \text{CuX}_2[(\text{-})\text{-sparteine-N,N'})] \\ \text{+} \quad \text{CH}_3\text{NO}_2 \end{array} \begin{array}{c} \text{CuX}_2[(\text{-})\text{-sparteine-N,N'})] \\ \text{Et}_3\text{N} \\ \text{O}_2\text{N} \end{array} \begin{array}{c} \text{OH} \\ \text{NO}_2 \end{array}$$

The Royal Society of Chemistry apologises for this error and any consequent inconvenience to authors and readers.

Additions and corrections can be viewed online by accessing the original article to which they apply.

**RSC** online shop now open

And to celebrate we're having a sale...

From best selling textbooks, to games and puzzles, the RSC online shop brings you class-leading products and services from the RSC. Each sale helps us continue our work in advancing the chemical sciences. Shop online during November and December and get a huge **25% discount\*** on all books, puzzles/games, videos or wall charts purchased.

\* Discount applied to your purchases when you check out. Offer ends December 31st, 2006



RSC Advancing the Chemical Sciences

www.rsc.org/shop