

ALLOPHANATES¹

HERBERT W. BLOHM² AND ERNEST I. BECKER³

Chemical Laboratories of the Polytechnic Institute of Brooklyn, Brooklyn, New York

Received July 25, 1952

CONTENTS

I. Introduction.....	471
II. Formation of allophanates.....	472
A. From cyanic acid.....	472
B. From certain acid chlorides.....	474
C. From carbamates.....	474
D. By ammonolysis or hydrolysis.....	475
E. By miscellaneous methods.....	476
F. Scope and limitations of the methods.....	476
III. Mechanism of the reactions leading to allophanates.....	478
IV. Properties and reactions of allophanates.....	481
V. Uses of allophanates and future problems.....	483
VI. Tables of allophanates.....	484
VII. References.....	499

I. INTRODUCTION

Allophanates are esters of allophanic acid, $\text{H}_2\text{NCONHCOOH}$, an acid which is known only in the form of its derivatives. Allophanates may be prepared from primary, secondary, and tertiary alcohols and from phenols. Although as derivatives they have certain advantages over those usually employed for the identification of alcohols, only a partial listing of the reported allophanates has been made (19, 20) and in only one text on qualitative organic analysis known to the authors are these compounds mentioned (33). Only one brief review of the reaction exists in the English language (20).

It was in 1830 that Liebig and Wöhler (129) first reported the isolation of a new compound resulting from the treatment of methanol with cyanic acid. No physical constants nor analytical data were given by them until sixteen years later (130), when they set out the correct formula and analysis. Having been surprised in isolating neither a cyanide, a cyanate, nor a cyanurate, they named the compound an allophanate.^{4, 5} Although this is the historical develop-


¹ This paper is based upon the B. S. thesis of Herbert W. Blohm.

² Present address: Kay-Fries, Inc., Haverstraw, New York.

³ Inquiries concerning this article should be addressed to Ernest I. Becker at the Polytechnic Institute of Brooklyn.

⁴ Allophane: Greek *allophanes*, "appearing otherwise," from *allos* (otherwise) + *phainesthai* (to appear).

⁵ The term has also been used to describe a class of inorganic aluminum silicates. These are not under consideration here. Although α - and γ -substituted allophanates are also known, the discussion has been limited to esters of the unsubstituted acid:



ment as usually cited, the work of Richardson (166) and of Schlieper (186) must be considered.

In 1837 Richardson published the first ultimate analysis for an allophanate but gave twice the molecular weight for the empirical formula. Schlieper, in 1846, gave the correct ultimate analysis for the product resulting from passing cyanic acid into fusel oil, probably isoamyl allophanate. Neither the melting point nor the correct formula was given, but several chemical reactions were discussed.

Between 1846 and 1919 new members of this class of compounds and new methods for preparing them were reported, but it remained for Béhal (11) to carry out the first extensive preparative study. Later, Locquin and Sung (135-139), Mastagli (141, 142), and Spielman, Barnes, and Close (194) made important contributions.


The mechanism of the formation of allophanates was first investigated by Hofmann (90) and Davis and Blanchard (43). E. A. Werner and A. E. A. Werner (225 and earlier papers) have most closely studied the reaction of cyanic acid with alcohols.

It is the purpose of this paper to review the known methods of preparation and chemistry of the allophanates, to tabulate the known allophanates with their physical constants, and to present some ideas as to the possible course of future work.

II. FORMATION OF ALLOPHANATES

A. From cyanic acid

The most frequently used procedure for the preparation of allophanates is the reaction of an alcohol with cyanic acid. Cyanuric acid is depolymerized to cyanic acid in a stream of carbon dioxide or nitrogen and passed into the alcohol (23, 84, 95, 129, 130, 165, 194, 210, 219) or into a solution of the alcohol in an inert solvent (108, 222). Alternatively, the cyanic acid may be condensed (b.p. 23.5°C.) into an inert solvent and the alcohol added (89, 106, 150, 173, 174, 181, 185, 191, 205, 221).



An apparatus similar to that of figure 1 may be used. The heating wire is conveniently operated from a variable transformer. By using the same length of wire and substituting for B a similar piece of tubing with a T in it for a thermometer, it is possible to determine what setting of the transformer will correspond to 360-400°C. Electrical heating is employed to avoid using a flame near the receiver, which may have in it a volatile solvent, such as ether or benzene.

The alcohol alone, or in a suitable inert solvent, is placed in a test tube or

other receiver at C so that the delivery tube is just below the liquid level. If the delivery tube becomes clogged it may expediently be cleared with a short length of wire. The receiver is cooled in an ice-water mixture. An excess of cyanuric acid is placed in B and a few pieces of dry ice in A, and the apparatus is heated. Cyanic acid is passed into the alcohol or the solution until no further solid separates.

The crude allophanate is filtered, washed with cold ether to remove unreacted alcohol and urethan, and then recrystallized from a suitable solvent. Allophanates have been recrystallized from water, ethanol, methanol, benzene, chloroform, acetone, ethyl ether, and petroleum ether. Water, ethanol, acetone, and benzene have proved adequate in the hands of the authors.

The cyanic acid may also be furnished by treating a cyanate with dry hydrogen chloride in ether (123, 224), by employing a cyanate in acetic acid solution (123), by treating a heavy metal cyanate with hydrogen sulfide (15), or by passing a stream of hydrogen chloride into a suspension of potassium cyanate





FIG. 1. Apparatus for the preparation of an allophanate from cyanic acid


in alcohol (225). The reaction of cyanic acid on a carbamate has also been reported as giving an allophanate in neutral medium (90, 207) and in acid medium (224).



Saytzeff (176, 177) reported that in the reaction of potassium cyanate with ethyl chloroacetate in 90 per cent ethanol solution two allophanates were obtained—ethyl and carbethoxymethyl allophanates.



Ethyl allophanate





Carbethoxymethyl
allophanate

Amato (1, 2) reported a similar experience somewhat later. Another reaction in which potassium cyanate figures as the source of cyanic acid was reported by Wilm (231). Refluxing a solution of the salt with ethyl chloroformate in absolute

ethanol gave potassium carbonate and potassium chloride, ethyl allophanate, ethyl carbonate, and a small quantity of cyanuric acid.



Another source of cyanic acid is the decomposition of urea in the presence of an alcohol (30, 90, 93, 236).




B. From certain acid chlorides

A second general procedure for the preparation of allophanates is by the treatment of an alcohol with an acid chloride. Carbamyl chloride may be used with the alcohol (70, 146, 190, 210, 231, 234)




or allophanyl chloride either in pyridine (46, 47, 158) or without pyridine (23, 82, 83, 150, 152, 154, 158).



In the absence of pyridine the reaction of allophanyl chloride with an unsaturated alcohol was reported to give addition of hydrogen chloride across the double bond (150) and in one instance the tertiary hydroxyl was replaced by chlorine in a dihydroxy compound (158). In the latter case, the reaction proceeded normally in pyridine to give a diallophanate.


Urea may be acylated to produce an allophanate. Thus, ethyl chloroformate (41, 180, 232) and 3-chloropropyl chloroformate (58) give the corresponding allophanates upon reaction with urea.




The reaction appears to give better yields in the presence of an excess of urea, which takes up the acid formed (180).

C. From carbamates

In 1886 Loeb (139) first reported that phosgene would react with ethyl carbamate to give an allophanate.




Later, thionyl chloride (68, 163, 187, 218, 224), sulfuryl chloride (60), and phosphorus pentoxide (14) were found capable of carrying out the conversion.




Phosphorus trichloride and phosphorus oxychloride do not react with urethan in the same way (187).

D. By ammonolysis or hydrolysis

A number of examples exist in which simpler compounds add ammonia or water to give an allophanate, or more complex molecules are ammonolyzed or hydrolyzed to give an allophanate. Thus, carbethoxy isocyanate gives the allophanate with ammonia (55),




carbethoxy cyanamide gives ethyl allophanate with water (145),




and triacylamides are ammonolyzed to the allophanate (52, 54),




while dicarbethoxy cyanamide may be selectively hydrolyzed to give, in one case, an allophanate (53).




Urea appears to be the ammonolytic reagent with ethyl oxalate (86), oxamide and ethyl allophanate being formed (see later).




The hydrolysis or ammonolysis of more complex ureides is shown in the following equations:






E. By miscellaneous methods


The electrolysis of formamide in ethanol is reported to give a "good yield" of the allophanate (179). Benzyl alcohol gives only meager yields, while polyhydroxy alcohols fail to give the allophanates. During an investigation of the rearrangement of potassium glycolobenzoylhydroxamate in boiling isoamyl alcohol, isoamyl allophanate was isolated (91). The reaction of α -diazotoluene with ethyl nitrocarbamate afforded a small quantity of benzyl allophanate (73). Linhard (131) claims that when carbamidosulfonic acid is added directly to ethanol a mixture of the carbamate and the allophanate is formed.



Azides have been shown to give allophanates (38, 39).



Métayer (147) has reported that distillation of a urethan, especially in the presence of Raney nickel, will give an allophanate. Davis and Blanchard (42) and Bishop (16) have reported that while nitrobiuret may be refluxed with absolute ethyl, propyl, butyl, and *tert*-butyl alcohols without reaction and likewise with the addition of 1 per cent of water, the allophanates are formed when more water is added. Most recently, Böttcher and Bauer (22) have reported that the pyrolysis of aryl diurethans can give an allophanate along with some carbamate and cyanuric acid.



Adelson (237, 238) has reported that allophanates may be prepared in improved yields when an alcohol and biuret are heated at reduced pressures.

F. Scope and limitations of the methods

Cyanic acid may be used directly to prepare the allophanates of primary, secondary, and tertiary alcohols, glycols, and phenols. Ethylene glycol and glycerol give monoallophanates (4). Because of the greater ease of hydrolysis of

aryl allophanates, these are most often made through use of carbamyl chloride or allophanyl chloride. For forming derivatives of small quantities of phenol the cyanic acid procedure is satisfactory, although some carbamate may also be present. An interesting synthesis is the preparation of the allophanate of 1,2,2,2-tetrachloroethanol by treating chloral with carbamyl chloride (212).

Carbamyl chloride and allophanyl chloride may be considered as forcing reagents; the yields of allophanates are generally better than with cyanic acid. These chlorides are preferred reagents for the reaction with phenols and thiophenols (128). As implied earlier, a base should be present to take up hydrogen chloride formed in the reaction to prevent further reaction at other sites in the molecule; for example, at a double bond by addition or at a tertiary alcohol by replacement.

Werner (224) claims that the highest yields of allophanate are achieved by means of the action of thionyl chloride on the corresponding carbamate. This method was tested for ethyl, isopropyl, butyl, and isoamyl allophanates. Since this procedure requires the prior preparation of the carbamate, it is difficult to recommend it as advantageous over the use of carbamyl chloride or allophanyl chloride directly on the alcohol.

Some disadvantages in the use of allophanates as derivatives of alcohols should be noted. The yields are often low. With alcohols such as $C_6H_5CH(R)OH$ only about 20 per cent yields have been obtained, using cyanic acid obtained by the pyrolysis of cyanuric acid. In preliminary experiments even lower yields have been obtained with other methods: namely, (a) carbinol and urea, (b) carbinol and potassium cyanate in the presence of acid, (c) carbinol and biuret, (d) chlorocarbonate and urea, (e) carbamate and thionyl chloride, and (f) carbamate and thionyl chloride and urea (34). Among the alcohols which give poor yields of the allophanates are 1-phenylethanol, 1-phenyl-1-propanol, 1-(*sec*-amyl)-1-propanol, allyldimethylmethanol, and cyclopropylphenylmethanol.

Another point is that the melting points are not always as unequivocal as desired. Close has supplied the fact that certain allophanates do not show the expected depression when mixed melting points are determined, as shown by the following example (34):

	Melting point °C.
<i>tert</i> -Amyl allophanate.....	168-169 (gas)
1-Ethylpropyl allophanate.....	175-176 (gas)
Mixture.....	171-172 (gas)


There is reason to believe that decomposition occurs with melting more frequently than has been reported.

The use of cyanic acid permits the formation of derivatives of a wide variety of substituted alcohols. Among the functional groups which do not appear to interfere with allophanate formation are double bonds, triple bonds, conjugated systems, halogens, ethers, quaternary nitrogen atoms, sulfide links, carbonyl groups, ester groups, and carboxyl groups.

One question which has not been answered is the following: "Can cyanic acid ($K_a = 2.2 \times 10^{-4}$ at room temperature (122)) cause the rearrangement of alcohols such as the allyl and propargyl alcohols and the *i*-steroids?"


III. MECHANISM OF THE REACTIONS LEADING TO ALLOPHANATES

The first studies on the mechanism of the reaction of cyanic acid with an alcohol to produce an allophanate were made by Hofmann (90). He stated that when an allophanate is prepared, the urethan is formed when the alcohol is in excess and the allophanate when the cyanic acid is in excess. The evidence offered was that a mixture of amyl allophanate and amyl alcohol in a sealed tube at 160–180°C. gave some amyl urethan, thus marking the urethan as the intermediate between the alcohol and the allophanate.



Eighteen years later, in 1889, Traube (207) reported that when benzyl carbamate was treated with cyanic acid, benzyl allophanate was formed. Further, he claimed that when 1 mole of allophanate was heated with 1 mole of alcohol, 2 moles of urethan were formed. These findings essentially corroborated those of Hofmann.


In 1929 Davis and Blanchard (43) reported on the inability to reproduce the results of Hofmann and proposed a novel course for the reaction.⁶ On the basis of the following evidence they postulated a non-isolable compound as the reactive allophanate-former. In aqueous solution cyanic acid ionizes as a weak acid, trimerizes to cyanuric acid, hydrolyzes to ammonia and carbon dioxide, thus forming ammonium cyanate, and finally dimerizes to dicyanic acid. The last compound can react with aniline to give phenylbiuret or with an alcohol to give an allophanate. It cannot, however, be isolated from aqueous solution, because it undergoes either depolymerization or hydrolysis during evaporation of the solvent. Further evidence cited was that when urea and 1-butanol are refluxed for 3 hr., the allophanate and carbamate are formed and less allophanate is present at the end of 6 hr. Also, 1-butanol and biuret gave a mixture of allophanate and carbamate with less allophanate at longer times. Thus, the dicyanic acid is presumed to form directly and independently of the carbamate as shown below:



In a series of papers A. E. A. Werner and coworkers (207, 225, 228) have refuted the work of Davis and Blanchard and also the work of Hofmann and Traube. Specifically, they were unable to duplicate the synthesis of ethyl allo-


⁶ It is curious that Davis and Blanchard (43) report that Béhal (11) was also unable to reproduce the work of Hofmann, but there is no record in reference 10 of any attempt to check Hofmann's work.

phanate by passing cyanic acid into molten ethyl carbamate or into a neutral solution of urethan in ether. Likewise, this conversion failed when they treated urethan with a mixture of finely ground potassium cyanate and acetic acid. Also, when urea and urethan were heated at 145–170°C. for 3 hr., only unchanged urethan, biuret, and cyanuric acid were recovered, with no trace of allophanate. However, the allophanate was readily prepared when the cyanic acid was produced in the presence of hydrogen chloride. Addition of potassium, sodium, or silver cyanate to a solution of urethan in anhydrous ether saturated with dry hydrogen chloride at room temperature gave the allophanate in about 70 per cent yield. Benzene or dioxane could also be used as the solvent. Werner and Gray (207) claim that the direct addition of cyanic acid to the carbonamide group—CONH₂—is brought about when the cyanic acid is liberated in the presence of hydrogen chloride, because a nascent carbamyl chloride is formed *in situ*.




The nascent carbamyl chloride then converts the carbamate to the allophanate. This intermediate is the one actually used by Gattermann (70) to convert ethanol to ethyl allophanate and also the one which he postulated as giving aromatic amides from aromatic hydrocarbons in the reaction now bearing his name.

Finally, by combining their observations with the work of Raiford and Freyer-muth (164), the Werners wrote the mechanism for the conversion of ethyl carbamate to ethyl allophanate as follows:



An interesting question is why the reaction with cyanic acid stops when two molecules have been added per molecule of alcohol. A. E. A. Werner (226) postulated that in the hydrolysis of diacylureas the formation of a six-membered chelate ring halted the hydrolysis at the removal of one acyl group. This concept may be applied to show that at the formation of a chelated six-membered ring the allophanate is stabilized.




Johnson and Ferstandig (101) claim that they have clearly elucidated the mechanism of this reaction, but their work has not yet appeared in print.

Certain conflicting experiments reported in the literature still remain unanswered. Even facts obtained by Cahours (30), Hofmann (90), Lane (123), and Traube (207) seem to refute parts of the mechanistic picture described by the Werners. Lane (123), for example, claims to have converted 2-methoxy-, 2-ethoxy-, 2-butoxy-, 2-phenoxy-, and 2,2'-butoxyethoxyethanols to the corresponding allophanates by treating the alcohols with suspensions of sodium cyanate in acetic acid, a reaction which the Werners claim does not work.


Birckenbach and Kolb (15) have reported that cyanic acid obtained by treating silver or mercuric cyanate with hydrogen sulfide or with hydrogen chloride in methanol gives only methyl urethan. The cyanic acid obtained by isolation and distillation and then addition to methanol gives urethan and allophanate. Further, the cyanic acid obtained from double decomposition of potassium cyanate, mercuric cyanate mixed with potassium salts, silver cyanate plus alkali metal salts, lead cyanate, and tetramethylammonium cyanate always gives allophanate, with occasionally some urethan. The authors emphasized that a urethan cannot be converted directly to the corresponding allophanate by subsequent addition of cyanic acid. Without further evidence the authors claimed that their results also supported the dicyanic acid postulate of Davis and Blanchard.

One factor which has not been investigated nor mentioned in any of the papers in the field is the purity of the cyanuric acid taken for depolymerization to cyanic acid. It is well known that when zinc chloride is heated with urea it gives cyanuric acid. Since the product is purified by extraction with water, it is doubtful that all of the zinc chloride is removed. During depolymerization of the cyanuric acid small quantities of residual zinc chloride may be volatilized into the stream of carbon dioxide and carried into the reaction mixture. If the reaction between an alcohol and cyanic acid is considered in the modern view as proceeding in two steps, the following equations may be written:⁷

⁷ See references 131-134, inclusive, for evidence that cyanic acid normally is HNCO rather than HOCl.



In the first step a pair of electrons on oxygen attacks the carbon atom of cyanic acid to give the unstable intermediate, which rearranges to the stable carbamate. In the second step, a pair of electrons on the amide nitrogen must attack the carbon of a second molecule of cyanic acid in advance of rearrangement to the allophanate. It is probable that, in the amide, the electrons on nitrogen are not as available for nucleophilic displacement as in an alcohol. This, then, is the slow step. Slight differences in basicity of the alcohols may account for the ease of conversion directly to the allophanate. This may be depicted as follows:



Zinc chloride as well as hydrogen chloride may serve as the Lewis acid A. To establish this hypothesis, it would be necessary to study the effect on the course of the reaction of a number of salts capable of acting as Lewis acids. It would be desirable to study the rate of formation of carbamate and of allophanate as a function of acid strength. It may thus be possible to demonstrate that cyanic acid ($K_a = 2.2 \times 10^{-4}$) at room temperature is capable of catalyzing the conversion of most alcohols to the carbamate but only of certain alcohols to the allophanates.

IV. PROPERTIES AND REACTIONS OF ALLOPHANATES

The chemistry of allophanates has been little studied. Hydrolysis, salt formation, ammonolysis, pyrolysis, reaction with thionyl chloride, acylation, and reaction with halogens have been tried and will be discussed in that order in this section.

The hydrolysis of allophanates may be effected readily by warm alkalies (4, 29, 118, 130, 171, 186). After purification and saponification, the pure alcohol and water-soluble products are obtained—an excellent way in which to purify an

alcohol. Rovira (171) found that potassium benzylate could be used to decompose and analyze the allophanates, since it led to the quantitative evolution of ammonia, which may be absorbed in standard sulfuric acid and titrated. Cold alkalies decompose the allophanates also, but only slowly, so that it is possible to reprecipitate the allophanates from the alkaline solutions if the solutions are neutralized promptly (60, 143). Baeyer (4) reported that with potassium hydroxide, potassium carbonate formed and no potassium allophanate was obtained. Lane (124) has patented the use of allophanates as a means for modifying cellulose fabrics and claims that they are not destroyed by hot alkaline soap solutions. This statement is in contrast to much previous work.

Even in neutral solution allophanates may be hydrolyzed, but this requires an elevated temperature (160°C .) beyond that found in crystallization from the usual solvents (90). Phenolic allophanates, on the other hand, are hydrolyzed by boiling water (24).

Concentrated sulfuric or nitric acid decomposes glyceryl allophanate (a mono-allophanate) with the evolution of carbon dioxide, a reaction which was interpreted as hydrolysis (4). However, cold dilute acids did not affect this allophanate.

Salts of allophanates have been prepared by the reaction of bases with ethyl allophanate (18, 60, 130, 176, 208). In this way, the barium (130, 176), calcium (130), potassium (130, 143), and sodium (60, 130) salts have been prepared. The sodium salt has also been prepared by double decomposition of the barium salt with sodium sulfate (130). The potassium salt has been obtained by adding ethyl allophanate to a solution of potassium amide in liquid ammonia (18). The disilver salt has been precipitated by treating ethyl allophanate with silver nitrate in hot water (40). It is not entirely surprising that treatment of the barium salt with neutral silver oxide or lead acetate failed to precipitate the new metal salt (130). It is interesting to note that a disodium salt of allophanic acid has been isolated by treating ethyl carbamate with sodium in benzene (60).

The acidity of the allophanates appears to arise from the diacyl nitrogen structure, ROCONHCONH_2 , which is well known to be responsible for the acidity of barbiturates and other classes of compounds.


Apparently acids can also form salts with allophanates. Mauquin (143) has reported that treatment of ethyl allophanate with nitric acid in ether gave beautiful crystals of the mononitrate.

Ammonolyses of allophanates have been effected with ammonia (17, 90, 207), aromatic amines (41, 90), aliphatic amines (90), and phenylhydrazine (41). Blair (17) reported that at 300°C . a mixture of ammonia, ammonium chloride, and ethyl allophanate produced a mixture of guanidine, urea, and ammonium carbamate, while at lower temperatures ammonia alone with ethyl or methyl allophanate gave biuret and ethanol. Hofmann (90) and Traube (207) also reported that heating with ammonia gave biuret. However, Schlieper (186) reports that ammonia does not react in warm solution.

Aniline at $120\text{--}125^{\circ}\text{C}$. has been shown to convert ethyl allophanate to diphenylbiuret, while at $160\text{--}170^{\circ}\text{C}$. it gives a mixture of monophenylurea and

diphenylurea (41). With *o*-anisidine, *o*-toluidine, and *m*-nitroaniline at about 125°C. the products were aryl-substituted allophanates, ArNHCONHCOOC₂H₅. As the temperature was raised, substituted biurets and arylureas were formed (41). With aniline, *m*-toluidine, and *p*-bromoaniline, none of the substituted allophanates were obtained; only substituted ureas and biurets were isolated. No explanation for this change in the course of the reaction was offered.

In 1920 Dains and Wertheim (41) showed that when phenylhydrazine reacts with methyl allophanate for a long time at 150°C., 1-phenylurazole is the product.



Pyrolysis of an allophanate yields the alcohol and cyanuric acid (78, 228). When ethyl allophanate is refluxed with thionyl chloride at atmospheric pressure it is unchanged, but when heated in a sealed tube with xylene at 140–150°C., 5 g. of ethyl allophanate is converted to 2 g. of cyanuric acid (187).

Waltmann and Wolf (217) have reported that stearyl allophanate may be treated with trioxymethylene and dry hydrogen chloride to give a chloromethyl derivative which may then be treated with pyridine at room temperature to give the quaternary salt.



Mauquin (143) has successfully acylated ethyl allophanate with the acid chloride of monoethyl oxalate.

Schlieper (186) has found that a warm aqueous suspension of amyl allophanate does not react with ammonia, nitric acid, chlorine, or sulfuric acid.

V. USES OF ALLOPHANATES AND FUTURE PROBLEMS

In addition to their application for the identification of alcohols, the allophanates may be used for their purification. As has been stated earlier, allophanates are readily hydrolyzed in acidic or basic solution.

In the patent literature claims have been made for the use of allophanates as intermediates in the preparation of water-repellent fabrics (124, 217), as softeners for hard deposits of carbon on the pistons of engines (61, 239), as additives to lubricants for use at extreme pressures (237, 238), for the conversion of medicinal agents of unpleasant odor or taste to odorless and tasteless compounds (153, 211, 212), and as depilatories (51).

Recently the investigation of allophanates as anticonvulsant drugs was reported (194). Benzyl allophanate, α -ethylbenzyl allophanate, and a few branched-chain aliphatic allophanates were found to be effective against electroshock,

metrazol, or both, but none was deemed worthy of clinical investigation. They were also found to have sleep-producing action.

In respect to the preparation of allophanates, the toxicity of cyanic acid should be mentioned; it is a corrosive liquid at 0°C. Sodium cyanate has a hypnotic action (188).

The discussion of the mechanism of the formation of allophanates makes it clear that additional work needs to be done to complete the picture. However, certain other studies are also obviously lacking.

A quantitative knowledge of the acidity of allophanates is not available. Such data would be of interest in studies of salt formation and in a study of the mechanism of the formation of allophanates as well. Although the allophanates presumably have one acidic hydrogen, only one reaction for their acylation has been carried out (143). The site of acylation was claimed to be the β -position, whereas it would appear that the α -position should carry the more active nitrogen. No alkylation reactions with reagents such as diazomethane⁸ and dimethyl sulfate have been reported. The reactions with nitrous acid or with coupling reagents, which conceivably could give interesting if not also useful compounds, have not been studied.

VI. TABLES OF ALLOPHANATES

Table 1 contains data on the allophanates of alcohols which are liquid at 25°C., while table 2 presents data on the allophanates of alcohols which are solid at this temperature. In each table the alcohols are arranged in the order of increasing boiling point. Table 3 contains data on the allophanates of a number of alcohols for which no physical constants are given in the literature. Tables 4 and 5 present data on the allophanates of phenols. Allophanates of a few miscellaneous alcohols are to be found in table 6.

⁸ Close (34) has stated that ethyl, isopropyl, and 1-phenylpropyl allophanates fail to react with diazomethane even in the presence of methanol.

TABLE 1
Allophanates of alcohols which are liquid at 25°C.

ALCOHOL	BOILING POINT °C.	MELTING POINT OF ALLOPHANATE °C.	REFERENCES
Methanol.....	65	208 210 212	(70) (224) (11)
1,1,1-Trifluoro-2-propanol.....	77.7-77.9	159.7 (dec.)	(200)
Ethanol.....	78	188 190 191 193 194 195 197-198	(218) (11, 39) (38) (225) (140) (23) (237)
<i>tert</i> -Butyl alcohol.....	83	190	(11, 194, 237)
2-Propanol.....	88	179 180 184-185	(224) (11) (194)
3-Buten-2-ol.....	96-97	151-152	(44)
Allyl alcohol.....	97	165	(11, 237)
1-Propanol.....	98	167 169.6-169.8 175 175.5	(7) (92) (224) (11)
2-Butanol.....	100	159.5	(11, 194)
2-Methyl-2-butanol.....	102	151-152 ^(a) 166-168	(11, 165) (194)
Cyclopropanol.....	100-103 53-55/100 mm.	179-181 (subliming with decom- position)	(194)
Isobutyl alcohol.....	107-108	174 178-179 179-180 180.5	(93) (165) (194) (11)
3-Methyl-2-butanol.....	111.5-112	163 ^(a) 169-170	(21) (194)
3-Penten-2-ol.....	112	158-159	(49)
3-Methyl-3-buten-2-ol.....	112-113	150-151	(126)
1-Penten-3-ol.....	114-116	151-152 152-153 155	(45) (44) (78)
3-Pentanol.....	116	171-172 179.5	(194) (11)
1-Butanol.....	117	149, 149- 150 149.5-150.5	(224, 164) (11)
2-Pentanol.....	119	154 157-158	(11) (194)

TABLE 1—Continued

ALCOHOL	BOILING POINT	MELTING POINT OF ALLOPHANATE	REFERENCES
	°C.	°C.	
2-Methyl-2-pentanol.....	123	125.6-126.6	(125)
		128	(11)
3-Methyl-3-pentanol.....	123	152	(78)
		153-154	(194)
2-Methoxyethanol.....	125	163	(123)
2-Methyl-3-pentanol.....	128	179 ^(b)	(11)
		185-186	(194)
2-Methyl-1-butanol.....	128	149-150	(194)
2-Chloro-1-ethanol.....	129	179-182	(194)
		182.5	(78)
3-Methyl-1-butanol.....	131.5	149.4-150	(92)
		150	(11)
		160	(224)
		162	(7)
4-Methyl-2-pentanol.....	132	161 ^(b)	(11)
		162-163	(194)
2,4-Dimethyl-2-pentanol.....	132-133	132	(11)
3,4-Dimethyl-1-pentyn-3-ol.....	133	144	(127)
1-Hexen-4-ol.....	76-77.8/95 mm.	169	(67)
1-Hexen-3-ol.....	133.5-134	139.5-140	(44)
2-Hexen-1-ol.....	101.4-101.8/95 mm.	154.5-155.5	(66)
4-Ethoxyethanol.....	135	143-145	(123)
3-Hexanol.....	135	185.5	(78)
1-Methylcyclopentanol.....	135-136	157	(32)
3-Ethyl-1-pentyn-3-ol.....	135-137	130-131	(138)
3-Methyl-1-hexyn-3-ol.....	137	133	(139)
2-Hexanol.....	137-138	170-171	(194)
		173 ^(b)	(11)
1-Pentanol.....	138	158-159	(194)
		159.3-159.8	(92)
		162	(90, 187)
2,4-Dimethyl-3-pentanol.....	138	173.5-174.5	(21)
3-Methyl-5-hexen-3-ol.....	139	132-135	(194)
4-Penten-1-ol.....	138.5-139.5	147-148 (dec.)	(159, 160)
Cyclopentanol.....	139-140	179.5	(11)
3-Methyl-3-hexanol.....	140	148	(78)
		149	(235)
2-Penten-1-ol.....	141-142	157-157.5	(45)
3-Ethyl-3-pentanol.....	142	156	(78)
		173-174	(194)
		182-183	(144)
3,4,4-Trimethyl-1-pentyn-3-ol.....	142-144	156	(135)
Methylpseudobutylvinylcarbinol.....	146-147	167-168	(136)
2-Ethyl-1-butanol.....	147.7	156-158	(194)
		161-161.5	(21)

TABLE 1—Continued

ALCOHOL	BOILING POINT °C.	MELTING POINT OF ALLOPHANATE	REFERENCES
		°C.	
2-Methyl-1-pentanol.....	147.7	150-151 155.8- 156.6 ^(c)	(194) (125)
5-Methyl-3-hexanol.....	147-148	185-187	(194)
3,5-Dimethyl-1-hexyn-3-ol.....	148	114	(127)
cis-o-Methylcyclopentanol.....	148-149	174	(77)
4-Methyl-3-hexanol.....	149-150	172-174	(194)
Ethyl α -hydroxyisobutyrate.....	150	125-126 (dec.)	(194)
trans-o-Methylcyclopentanol.....	150-151	174	(77)
Trichloroethanol.....	151	182-183	(230)
4-Methyl-1-pentanol.....	151-152	155-156 162	(185) (11)
2,2,3-Trimethyl-3-pentanol.....	152	134-135	(135)
3-Heptanol.....	156	185.4- 185.6 187	(125) (78)
1-Hepten-3-ol.....	153.5-154	156.5-157	(44)
4-Heptanol.....	154-155	206	(11)
2-Ethyl-2-methyl-1-butanol.....	155	102	(64)
Ethyl lactate.....	155	170	(207)
1-Hexanol.....	156	159-160 163.3-163.8 165	(147) (125) (11)
3-Hexen-1-ol.....	156-157	143-144 (synthetic) 134-135 (natural) 143 (cis) 146 (trans) 146 (natural)	(174, 204) (174) (203) (203) (204)
2,4-Dimethylpentanol.....	157	116-117	(194)
4-Hexen-1-ol.....	155-158	161.5	(167, 168)
2-Heptanol.....	158-160	148-150	(19, 194)
Ethyl glycolate.....	160	144	(207)
3-Ethyl-3-hexanol.....	161	149	(11)
3-Chloro-1-propanol.....	160-162	166	(58)
1,1,1-Trichloro-2-propanol.....	161	186	(212)
Cyclohexanol.....	161	173 179	(194) (11)
2-Diethylaminoethanol.....	162	137-138	(146)
3-Methyl-3-heptanol.....	164	130	(78)
2-Methylcyclohexanol.....	166.5	177	(78)
Furfuryl alcohol.....	170	167.5 ^(a)	(78)
2-Butoxyethanol.....	171	118	(123)
3,5-Dimethyl-4-heptanol.....	171	163	(209)

TABLE 1—Continued

ALCOHOL	BOILING POINT °C.	MELTING POINT OF ALLOPHANATE °C.	REFERENCES
<i>cis</i> -3-Methylcyclohexanol.....	171–171.4	141.6	(121)
3-Ethyl-5-methyl-3-hexanol.....	172	145	(11)
3-Methyl-1-hexanol.....	172	137	(235)
<i>trans</i> -3-Methylcyclohexanol.....	173.6–174.6	178.8	(121)
<i>cis</i> -1,4-Dimethyl-2-cyclohexanol.....	175	158 ^(d)	(74, 75)
<i>cis</i> -1,3-Dimethylcyclohexan- <i>trans</i> -4-ol.....	175–176	207 ^(d)	(76)
<i>cis</i> -1,3-Dimethylcyclohexan- <i>cis</i> -4-ol.....	176	184 ^(d)	(76)
1-Heptanol.....	176	160	(11)
1,3-Dimethyl-4-cyclohexanol.....	176–177	149 ^(d)	(76)
<i>trans</i> -1,4-Dimethyl-2-cyclohexanol.....	177	125 ^(d)	(74, 75)
2-Methyl-2-hepten-6-ol.....	177–178	99–100	(56)
1,3-Dichloro-2-propanol.....	178	182	(190)
Tetrahydrofurfuryl alcohol.....	178	161– 161.5 ^(e)	(92)
2,6-Dimethyl-4-heptanol.....	178	156 ^(b)	(11)
2-Octanol.....	179	155	(11)
3-Octanol.....	178.5–179.5	167	(12)
<i>d</i> -3-Octanol.....	88–92/33 mm.	182	(29)
2-Hydroxyethyl formate.....	179–180	151	(83)
3-Methyl-4-octanol.....	180	160	(209)
4-Ethyl-4-heptanol.....	180	124	(135)
2-Ethyl-1-hexanol.....	181–183	124 125	(147) (142)
Cycloheptanol.....	185	184	(173)
3,7-Dimethyl-1-octyn-3-ol.....	187–188	114–115	(135)
2-Hydroxyethyl acetate.....	187–189	164	(83)
3,5,5-Trimethyl-1-hexanol.....	191	148.5–149	(21)
5-Nonanol.....	194	158	(209)
2,2'-Ethoxyethoxyethanol.....	195	102–104	(123)
1-Octanol.....	194–196	155–156 157	(70) (11)
Ethylene glycol.....	197	153.8–154.8 160	(92) (4, 237)
1-Phenylethanol.....	204	181–182 181.5 185–186	(193) (11) (194)
Benzyl alcohol.....	206	121 182–183 183 191.5 191–191.5	(11) (194) (207) (23) (152)
1-Nonanol.....	211	158	(11)
<i>d</i> -Neomenthol.....	211.5–211.8	215.5	(236)
α -Allyl- α -methylbenzyl alcohol.....	217–221	139–143 (dec.)	(194)
1-Phenyl-1-propanol.....	217–221	149–150	(194)
Phenethyl alcohol.....	219–221	186 197.5	(11) (23)

TABLE 1—Continued




ALCOHOL	BOILING POINT °C.	MELTING POINT OF ALLOPHANATE °C.	REFERENCES
Carvomenthol.....	222	192.5	(11)
3-Phenyl-3-pentanol.....	223-224	151-152	(194)
Nerol.....	224-225	84-84.5	(152)
Undecanol.....	225.4	155.5-158	(11)
Geraniol.....	229	120-121	(25, 184)
		124-124.5	(149, 152, 237)
2,2'-Butoxyethoxyethanol.....	230.4	88	(123)
1-Decanol.....	233	159	(11)
Phenylpropanol.....	236	165	(11)
2-Phenoxyethanol.....	237	123	(82)
		219-220	(123)
Cinnamyl alcohol.....	258	185	(11, 237)
1-Dodecanol.....	259	159	(97)
		159.5	(11)
Anisyl alcohol.....	259	180.25	(11)
Diethyl tartrate.....	280	188	(207)
Glycerol.....	290	About 160	(4)
2,3-Dimethyl-1-penten-3-ol.....	44-45/18 mm.	147	(36)
1,10-Dimethyl-5,10-cis-7-naphthiten-1-ol	77-78/0.3 mm.	178.8-179.6 (dec.)	(69)
3,6,7-Trimethyl-2,6-octadien-8-ol.....	61-62/0.009 mm.	108-109	(182)
2-(2',6',6'-Trimethyltetrahydro-2'-pyranyl)ethanol.....	62-63/0.2 mm.	165-165.5	(31)
2-Hydroxymethyl-5,5-dimethylcyclohexanone.....	66/0.24 mm.	154	(197)
	95-96/0.05 mm.	162-163	(35)
	98-100/0.05 mm.	183-185 (not con- stant)	(35)
	104-105/0.05 mm.	167	(31)

TABLE 1—Continued


ALCOHOL	BOILING POINT	MELTING POINT OF ALLOPHANATE	REFERENCES
	°C.	°C.	
Phytol.....	155–157/1 mm.	74 ^(f)	(116)
		77–78	(206)
Dihydrophytol.....	157–159/0.8 mm.	73	(104)
3-Hydroxymethyl-2,6,10-trimethylhendeca-1,5,9-triene.....	95.5–96.5/0.01 mm.	80	(35)
2-Acetyl-5,9-dimethyldeca-4,8-dien-1-ol.....	104–106/0.01 mm. 122/0.15 mm.	99–100 161–162 ^(g)	(35) (181)
3-Hydroxymethyl-2,6,10-trimethylhendeca-5,9-dien-2-ol.....	127–130/0.01 mm.	161–162 ^(g)	(35)
			
An alcohol isomeric with the above ^(h)	175–178/0.45 mm.	165	(199)
2-Methyl-3-(4'-isopropylcyclohexyl)propan-1-ol.....	116–117/1.7 mm.	182	(199)
3,7-Dimethyl-1,7-octanediol.....	124–125/1 mm.	120.5–121 196–197 (mono-allophamate) ⁽ⁱ⁾ (diallophamate)	(151) (158)
3(4?)-Bromo-3-hexen-1-ol.....	68–69/3 mm.	113 171	(203)
Cyclocitronellol.....	84/4 mm.	172 132 (stereo-isomer)	(5) (5)
cis-Cryptol.....	86/6 mm.	166–168	(72)
l- α -Terpineol.....	78–81/2.6 mm.	133–134	(152)
trans-Cryptol.....	90/4 mm.	196–197 (dec.)	(72)
2-Methyl-3-(4'-isopropylphenyl)propen-1-ol..	130–130.5/4 mm.	171–172	(151)
3-Methyl-1-heptyn-3-ol.....	63/12 mm.	126–127	(127)
3-Hexyn-1-ol.....	66–67/13 mm.	187	(203)
2-Iodoethanol.....	85/25 mm.	182	(219)
1-Ethynylcyclohexanol.....	73–75/15 mm.	197	(138)
3-Propyl-1-hexyn-3-ol.....	69–71/12 mm.	143	(135)
3-Propyl-1-hexanol.....	78–79/16 mm.	133–134	(137)
4-Hepten-1-ol.....	81.5/17 mm.	168.8	(168)
4-Hepten-1-ol.....	81.5/17 mm.	168.5	(167)
Ethyl 2-hydroxy-3-methyl-3-butenoate.....	71–76/12 mm.	121	(215)
3-Propyl-1-hexen-3-ol.....	75–76/12 mm.	112	(136)

TABLE 1—Continued

ALCOHOL	BOILING POINT °C.	MELTING POINT OF ALLOPHANATE °C.	REFERENCES
2,2,3-Trimethyl-6-hepten-3-ol	74/10 mm.	112	(37)
3-Methyl-1-octyn-3-ol	75/10 mm.	120	(127)
3,3-Dimethyl-1-ethynylcyclohexanol	81/11 mm.	157	(214)
2-Ethyl-2-hydroxymethyldioxolane	82/11 mm.	147	(229)
3,4,4-Trimethyl-2-penten-1-ol	84/12 mm.	77	(137)
2-(2'-Methyl-2'-dioxolanyl)ethanol	86/11 mm.	149	(229)
2,7-Dimethyl-1-octen-3-ol	85-87/11 mm.	87-88	(185)
d-2-Octanol	87/17 mm.	182-182.5	(148)
4-Octen-1-ol	95/17 mm.	160	(167, 168)
5-Methoxy-4-penten-1-ol	95.5/16 mm.	151	(170)
l-Tetrahydrolinaloöl	86-88/12 mm.	88	(6)
1-Hydroxymethyl-3-methylcyclohexene	88/11 mm.	172-173	(240)
1-Hydroxymethyl-3, 3-dimethylcyclohexene	88-92/11 mm.	163-164	(240)
2-Isopropenyl-2, 3-dimethyl-4-penten-1-ol	89/12 mm.	179-180	(25)
2-Isopropyl-3, 3-dimethyl-1-pentanol	89/11 mm.	163	(25)
4,7-Octadien-1-ol	90/12 mm.	153	(169)
3,7-Dimethyl-3-octanol	92-93/12 mm.	110-111	(6, 135)
(d, l-Tetrahydrolinaloöl)		108-109	(120)
5-Ethoxy-4-penten-1-ol	110/26 mm.	145	(170)
1-Cyclohexyl-2-propen-1-ol	93-94/13 mm.	169-170	(48)
2-Isopropenyl-4-methyl-4-hexen-1-ol	93/12 mm.	123	(126)
2-Isopropyl-4-methyl-1-hexanol	97-98/14 mm.	125	(126)
4-Bromo-4-penten-1-ol	97/15 mm.	138-139	(160)
Cyclooctanol	99/16 mm.	183.5	(173)
2-Methylene-4, 4-dimethyl-1-hydroxymethyl-cyclohexane	99-100/15 mm.	163-164	(197)
d, l- γ , δ -Dihydrolavandulol	94/12 mm.	121	(183)
(5-Methyl-2-isopropenyl-1-hexanol)			
5-Methyl-2-isopropyl-3-hexen-1-ol	92-95/12 mm.	143-144	(191)
d, l- β , γ -Dihydrolavandulol	95/12 mm.	121-122	(191)
(5-Methyl-2-isopropyl-4-hexen-1-ol)			
Isolavandulol	95/11 mm.	120	(184)
d, l-Lavandulol	100-101/16 mm.	114-115	(26)
		119	(183)
		119-120	(184)
d, l-Tetrahydrolavandulol	96-97/12 mm.	99-100	(183, 191)
		100-101	(184)
1-Hydroxymethyl-2, 2, 4-trimethylcyclohexane	95-96/10 mm.	166-167	(196)
4-Nonen-1-ol	111-113/22 mm.	149.5	(161)
2-Hydroxyethyl butyrate	98-100/14 mm.	142	(83)
CH ₃ CH ₃	CH ₂ C(CH ₃) ₂ OH	106-108/16 mm.	117
			(213)
trans-2-Isopropyl-5-methyl-3, 5-hexadien-1-ol	94-97/11 mm. (lower-boiling fraction)	114-116	(191)

TABLE 1—Continued

ALCOHOL	BOILING POINT	MELTING POINT OF ALLOPHANATE	REFERENCES
	°C.	°C.	
cis-2-Isopropyl-5-methyl-3,5-hexadien-1-ol...	94-97/11 mm. (higher-boiling fraction)	78-80	(191)
2,3,6-Trimethylocta-6-en-1-yn-3-ol.....	96-97/11 mm.	138	(182)
3,5,5-Trimethylhepta-2,6-dien-1-ol.....	98-101/12 mm.	137-138	(26)
3-Propyl-2-hexen-1-ol.....	99-101/12 mm.	147-148	(137)
(+)-Cyclolavandulol.....	96-99/11 mm.	157-158	(24)
2,6-Nonadien-1-ol.....	95.5-100/11 mm.	140 145, 148	(201, 202) (174)
d, l-Cyclolavandulol.....	98-100/11 mm.	160 163-164	(24) (24)
2-(3',3'-Dimethylcyclohexylidene)ethanol...	102-103/13 mm.	158-159	(214)
1,5,5-Trimethyl-4-hydroxymethylcyclo- hexene.....	110/15 mm.	157-158	(196)
2,7-Dimethyl-2-octen-1-ol.....	103-104/11 mm.	124-125	(185)
5,5-Dimethyl-1-cyclohexeneethanol.....	102-103/12 mm.	183-184	(214)
2,7-Dimethyl-2,6-octadien-1-ol.....	106-108/12 mm.	122-123	(185)
1,3,3-Trimethyl-4-hydroxymethylcyclo- hexene.....	105-106/11 mm.	146-147	(196)
Citronellol.....	107-108/12 mm.	108-108.5 112-113	(150) (80)
3-(2'-Methyl-2'-dioxolanyl)-1-propanol.....	108/11 mm.	119	(229)
Tetrahydrogeraniol.....	98-100/7 mm.	117 117-118	(137) (25, 120)
4-Nonen-1-ol.....	118-119/23 mm.	149.5	(168)
2-Phenylpropan-1-ol.....	116/18 mm.	175	(141, 142)
d,l-β-Citronellol.....	118-119/18 mm.	104.5-105.5	(56)
Cyclononanol.....	112-113/13 mm.	177	(173)
4-Butyl-3-oxo-1-butanol.....	112-113/13 mm.	135	(229)
7-Hydroxycineole.....	108-111/9 mm.	169	(46)
l-Citronellol.....	108-109/10 mm.	106-107	(57)
d-Citronellol.....		105-106	(57)
2-(4'-Methylphenyl)propan-1-ol.....	124/17 mm.	157	(141, 142)
5-Butoxy-4-penten-1-ol.....	124-127/17 mm.	144	(170)
Myrcenol.....	123-128/17 mm.	110-111	(47)
Lactic acid.....	122/14 mm.	190	(207)
Cuminalcohol.....	122.5/13 mm.	201	(142)
2-Acyl-5-methyl-4-hexen-1-ol.....	122-123/11 mm.	146	(188)
2-(2',4'-Dimethylphenyl)ethanol.....	125/10 mm.	222	(85)
2-Acetyl-4-methyl-4-hexen-1-ol.....	123-124/13 mm.	151	(126)
α-Ionol.....	126.6-127/14.5 mm.	137-138 ⁽¹⁾	(175)
β-Ionol.....	130.5/14.5 mm.	144-145	(175)
2-Butyl-1-octanol.....	132/15 mm.	119	(142)
2-Ethyl-3-phenyl-2-propen-1-ol.....	134/15 mm.	147	(141, 142)
Tetrahydroionol.....	134.5/15 mm.	164	(156)
2-Ethyl-3-phenyl-1-propanol.....	134-135/15 mm.	135	(142)

TABLE 1—Continued

ALCOHOL	BOILING POINT	MELTING POINT OF ALLOPHANATE	REFERENCES
			°C.
2-Methyl-2-undecanol.....	134–135/15 mm.	113.5	(11)
2-Benzoyloxyethanol.....	137–138/17 mm.	156	(82, 155)
2-(2',2',3'-Trimethyl-1-cycloheptylidene)-ethanol.....	131–132/10 mm.	170	(178)
2-(2',2',3'-Trimethyl-1-cycloheptyl)ethanol.....	131–132/10 mm.	168	(178)
2- <i>o</i> -Anisylethanol.....	136/12 mm.	176–177	(84)
2-Methyl-2-(1'-hydroxy-6'-methyl-5-hepten-2'-yl)-1,3-dioxolane.....	136/11 mm.	102–103	(183)
2-Phenethyloxyethanol.....	140–142/15 mm.	150	(82, 155)
Irol.....	136–140/13 mm.	197	(175)
2-Hydroxyethyl pelargonate.....	143–144/14 mm.	142	(83)
1-Benzoyloxy-2-butanol.....	151–152/18 mm.	102	(155)
3-Benzoyloxy-1-propanol.....	155–157/20 mm.	119	(155)
2-(3'-Phenylpropoxy)ethanol.....	154–155/18 mm.	131	(82)
		131.5	(81, 155)
2-Butyl-3-phenyl-2-propen-1-ol.....	155/15 mm.	155	(141, 142)
2-Butyl-3-phenyl-1-propanol.....	155/15 mm.	144	(142)
3-(3'-Phenylpropoxy)-1-propanol.....	160/20 mm.	113	(155)
2-Amyl-2-nonen-1-ol.....	154/13 mm.	111.5	(142, 157)
2-Amyl-1-nonanol.....	154/13 mm.	120	(142, 157)
5-Phenyl-4-penten-1-ol.....	157/16 mm.	170.5	(167, 168)
2-Amyl-3-cyclohexyl-1-propanol.....	158/15 mm.	108	(142)
2-Amyl-3-phenyl-1-propanol.....	162/13 mm.	125	(142, 157)
2-Amyl-3-phenyl-2-propen-1-ol.....	162/12 mm.	160	(141, 142)
2-Hydroxyethyl phenylacetate.....	165–166/15 mm.	157–158	(83)
α-Santalol.....	166–167/14 mm.	162–163	(28)
4-(3'-Phenylpropoxy)-2-butanol.....	175/19 mm.	165	(155)
Lanceol.....	175–176/17 mm.	114–115	(27)
Santalol.....	170–178/17 mm.	162 ^(k)	(210)
β-Santalol.....	177–178/17 mm.	159–160	(28)
2-Hexyl-3-phenyl-1-propanol.....	175–176/15 mm.	124	(142)
2-Hexyl-3-phenyl-2-propen-1-ol.....	176/15 mm.	142	(141, 142)
2-Hexyl-1-decanol.....	177/15 mm.	90	(142)
2-Heptyl-3-phenyl-2-propen-1-ol.....	186/15 mm.	139	(142)
2-Heptyl-3-phenyl-1-propanol.....	187/15 mm.	128	(142)
1,3-Diphenyl-1-propanol.....	194/15 mm.	99	(142)
2-Heptyl-1-undecanol.....	198/15 mm.	80	(142)
2-Octyl-3-phenyl-2-propen-1-ol.....	198/15 mm.	138	(141, 142)
2,3-Diphenyl-1-propanol.....	190–210/15 mm.	162	(142)
2-Octyl-3-phenyl-1-propanol.....	200–201/15 mm.	117	(142)
2,2-Dibenzylethanol.....	202/15 mm.	140	(142)
1,3-Diphenyl-1-butanol.....	202/15 mm.	148	(142)
1-(4'-Methylphenyl)-3-phenyl-1-propanol.....	200–202/13 mm.	111	(142)
2-Nonyl-3-phenyl-1-propanol.....	207–208/15 mm.	97	(142)
2-(8'-Nonen-1'-yl)-3-phenyl-2-propen-1-ol....	212/17 mm.	127	(141, 142)
2-Nonyl-3-phenyl-2-propen-1-ol.....	212/17 mm.	132	(141)
		136	(142)

TABLE 1—Concluded

ALCOHOL	BOILING POINT °C.	MELTING POINT OF ALLOPHANATE	REFERENCES
		°C.	
2-(8'-Nonen-1'-yl)-3-phenyl-1-propanol.....	211/15 mm.	109	(142)
2-Octyl-1-dodecanol.....	215/15 mm.	69	(142)
2-Decyl-3-phenyl-2-propen-1-ol.....	221/15 mm.	137	(141, 142)
2-Decyl-3-phenyl-1-propanol.....	221-222/15 mm.	109	(142)
2-Benzyl-5-phenyl-1-pentanol.....	224/15 mm.	110	(142)
2-(8'-Nonen-1'-yl)-12-tridecen-1-ol.....	235/15 mm.	75	(142)
2-Nonyl-1-tridecanol.....	235/15 mm.	80	(142)
2-Decyl-1-tetradecanol.....	250/15 mm.	72	(142)
1-Phenyl-2,2-dibenzyl-1-ethanol.....	254-255/15 mm.	140	(142)

(a) Checked by reference 92.

(b) Checked by reference 125.

(c) Analysis: Calculated for $C_8H_{16}N_2O_3$: C, 51.04; H, 8.57; N, 14.89. Found: C, 50.65; H, 8.70; N, 15.1.

(d) The structure of this alcohol is in doubt, since it was made by treating a cyclohexene oxide with a Grignard reagent, a procedure which may lead to ring contraction. See reference 71 for a typical example.

(e) Analysis: Calculated for $C_7H_{12}N_2O_4$: C, 44.67; H, 6.43; N, 14.89. Found: C, 44.92; H, 6.13; N, 14.9.(f) Karrer (116) reported "triticol" as $C_{19}H_{38}O$ or $C_{20}H_{40}O$ (phytol) and probably the latter.

(g) A diallophanate.

(h) See reference 198 for probable structure.

(i) In the monoallophanate one of the hydroxyl groups—probably the tertiary one—has been replaced by chlorine.

(j) J. Kandel (Compt. rend. 205, 994 (1937)) reports the melting point as 151-152°C.

(k) Probably a mixture of the allophanates of α - and β -santalols.

TABLE 2
Allophanates of alcohols which are solid at 25°C.

ALCOHOL	MELTING POINT	MELTING POINT OF ALLOPHANATE	REFERENCES
	°C.	°C.	
Anisyl alcohol.....	25	180.25	(11)
1-Methylcyclopentanol.....	36	157	(32)
2-Hydroxyethyl benzoate.....	36-37	179-180	(83)
2-Phenyl-2-propanol.....	37	124-126 (dec.)	(194)
Myristyl alcohol.....	38	156.0-156.5 ^(a)	(21)
Menthol.....	43	133 ^(b) , 177 ^(b)	(9, 10)
		213	(11)
		215	(145)
Cetyl alcohol.....	49	70	(70)
		153	(95)
1,1,1-Trifluoro-2-propanol.....	52	159.7 (dec.)	(200)
Piperonyl alcohol.....	52-53	176.5 (dec.)	(11)
1-Octadecanol.....	58.5	154.3-155.5	(237)
1-Phenyl-1,2-ethanediol.....	67	168 (monoallophanate) ^(c)	(154)
		240-241 (diallophanate)	(154)
Benzhydrol.....	69	181-182 (dec.)	(194)
Glycolic acid.....	79	192	(207)
1,5-Bis-(4'-methoxyphenyl)-3-pentanol.....	80-81	136 (block) 137 (capillary)	(50) (50)
<i>l</i> -4-(2',3',6'-Trimethylphenyl)-2-butanol.....	88.5-90.5	173-175	(162, 163)
1,1,1-Trichloro-2-methyl-2-propanol.....	97	114	(234)
(Chlorobutanol)			
Coprosterol.....	103-104	210-211	(205)
1,2-Diphenyl-1,2-ethanediol.....	134	280 (diallophanate)	(154)
Sitosterol.....	137	246-247 (charring)	(205)
Dihydrocholesterol.....	141.5-142	255-256 ^(d)	(205)
(Cholestanol)			
Dihydrocryptosterol.....	145-146	249 (dec.)	(172)
(Dihydrolanosterol)			
Cholesterol.....	148.5	214-215 ^(e) 235-236 ^(d, f) 277-278	(102) (205) (63)
Testosterone.....	153.5-154.5	229.8-230.3 ^(g)	(92)
Ergosterol.....	163	250 ^(h)	(233)
Stigmasterol.....	169-170	241-242 ⁽ⁱ⁾	(102)
Pregnenolone.....	183-186	239.8-240.3 (dec.) ^(j)	(125)

^(a) Analysis: Calculated for C₁₆H₃₂N₂O₃: C, 63.96; H, 10.74; N, 9.33. Found: C, 64.00, 63.66; H, 10.59, 10.51; N, 9.49.

^(b) The structure of this alcohol is in doubt, since it was made by treating a cyclohexene oxide with a Grignard reagent, a procedure which may lead to ring contraction. See reference 71 for a typical example.

^(c) The position of the ester was not specified.

^(d) Checked by reference 125.

^(e) This is cholestry carbamate, which was obtained by Kaighin, using four-times-recrystallized cyanuric acid as the source of cyanic acid. Analysis: Calculated for C₂₈H₄₄NO₂: C, 78.27; H, 11.02; N, 3.26. Found: C, 78.10; H, 11.12; N, 3.28.

^(f) Checked by reference 92.

^(g) Analysis: Calculated for C₂₁H₃₀N₂O₄: C, 67.34; H, 8.09; N, 7.48. Found: C, 67.65; H, 7.90; N, 7.37.

^(h) Checked by reference 102.

⁽ⁱ⁾ Analysis: Calculated for C₃₁H₅₀N₂O₃: C, 74.65; H, 10.11; N, 5.62. Found: C, 74.62; H, 10.08; N, 6.09.

^(j) Analysis: Calculated for C₂₃H₃₄N₂O₄: C, 68.63; H, 8.51; N, 6.96. Found: C, 68.12; H, 8.38; N, 7.04.

TABLE 3
Allophanates of alcohols for which no physical constants are given

ALCOHOL	MELTING POINT OF ALLOPHANATE	REFERENCES
	°C.	
Ricinolein.....	61-62	(211)
6-Nonen-3-ol.....	96	(202)
<i>l</i> -Tetrahydrolavandulol.....	101-102	(183)
2-(<i>p</i> - <i>tert</i> -Butylphenoxy)ethanol.....	117-121	(237)
3-Methyl-2-oxo-3-pentanol.....	120.5	(189)
1,6-Nonadien-3-ol.....	125	(202)
Amyl lactate.....	131	(207)
9-Octadecen-1-ol.....	135	(3)
Undecenyl alcohol.....	143	(11)
5-Methyl-1-hexen-3-ol.....	147.5	(46)
Cyclohexylbutanol.....	148	(11)
1,2,2,2-Tetrachloroethanol.....	160	(212)
2,4,4-Trimethyl-1-hydroxymethylcyclohexane.....	163-164	(197)
<i>d,l</i> -Dihydrocyclolavandulol.....	163-164	(24)
2-Methyl-2-hepten-4-ol.....	171	(79)
Ethylvanillyl alcohol.....	173	(11)
Methyl saligenin.....	180	(11)
<i>tert</i> -Menthol.....	187	(11)
"Oxycamphor".....	204	(145)
Isopulegyl alcohol.....	219	(11)
β -Amyrin.....	250	(62)
	273-275	(13, 206)

TABLE 4
Allophanates of phenols

PHENOL	BOILING POINT	MELTING POINT OF ALLOPHANATE	REFERENCES
			°C.
2-Chlorophenol.....	175-176	179	(78)
Phenol.....	181	150 (dec.)	(208)
		180	(11)
		194	(23)
<i>o</i> -Cresol.....	191	185	(11)
<i>p</i> -Cresol.....	202	194	(11)
Methyl salicylate.....	223	175	(145)
Monomethoxytoluol.....	180-190/0.03 mm.	105-107	(105)
<hr/>			
	MELTING POINT		
2-Methoxyphenol.....	28	176	(145)
		200	(23)
2-Ethoxyphenol.....	28	212	(23)
2,2-Didodecyl-5,7,8-trimethyl-6-hydroxychroman.....	28	116	(97)
<i>o</i> -Cresol.....	30	185	(11)
<i>p</i> -Cresol.....	35-36	194	(11)
Phenol.....	42-43	180	(11)
		194	(23)
Thymol.....	51.5	190	(70)
2-Dodecyl-2,5,7,8-tetramethyl-6-hydroxychroman.....	60-61	180	(97)
2,3,5,6-Tetramethyl-4-chaulmooglyoxyphenol.....	61	174	(221)
2,3,6(2,3,5?)-Trimethyl-4-dodecyloxyphenol.....	81	184	(220)
2,2,5,7,8-Pentamethyl-6-hydroxychroman.....	93-94	230	(97)
2,3,5,6-Tetramethyl-4-dihydrophytyloxyphenol.....	97	178-180 182-183	(117) (65, 221)
Resorcinol.....	111	120	(207)
2-Ethyl-4,6,7-trimethyl-5-hydroxychroman.....	120	214	(106)
2,5,7,8-Tetramethyl-6-hydroxychroman.....	145	220	(97)
1,7-Dihydroxynaphthalene.....	178	243 (diallophanate)	(128)

TABLE 5
Allophanates of phenols for which no physical constants are given

PHENOL	MELTING POINT OF ALLOPHANATE	REFERENCES
Methylmethoxytocol.....	103 (not sharp)	(105)
<i>d,l</i> -5,7-Diethyltocol.....	107	(118)
2,2-Dicetyl-5,7,8-trimethyl-6-hydroxychroman.....	109	(96)
2,3,6-Trimethyl-4-[3'-methyl-5'-(2",2",6"-trimethylcyclohex-1"-yl)amyl-1'-oxy]phenol.....	128	(222)
γ-Tocopherol.....	135	(116)
	136-138	(8)
	137-140	(59)
2-Tridecyl-5,7-dimethyl-6-hydroxychroman..... (Isophytol)	143	(111)
7,8-Dimethyltocol.....	146	(110)
β-Tocopherol.....	137	(116)
	138-139	(8)
	143-144	(109, 110, 116)
	143.5-144.5	(206)
	147	(94)
2-Propyl-2-(4',8',12'-trimethyltridecyl)-5,7,8-trimethyl-6-hydroxychroman.....	146-148	(119)
2-Methyl-2-(4',8',12'-trimethyltridecyl)-7,8-trimethylene-6-hydroxychroman.....	149	(113)
2-Ethyl-2-(4',8',12'-trimethyltridecyl)-5,7,8-trimethyl-6-hydroxychroman.....	150-152	(119)
5,8-Dimethyltocol.....	154	(110)
<i>d,l</i> -5,7-Dimethyltocol..... (β-Tocopherol?)	150	(107)
	154-155	(107)
5,7-Dimethyl-8-methoxytocol.....	159-161	(105)
2,5,7,8-Tetramethyl-2-(4',8',12',16'-tetramethylheptadecyl)-6-hydroxychroman.....	161	(88)
2-Methyl-2-(4',8',12'-trimethyltridecyl)-7,8-tetramethylene-6-hydroxychroman.....	162-163	(113)
<i>d,l</i> -3,4-Dehydro-α-tocopherol.....	163	(89, 114)
<i>d,l</i> -5,8-Diethyl-7-methyltocol or <i>d,l</i> -7,8-Diethyl-5-methyltocol }	165	(108)
2,5,7,8-Tetramethyl-2-(4',8'-dimethylnonyl)-6-hydroxychroman.....	170	(87)
2,5,7-Trimethyl-8-ethyl-2-(4',8',12'-trimethyltridecyl)-6-hydroxychroman.....	170-171	(103)
2,3,5,6-Tetramethyl-4-[3'-methyl-5'-(2",2",6"-trimethylcyclohex-1"-yl)amyl-1'-oxy]phenol.....	173-174	(222)
2,3,5,6-Tetramethyl-4-phytyloxyphenol.....	176	(117)
α-Tocopherol.....	157-158	(8)
	158-159	(13)
	158-160	(62, 206)
	170-172	(98)
	172	(109, 114, 115)
Iso-α-tocopherol.....	175-176 (<i>d,l</i>)	(99)
	174-175	(95)
	176	(96, 100)

TABLE 5—*Concluded*

PHENOL	MELTING POINT OF ALLOPHANATE	REFERENCES
	°C.	
2-(6',10',14'-Trimethyl-2'-pentadecyl)-4,6,7-tri-methyl-5-hydroxycoumaran.....	176-180	(192)
5,7,8-Trimethyl-2,2-pentamethylene-6-hydroxy-chroman.....	184-186	(112)
2,3,5,6-Tetramethyl-4-(4'-methyl-2-pentyloxy)-phenol.....	206	(222)
2,3,5,6-Tetramethyl-4-dodecyloxyphenol.....	223	(221)
2,4,6,7-Tetramethyl-5-hydroxycoumaran.....	230	(222)
2,4,6,7-Tetramethyl-5-hydroxychroman.....	230	(222)

TABLE 6
Allophanates of miscellaneous alcohols

ALCOHOL	MELTING POINT OF ALLOPHANATE	REFERENCES
	°C.	
NH ₂ COOCH(CH ₃)S ₂ CH(CH ₃)OCOCONHCONH ₂	192-193	(216)
NH ₂ COCH(CH ₃)S ₂ CH(CH ₃)OCOCONHCONH ₂	213	(216)
β-Hydroxyethyltrimethylammonium iodide.....	254-255 (dec.)	(219)

VII. REFERENCES

- (1) AMATO, D.: Ber. 5, 290 (1872).
- (2) AMATO, D.: Gazz. chim. ital. 3, 467 (1873).
- (3) ANDRÉ, F.: Compt. rend. 185, 281 (1927).
- (4) BAAYER, A.: Ann. 114, 156 (1860).
- (5) BARBIER, H.: Helv. Chim. Acta 23, 524 (1940).
- (6) BARBIER, P., AND LOCQUIN, R.: Compt. rend. 158, 1554 (1914).
- (7) BASTERFIELD, S., AND WHELAN, M. S.: J. Am. Chem. Soc. 49, 3177 (1927).
- (8) BAXTER, J. G., ROBESON, C. D., TAYLOR, J. D., AND LEHMAN, R. W.: J. Am. Chem. Soc. 65, 918 (1943).
- (9) BEDOS, P.: Compt. rend. 181, 117 (1925).
- (10) BEDOS, P.: Bull. soc. chim. France [4] 39, 674 (1926).
- (11) BÉHAL, A.: Bull. soc. chim. France [4] 25, 473 (1919).
- (12) BÉNEZET, L.: Parfumerie 1, 153 (1943).
- (13) BERGEL, F., TODD, A., AND WORK, T.: J. Chem. Soc. 1938, 253.
- (14) BIILMANN, E., AND BJERRUM, J.: Ber. 50, 503 (1917).
- (15) BIRCKENBACH, L., AND KOLB, H.: Ber. 68, 895 (1935).
- (16) BISHOP, H. S.: Thesis, Massachusetts Institute of Technology, 1925.
- (17) BLAIR, J. S.: J. Am. Chem. Soc. 48, 87 (1926).
- (18) BLAIR, J. S.: J. Am. Chem. Soc. 48, 96 (1926).
- (19) BLOHM, H. W.: Thesis for the B.S. degree, Polytechnic Institute of Brooklyn, June, 1950.
- (20) BLOHM, H. W., AND BECKER, E. I.: Am. Doc. Inst., Doc. 2858; Chem. Abstracts 44, 8284i (1950).
- (21) BLOHM, H. W., AND BECKER, E. I.: J. Am. Chem. Soc. 72, 5342 (1950).
- (22) BÖTTCHER, B., AND BAUER, F.: Chem. Ber. 84, 289 (1951).
- (23) BOUGAULT, J., AND LEBOUCHQ, J.: Bull. soc. chim. France [4] 47, 594 (1930).

- (24) BOURQUIN, J.-P., SIMON, H. L., SCHÄPPI, G., STEINER, U., AND SCHINZ, H.: *Helv. Chim. Acta* **32**, 1564 (1949).
- (25) BRACK, K., AND SCHINZ, H.: *Helv. Chim. Acta* **34**, 1523 (1951).
- (26) BRACK, K., AND SCHINZ, H.: *Helv. Chim. Acta* **34**, 2009 (1951).
- (27) BRADFIELD, A. E., FRANCIS, E. M., PENFOLD, A. R., AND SIMONSEN, J. L.: *J. Chem. Soc.* **1936**, 1619.
- (28) BRADFIELD, A. E., PENFOLD, A. R., AND SIMONSEN, J. L.: *J. Chem. Soc.* **1935**, 309.
- (29) BRYAN, C. E., BURNETT, N. C., AND WEARN, R. B.: *J. Am. Chem. Soc.* **73**, 1848 (1951).
- (30) CAHOURS, A.: *Compt. rend.* **76**, 1383 (1873).
- (31) CALIEZI, A., LEDERER, E., AND SCHINZ, H.: *Helv. Chim. Acta* **34**, 879 (1951).
- (32) CHAVANNE, G., AND DEVOGEL, L.: *Bull. soc. chim. Belg.* **37**, 141 (1928).
- (33) CHERONIS, N. D., AND ENTRIKIN, J. B.: *Semimicro Qualitative Analysis*, pp. 371, 373, 375. Thomas Y. Crowell Company, New York (1947).
- (34) CLOSE, W. J.: Private communication.
- (35) COLOMBI, L., AND SCHINZ, H.: *Helv. Chim. Acta* **35**, 1066 (1952).
- (36) COLONGE, J., AND DREUX, J.: *Compt. rend.* **230**, 1776 (1950).
- (37) COLONGE, J., AND LAGIER, A.: *Bull. soc. chim. France* **1949**, 15.
- (38) CURTIUS, T., AND HOCHSCHWENDER, K.: *J. prakt. Chem.* **91**, 415 (1915).
- (39) CURTIUS, T., AND MÜLLER, C.: *Ber.* **34**, 2794 (1901).
- (40) DAINS, F. B., GREIDER, H. W., AND KIDWELL, C. H.: *J. Am. Chem. Soc.* **41**, 1004 (1919).
- (41) DAINS, F. B., AND WERTHEIM, E.: *J. Am. Chem. Soc.* **42**, 2303 (1920).
- (42) DAVIS, T. L., AND BLANCHARD, K. C.: *J. Am. Chem. Soc.* **51**, 1801 (1929).
- (43) DAVIS, T. L., AND BLANCHARD, K. C.: *J. Am. Chem. Soc.* **51**, 1806 (1929).
- (44) DELABY, R.: *Compt. rend.* **175**, 967 (1922).
- (45) DELABY, R.: *Ann. chim.* **20**, 196 (1923).
- (46) DELABY, R., AND BILLUART, A.: *Bull. soc. chim. France* **10**, 567 (1943).
- (47) DELABY, R., AND DUPIN, E.: *Bull. soc. chim. France* [5] **5**, 931 (1938).
- (48) DELABY, R., AND JANOT, M.: *Compt. rend.* **181**, 1146 (1925).
- (49) DELABY, R., AND MOREL, G.: *Bull. soc. chim. France* [4] **39**, 220 (1926).
- (50) DELÉPINE, M., AND SOSA, A.: *Bull. soc. chim. France* **9**, 771 (1942).
- (51) DEMUTH, R. L. (to R. L. Demuth Ltd.): British patent 636,181 (April 26, 1950).
- (52) DIELS, O.: *Ber.* **36**, 736 (1903).
- (53) DIELS, O., AND GOLLMANN, R.: *Ber.* **44**, 3158 (1911).
- (54) DIELS, O., AND NAVIASKY, P.: *Ber.* **37**, 3672 (1904).
- (55) DIELS, O., AND WOLF, B.: *Ber.* **39**, 686 (1906).
- (56) DOEVRE, J.: *Bull. soc. chim. France* [4] **45**, 351 (1929).
- (57) DOEVRE, J.: *Compt. rend.* **208**, 1658 (1939).
- (58) DOX, A. W., AND YODER, L.: *J. Am. Chem. Soc.* **45**, 723 (1923).
- (59) EMERSON, O. H., AND SMITH, L. I.: *J. Am. Chem. Soc.* **62**, 1869 (1940).
- (60) EPHRAIM, F.: *Ber.* **35**, 776 (1902).
- (61) ESS, P. R. VAN, AND WHITE, E. R. (to Shell Development Co.): U. S. patent 2,380,205 (July 10, 1945).
- (62) EVANS, H. M., EMERSON, O. H., AND EMERSON, G. A.: *J. Biol. Chem.* **113**, 319 (1926).
- (63) FABRE, R.: *Compt. rend.* **183**, 679 (1926).
- (64) FAVORSKIÍ, AL., AND ZALESSKIÍ-KIBARDINE, J.: *Bull. soc. chim. France* [4] **37**, 1227 (1925).
- (65) FERNHOLZ, E. (to Merck & Co.): U. S. patent 2,280,814 (April 28, 1942).
- (66) FREEDMAN, R. W., AND BECKER, E. I.: *J. Am. Chem. Soc.* **73**, 2366 (1951).
- (67) FREEDMAN, R. W., AND BECKER, E. I.: *J. Org. Chem.* **16**, 1701 (1951).
- (68) FREYERMUTH, H. B., AND RAIFORD, L. C.: *Proc. Iowa Acad. Sci.* **48**, 248 (1941).
- (69) GADDIS, A. M., AND BUTZ, L. W.: *J. Am. Chem. Soc.* **69**, 117 (1947).
- (70) GATTERMANN, L.: *Ann.* **244**, 29 (1888).
- (71) GAYLORD, N. G., AND BECKER, E. I.: *Chem. Revs.* **49**, 413 (1951).

- (72) GILLESPIE, D. T. C., MACBETH, A. K., AND MILLS, J. A.: *J. Chem. Soc.* **1948**, 996.
 (73) GILLIBRAND, M. I., AND LAMBERTON, A. H.: *J. Chem. Soc.* **1949**, 1883.
 (74) GODCHOT, M., AND BEDOS, P.: *Compt. rend.* **178**, 1184 (1924).
 (75) GODCHOT, M., AND BEDOS, P.: *Bull. soc. chim. France [4]* **37**, 1637 (1925).
 (76) GODCHOT, M., AND BEDOS, P.: *Compt. rend.* **180**, 751 (1925).
 (77) GODCHOT, M., AND BEDOS, P.: *Compt. rend.* **182**, 393 (1926).
 (78) GRANDIERE, J.: *Bull. soc. chim. France [4]* **35**, 187 (1924).
 (79) GRIGNARD, V., AND DUBIEN, M.: *Ann. chim.* **2**, 282 (1924).
 (80) GRIGNARD, V., AND ESCOURROU, R.: *Bull. soc. chim. France [4]* **37**, 546 (1925).
 (81) HALASZ, A.: *Compt. rend.* **209**, 319 (1939).
 (82) HALASZ, A.: *Bull. soc. chim. France [5]* **8**, 170 (1941).
 (83) HALASZ, A., AND ROVIRA, S.: *Bull. soc. chim. France [5]* **8**, 185 (1941).
 (84) HARDEGGER, E., REDLICH, D., AND GAL, A.: *Helv. Chim. Acta* **28**, 628 (1945).
 (85) HARISPE, J. V.: *Ann. chim.* **6**, 249 (1936).
 (86) HLASIWEITZ, H.: *Ann.* **134**, 115 (1865).
 (87) HOFFMANN-LA ROCHE & Co. A.-G.: Swiss patent 219,860 (June 16, 1942).
 (88) HOFFMANN-LA ROCHE & Co. A.-G.: Swiss patent 220,352 (July 1, 1942).
 (89) HOFFMANN-LA ROCHE & Co. A.-G.: Swiss patent 221,846 (September 1, 1942).
 (90) HOFMANN, A. W.: *Ber.* **4**, 262 (1871).
 (91) HURD, C. D., AND FAN, J. W.: *J. Am. Chem. Soc.* **73**, 110 (1951).
 (92) HWODZEK, W. P.: Thesis for the B.S. degree, Polytechnic Institute of Brooklyn, June, 1951.
 (93) JACOBSON, R. A.: *J. Am. Chem. Soc.* **60**, 1742 (1938).
 (94) JOHN, W.: *Z. physiol. Chem.* **250**, 11 (1937).
 (95) JOHN, W., AND GUNTHER, P.: *Ber.* **74B**, 879 (1941).
 (96) JOHN, W., GUNTHER, P., AND RATHMANN, F. H.: *Z. physiol. Chem.* **268**, 104 (1941).
 (97) JOHN, W., GUNTHER, P., AND SCHMEIL, M.: *Ber.* **71B**, 2637 (1938).
 (98) JOHN, W., AND HERRMANN, H.: *Z. physiol. Chem.* **273**, 191 (1942).
 (99) JOHN, W., AND PINI, H.: *Z. physiol. Chem.* **273**, 225 (1942).
 (100) JOHN, W., AND RATHMANN, F. H.: *Ber.* **74B**, 890 (1941).
 (101) JOHNSON, J. R., AND FERSTANDIG, L. L.: *Science* **110**, No. 2861, 441 (1949).
 (102) KAIGHIN, H. Y.: Thesis for the B.S. degree, Polytechnic Institute of Brooklyn, June, 1951.
 (103) KARRER, P. (to Hoffmann-La Roche, Inc.): U. S. patent 2,287,106 (June 23, 1942).
 (104) KARRER, P., AND BRETSCHER, E.: *Helv. Chim. Acta* **26**, 1758 (1943).
 (105) KARRER, P., AND DÜRR, K.: *Helv. Chim. Acta* **32**, 1361 (1949).
 (106) KARRER, P., ESCHER, R., AND RENTSCHLER, H.: *Helv. Chim. Acta* **22**, 1287 (1939).
 (107) KARRER, P., AND FRITZSCHE, H.: *Helv. Chim. Acta* **22**, 260 (1939).
 (108) KARRER, P., AND HOFFMANN, O.: *Helv. Chim. Acta* **23**, 1126 (1940).
 (109) KARRER, P., AND ISLER, O. (to Hoffmann-La Roche, Inc.): U. S. patent 2,411,967 (December 3, 1946).
 (110) KARRER, P., AND ISLER, O. (to Hoffmann-La Roche, Inc.): U. S. patent 2,411,968 (December 3, 1946).
 (111) KARRER, P., AND ISLER, O. (to Hoffmann-La Roche, Inc.): U. S. patent 2,411,969 (December 3, 1946).
 (112) KARRER, P., AND KEHRER, F.: *Helv. Chim. Acta* **25**, 29 (1942).
 (113) KARRER, P., AND KUGLER, A.: *Helv. Chim. Acta* **28**, 436 (1945).
 (114) KARRER, P., LEGLER, R. G., AND SCHWAB, G.: *Helv. Chim. Acta* **23**, 1132 (1940).
 (115) KARRER, P., AND RENTSCHLER, H.: *Helv. Chim. Acta* **26**, 1750 (1943).
 (116) KARRER, P., SALOMON, H., AND FRITZSCHE, H.: *Helv. Chim. Acta* **20**, 1422 (1937).
 (117) KARRER, P., SALOMON, H., AND FRITZSCHE, H.: *Helv. Chim. Acta* **21**, 309 (1938).
 (118) KARRER, P., AND SCHLÄPFER, R.: *Helv. Chim. Acta* **24**, 298 (1941).
 (119) KARRER, P., AND STÄHELIN, M.: *Helv. Chim. Acta* **28**, 438 (1945).

- (120) KIMEL, W., AND COPE, A. C.: J. Am. Chem. Soc. **65**, 1992 (1943).
- (121) LACOURT, A.: Bull. soc. chim. Belg. **36**, 346 (1927).
- (122) LANDOLT-BÖRNSTEIN: *Physikalisch-Chemische Tabellen*, 2nd supplement, Vol. II, p. 1079. Julius Springer Verlag, Berlin (1931).
- (123) LANE, E. S.: J. Chem. Soc. **1951**, 2764.
- (124) LANE, S. (to Imperial Chemical Industries, Ltd.) British patent 596,153 (December 30, 1947).
- (125) LAPKIN, M.: Thesis for the B.S. degree, Polytechnic Institute of Brooklyn, June 1951.
- (126) LAUCHENAUER, A., AND SCHINZ, H.: Helv. Chim. Acta **34**, 1514 (1951).
- (127) LEERS, L.: Bull. soc. chim. France [4] **39**, 421 (1926).
- (128) LEMAN, A.: Ann. chim. **9**, 357 (1938).
- (129) LIEBIG, J., AND WÖHLER, F.: Poggendorf's Ann. **20**, 396 (1830).
- (130) LIEBIG, J., AND WÖHLER, F.: Ann. **59**, 291 (1846).
- (131) LINHARD, M.: Ann. **535**, 267 (1938).
- (132) LINHARD, M.: Z. anorg. allgem. Chem. **236**, 200 (1938).
- (133) LINHARD, M.: Z. anorg. allgem. Chem. **239**, 155 (1938).
- (134) LINHARD, M., AND BETZ, K.: Ber. **73**, 177 (1940).
- (135) LOCQUIN, R., AND SUNG, W.: Compt. rend. **174**, 1427 (1922).
- (136) LOCQUIN, R., AND SUNG, W.: Compt. rend. **174**, 1551 (1922).
- (137) LOCQUIN, R., AND SUNG, W.: Compt. rend. **174**, 1713 (1922).
- (138) LOCQUIN, R., AND SUNG, W.: Bull. soc. chim. France [4] **35**, 597 (1924).
- (139) LOCQUIN, R., AND SUNG, W.: Bull. soc. chim. France [4] **35**, 604 (1924).
- (140) LOEB, M.: Ber. **19**, 2340 (1886).
- (141) MASTAGLI, P.: Compt. rend. **205**, 802 (1937).
- (142) MASTAGLI, P.: Ann. chim. **10**, 281 (1938).
- (143) MAUQUIN, C.: Ann. chim. phys. [8] **22**, 297 (1911).
- (144) MAVRODIN, A.: Compt. rend. **192**, 363 (1931).
- (145) MERCK, E.: German patent 248,164 (August 11, 1911).
- (146) MERCK, E. (O. DALMER AND C. DIEHL): German patent 553,148 (May 9, 1930).
- (147) MÉTAYER, M.: Bull. soc. chim. France **1951**, 802.
- (148) NAVES, Y. R.: Helv. Chim. Acta **26**, 162 (1943).
- (149) NAVES, Y. R.: Helv. Chim. Acta **29**, 553 (1946).
- (150) NAVES, Y. R.: Helv. Chim. Acta **29**, 1447 (1946).
- (151) NAVES, Y. R.: Compt. rend. **224**, 1830 (1947).
- (152) NAVES, Y. R., AND GRAMPOLOFF, A. V.: Helv. Chim. Acta **25**, 1500 (1942).
- (153) OVERLACH, M.: Berl. klin. Wochschr. **45**, 1403 (July 27); Chem. Abstracts **2**, 3259⁴ (1908).
- (154) PALFRAY, L., AND PANNELIER, R.: Bull. soc. chim. France [5] **4**, 1913 (1937).
- (155) PALFRAY, L., SABETAY, S., AND HALASZ, A.: Compt. rend. **208**, 289 (1939).
- (156) PALFRAY, L., SABETAY, S., AND KANDEL, J.: Compt. rend. **203**, 1376 (1936).
- (157) PALFRAY, L., SABETAY, S., AND MASTAGLI, P.: Compt. rend. **203**, 1523 (1936).
- (158) PALFRAY, L., SABETAY, S., AND RANGEL, A.: Compt. rend. **212**, 911 (1941).
- (159) PAUL, R.: Compt. rend. **192**, 1574 (1931).
- (160) PAUL, R.: Ann. chim. **18**, 303 (1932).
- (161) PAUL, R., AND RIOBÉ, O.: Compt. rend. **224**, 474 (1947).
- (162) PRELOG, V. von, AND FÜHRER, J.: Helv. Chim. Acta **28**, 583 (1945).
- (163) PRELOG, V. von, FÜHRER, J., HAGENBACH, R., AND FRICK, H.: Helv. Chim. Acta **30**, 113 (1947).
- (164) RAIFORD, L. C., AND FREYERMUTH, H. B.: J. Org. Chem. **8**, 174 (1943).
- (165) REMFRY, F. G. P.: J. Chem. Soc. **99**, 610 (1911).
- (166) RICHARDSON, T.: Ann. **23**, 138 (1837).
- (167) RIOBÉ, O.: Compt. rend. **225**, 384 (1947).
- (168) RIOBÉ, O.: Ann. chim. [12] **4**, 593 (1949).

- (169) RIOBÉ, O.: Compt. rend. **231**, 1312 (1950).
- (170) RIOBÉ, O.: Bull. soc. chim. France **1952**, 305.
- (171) ROVIRA, S.: Ann. chim. **20**, 660 (1945).
- (172) RUZICKA, L., DENSS, R., AND JEGER, O.: Helv. Chim. Acta **28**, 759 (1945).
- (173) RUZICKA, L., PLATTNER, PL. A., AND WILD, H.: Helv. Chim. Acta **28**, 395 (1945).
- (174) RUZICKA, L., SCHINZ, H., AND SUSZ, B. P.: Helv. Chim. Acta **27**, 1561 (1944).
- (175) RUZICKA, L., SEIDEL, C. F., SCHINZ, H., AND TAVEL, CH.: Helv. Chim. Acta **31**, 257 (1948).
- (176) SAYTZEFF, A.: Ann. **135**, 229 (1865).
- (177) SAYTZEFF, A.: Bull. soc. chim. **3**, 350 (1865).
- (178) SCHÄPPI, G., AND SEIDEL, C. F.: Helv. Chim. Acta **30**, 2199 (1947).
- (179) SCHAUM, K., AND SCHNEIDER, H.: Ber. **56**, 2460 (1923).
- (180) SCHIFF, H.: Ann. **291**, 367 (1896).
- (181) SCHINZ, H., AND MÜLLER, P. H.: Helv. Chim. Acta **27**, 57 (1944).
- (182) SCHINZ, H., RUZICKA, L., SEIDEL, C. F., AND TAVEL, CH.: Helv. Chim. Acta **30**, 1810 (1947).
- (183) SCHINZ, H., AND SCHÄPPI, G.: Helv. Chim. Acta **30**, 1483 (1947).
- (184) SCHINZ, H., AND SEIDEL, C. F.: Helv. Chim. Acta **25**, 1572 (1942).
- (185) SCHINZ, H., AND SIMON, H. L.: Helv. Chim. Acta **28**, 774 (1945).
- (186) SCHLIEPER, A.: Ann. **59**, 23 (1846).
- (187) SCHROETER, G., AND LEWINSKI, M.: Ber. **26**, 2171 (1893).
- (188) SCHÜTZ, F.: Nature **155**, 759 (1945).
- (189) SEIFERT, P., VOGEL, E., ROSSI, A., AND SCHINZ, H.: Helv. Chim. Acta **33**, 725 (1950).
- (190) SEN, H. K., AND BARAT, C.: Quart. J. Indian Chem. Soc. **2**, 77 (1925).
- (191) SIMON, H. L., KAUFMANN, AD., JR., AND SCHINZ, H.: Helv. Chim. Acta **29**, 1133 (1946).
- (192) SMITH, L. I., AND BOYACK, G. A.: J. Am. Chem. Soc. **70**, 2690 (1948).
- (193) SONTAG, D.: Ann. chim. [11] **1**, 359 (1934).
- (194) SPIELMAN, M. A., BARNES, J. D., AND CLOSE, W. J.: J. Am. Chem. Soc. **72**, 2520 (1950).
- (195) STAHL, G. W., AND COTTLE, D. S.: J. Am. Chem. Soc. **65**, 1782 (1943).
- (196) STEINER, U., AND SCHINZ, H.: Helv. Chim. Acta **34**, 1176 (1951).
- (197) STEINER, U., AND SCHINZ, H.: Helv. Chim. Acta **34**, 1508 (1951).
- (198) STOLL, M., AND HINDER, M.: Helv. Chim. Acta **33**, 1251 (1950).
- (199) STOLL, M., HINDER, M., AND PALLUY, ED.: Helv. Chim. Acta **34**, 1664 (1951).
- (200) SWARTS, F.: Bull. soc. chim. Belg. **38**, 99 (1929).
- (201) TAKEI, S., AND ONO, M.: J. Agr. Chem. Soc. Japan **15**, 193 (1939).
- (202) TAKEI, S., ONO, M., KUROIWA, Y., TAKAHATA, T., AND SIMA, T.: J. Agr. Chem. Soc. Japan **14**, 717 (1938).
- (203) TAKEI, S., ONO, M., AND SINOSAKI, K.: Ber. **73B**, 950 (1940).
- (204) TAKEI, S., SAKATO, Y., ONO, M., AND YOSIRO, K.: J. Agr. Chem. Soc. Japan **14**, 709 (1938).
- (205) TANGE, U., AND MCCOLLUM, E. V.: J. Biol. Chem. **76**, 445 (1928).
- (206) TODD, A., BERGEL, F., AND WORK, T. S.: Biochem. J. **31**, 2257 (1937).
- (207) TRAUBE, W.: Ber. **22**, 1572 (1889).
- (208) TUTTLE, D. K.: Jahresbericht **1857**, 451.
- (209) VAVON, G., AND IVANOW, D.: Compt. rend. **177**, 453 (1923).
- (210) VEREINIGTE CHININFABRIKEN ZIMMER & Co.: German patent 204,922 (August 31, 1907).
- (211) VEREINIGTE CHININFABRIKEN ZIMMER & Co.: German patent 211,197 (December 28, 1907).
- (212) VEREINIGTE CHININFABRIKEN ZIMMER & Co.: German patent 225,712 (June 27, 1908).
- (213) VODOZ, CH. A., AND SCHINZ, H.: Helv. Chim. Acta **33**, 1040 (1950).

- (214) VODOZ, CH. A., AND SCHINZ, H.: *Helv. Chim. Acta* **33**, 1321 (1950).
- (215) VOGEL, E., AND SCHINZ, H.: *Helv. Chim. Acta* **33**, 116 (1950).
- (216) WAGNER-JAUREGG, TH.: *Ann.* **561**, 87 (1948).
- (217) WALTSMANN, E., AND WOLF, E. (to Heberlein Patent Corporation): U. S. patent 2,380,133 (July 10, 1945).
- (218) WARREN, W. H., AND WILSON, F. E.: *Ber.* **68B**, 957 (1935).
- (219) WEIZMANN, M., AND BOGRACHOV, E.: *J. Am. Chem. Soc.* **67**, 154 (1945).
- (220) WERDER, F. VON (to Merck & Co.): U. S. patent 2,212,532 (August 27, 1940).
- (221) WERDER, F. VON, AND MOLL, T.: *Z. physiol. Chem.* **254**, 39 (1938).
- (222) WERDER, F. VON, MOLL, T., AND JUNG, F.: *Z. physiol. Chem.* **257**, 129 (1939).
- (223) WERNER, A. E. A.: *Sci. Proc. Roy. Dublin Soc.* **23**, 131 (1943).
- (224) WERNER, A. E. A., AND GRAY, J.: *Sci. Proc. Roy. Dublin Soc.* **24**, 77 (1946).
- (225) WERNER, A. E. A., AND GRAY, J.: *Sci. Proc. Roy. Dublin Soc.* **24**, 209 (1947).
- (226) WERNER, E. A.: *J. Chem. Soc.* **113**, 622 (1918).
- (227) WERNER, E. A., AND FEARON, W. R.: *J. Chem. Soc.* **117**, 1356 (1920).
- (228) WERNER, E. A., AND WERNER, A. E. A.: *Sci. Proc. Roy. Dublin Soc.* **23**, 137 (1943).
- (229) WILLIMANN, L., AND SCHINZ, H.: *Helv. Chim. Acta* **32**, 2151 (1949).
- (230) WILLSTÄTTER, R., AND DUISBERG, W.: *Ber.* **56B**, 2283 (1923).
- (231) WILM, TH.: *Ann.* **192**, 243 (1878).
- (232) WILM, TH., AND WISCHIN, G.: *Ann.* **147**, 150 (1868).
- (233) WINDAUS, A., AND RYGH, O.: *Nachr. Ges. Wiss. Göttingen, Math.-physik. Klasse* **1928**, No. 2, 202 (1928); *Chem. Abstracts* **24**, 4789⁷ (1930).
- (234) WOLFFENSTEIN, R., LOEWY, A., AND BACHSTEZ, M.: *Ber.* **48**, 2035 (1915).
- (235) ZEILE, K., AND MEYER, H.: *Chem. Ber.* **82**, 267 (1949).
- (236) ZEITSCHEL, O., AND SCHMIDT, H.: *Ber.* **59B**, 2298 (1926).

References added in proof

- (237) ADELSON, D. E. (to Shell Development Co.): U. S. patent 2,599,737 (June 10, 1952).
- (238) ADELSON, D. E., AND LARSEN, R. G. (to Shell Development Co.): U. S. patent 2,599,-736 (June 10, 1952).
- (239) N. V. DE BATAAFSCHE PETROLEUM MAATSCHAPPI: Dutch patent 66,832 (December 15, 1950).
- (240) SEIFERT, P., AND SCHINZ, H.: *Helv. Chim. Acta* **34**, 728 (1951).
- (241) TAKEI, S., AND ONO, M.: *J. Agr. Chem. Soc. Japan* **18**, 625 (1942); *Chem. Abstracts* **45**, 4201f (1951).