

THE REACTIONS OF MESITYL OXIDE

MARTIN HAUSER

*Applied Products Research Department, Stamford Research Laboratories
American Cyanamid Company, Stamford, Conn.*


Received July 31, 1962

CONTENTS

I. Introduction	311
II. The Reactivity of Mesityl Oxide	311
III. Reactions at the Double Bond	312
A. Simple Additions	312
B. Addition Accompanied by Secondary Reactions	314
C. Hydrogen Substitution and Miscellaneous Reactions	314
IV. Reactions at the Carbonyl Group	314
A. Grignard Reactions	314
B. Miscellaneous Reactions	315
V. Reactions at the α -Methyl Group	315
VI. Reactions at Several Sites—Cyclizations	315
A. Self-condensation	316
B. Condensations Giving Carbocyclic Compounds	317
1. Cyclopentane Derivatives	317
2. Cyclohexane Derivatives	317
3. Polycyclic Derivatives	318
C. Condensations Giving Heterocyclic Compounds	318
1. Five-membered or Smaller Rings—One Hetero Atom	318
2. Five-membered or Smaller Rings—Two Hetero Atoms	318
3. Six-membered Rings—One Hetero Atom	319
4. Six-membered Rings—Two Hetero Atoms	320
5. Seven-membered Rings	320
6. Polycyclic Compounds	320
VII. Reduction, Oxidation, and Pyrolysis	321
A. Reduction	321
B. Oxidation	322
C. Pyrolysis	322
VIII. References	322

I. INTRODUCTION

Mesityl oxide (I) is a tri-functional intermediate of considerable utility in the synthesis of a number of different types of organic compounds.



The possibility of effecting reaction at the double bond, the carbonyl group, the α -methyl group or several of these sites simultaneously is the key to the versatility of this material.

The chemistry of mesityl oxide has not previously been reviewed and this article will attempt to cover the literature through 1960. Where possible, significant post-1960 references have been included. Emphasis has been given to more recent work in the area since such work is, in general, more complete and offers better description of the experimental procedures.


Mesityl oxide is actually 4-methyl-3-pentene-2-one, but the name mesityl oxide is so well-established that it will be used throughout this review. All other

nomenclature conforms to "Chemical Abstracts" systems.

The poly-functionality of mesityl oxide provides the logical basis for this review of its reactions. For purposes of brevity, preparations of mesityl oxide and the well-known preparations of mesitylene and phorone from mesityl oxide have been omitted.

II. THE REACTIVITY OF MESITYL OXIDE

The majority of the reactions of mesityl oxide are those of either an α,β -unsaturated carbonyl compound or those of a methyl ketone. From structure II, it can be seen that both the carbonyl carbon atom and the β -ethylenic carbon atom are activated for nucleophilic attack.




Accordingly, mesityl oxide may undergo 1,2- or 1,4-

addition. The general tendency for nucleophilic reagents to add at the end of a conjugated system (β -carbon) is partially balanced by the greater charge on the carbonyl carbon closer to the electronegative oxygen atom which initiates the electronic shift. Due to these opposing factors, quantitative prediction of the site of reaction is difficult. However, as will be seen, the β -carbon atom appears more favored for nucleophilic addition.


The further description of mesityl oxide as a methyl ketone leads immediately to its reactivity in aldol- and Claisen-type condensations. The methyl group, rendered electron-deficient by the carbonyl oxygen, readily loses a proton to form the carbanion which is the attacking species in these reactions. The combination of the high electrophilicity of the β -carbon with this reactivity of the α -methyl group is an important factor in many of the cyclization reactions of mesityl oxide.

Some of the reactions of mesityl oxide are also complicated by the several tautomeric forms possible for the molecule. At equilibrium, at room temperature, mesityl oxide is a nine-to-one mixture (142) of I and 4-methyl-4-pentene-2-one (III, isomesityl oxide). Freshly prepared, mesityl oxide is primarily I and equilibrates with III very slowly unless base is present.



When base-catalyzed equilibration is combined with an attacking species which is a good hydrogen acceptor, the major reaction product may be derived from III rather than I. This will be illustrated later.

One further structural tautomer of mesityl oxide, the enol form IV, is possible. However, enolic reactions of mesityl oxide are very limited.



III. REACTIONS AT THE DOUBLE BOND

A. SIMPLE ADDITIONS

Mesityl oxide undergoes a variety of simple nucleophilic additions at the double bond. These reactions all proceed by Markownikoff addition as would be expected from the polarization of the molecule (structure II). Furthermore, the activating influence of the carbonyl oxygen makes addition facile and rapid. In only a few cases is a small amount of the counter-Markownikoff product isolated.


Mesityl oxide readily adds halogens, hydrohalic acids and hypohalic acids. The results of these reactions are summarized in Table I. Often, the reaction product is unstable and yields have either been omitted or refer to crude products.

This instability often allows secondary reactions to be performed directly in solution. For example, chlorina-


TABLE I
REACTIONS OF MESITYL OXIDE WITH HALOGENS, HYDROHALIC ACIDS AND HYPOHALIC ACIDS

Reagent	Product:		Yield, %	References
	(CH ₃) ₂ CXCH ₂ COCH ₃	Y		
Cl ₂	X	Y	30	(41)
Br ₂	Cl	Cl	..	(117, 48)
I ₂	Br	Br	..	(49)
HCl	I	I	..	(128)
HBr	Cl	H	..	(131)
HI	Br	H	..	(118)
HOCl	I	H	..	(115, 139)
HOBr	OH	Cl	60	(115)
HOI	OH	Br	25	(115)
	OH	I	..	(115)

tion of mesityl oxide in the presence of sodium bicarbonate (145) gives 60-70% of 3-chloro-4-methyl-3-pentene-2-one (V), formed by dehydrochlorination of the original dichloro-adduct.




The addition of ammonia and simple amines to mesityl oxide is straightforward and requires no catalyst. However, the 4-amino-4-methyl-2-pentanones (VI) obtained as products are again unstable and are usually isolated as salts.



R ₁	R ₂	Yield, %	Reference
H	H	69	(67)
Me	H	"Quantitative"	(71)
Me	Me	"Good"	(71)

"Good" yields of 4-(N-benzyl-N-hydroxylamino)-4-methyl-2-pentanone (VII) have been reported in a similar reaction of mesityl oxide and N-benzyl-hydroxylamine (134). The yield of the corresponding reaction with N-phenylhydroxylamine is described as very poor.



The addition of alcohols to mesityl oxide to give 4-alkoxy-4-methyl-2-pentanones (VIII) may be acid- (73) or base-catalyzed (144). In general, basic catalysis appears to produce better yields. Some typical reactions are listed in Table II.


TABLE II
REACTIONS OF MESITYL OXIDE WITH ALCOHOLS

ROH, R =	Yield of (CH ₃) ₂ C(OR)CH ₂ - COCH ₃ (VIII), %	Reference
Me	a	(144)
Et	21	(73)
Et	82	(144)
Pr	..	(73)
Bu	25	(73)
Bu	b	(144)
iso-Bu	..	(73)
iso-Amyl	..	(73)
Benzyl	..	(73)



a "High" yield based on 45-50% conversion. b "High" yield based on 32% conversion.

More recently (101), anion exchange resins have been used to catalyze the additions. A yield of 90–95% (40–60% conversion) is obtained with methanol and 82% (27% conversion) with ethanol.




Hydrogen sulfide adds to mesityl oxide with triethylamine as catalyst (53) to yield 80% of 4-mercapto-4-methyl-2-pentanone (IX). With a potassium hydroxide catalyst (7), an unspecified amount of the bis-compound X is formed as by-product.




The reaction of mesityl oxide and thiocyanic acid (24) results in a 72% yield of 4-methyl-4-thiocyanato-2-pentanone (XI). Similarly, mesityl oxide adds sulfuric acid (86) to give 2-methyl-4-oxo-2-pentane-sulfonic acid (XII, as the sodium salt). The identical product is obtained by direct addition of sodium bisulfite (61).



The addition of aromatic compounds to mesityl oxide is illustrated by aluminum chloride-catalyzed reactions with benzene (74) and chlorobenzene (40). The products are, respectively, 4-methyl-4-phenyl-2-pentanone (XIII, 72–89% yield) and 4-(*p*-chlorophenyl)-4-methyl-2-pentanone (XIV, 62% yield). Fluorene adds to mesityl oxide in 15–20% yield using a potassium hydroxide catalyst (54). The product, 4-(9-fluorenyl)-4-methyl-2-pentanone (XV), may be obtained in 55% yield by the use of 9-fluorenylsodium.



The heterocyclic 2-methylfuran (97, 165) and 2-ethylfuran (165) undergo acid-catalyzed additions to give XVIa (50%) and XVIb, respectively.





A variety of phosphorus- and/or sulfur-containing organic acids, including dialkyl phosphites (124), dialkyl dithiophosphates (72), thiolacetic acid (23, 150) and ethyl alkylphosphonites (125), have been allowed to react with mesityl oxide. High yields of the simple addition products are generally obtained.

These reactions may be uncatalyzed, base-catalyzed or free radical-catalyzed and are summarized in Table III.

TABLE III
REACTIONS OF MESITYL OXIDE WITH PHOSPHORUS AND/OR SULFUR-CONTAINING ORGANIC ACIDS

Reagent	Catalyst	Product yield, %	Reference
(RO) ₂ POH, R =		(CH ₃) ₂ C(P(O)(OR) ₂)CH ₂ COCH ₃	
Me	NaOMe	70	(124)
Bu	NaOBu	77	(124)
iso-Bu	NaOBu-iso	69	(124)
(RO) ₂ PS ₂ H		(CH ₃) ₂ C(SP(S)(OR) ₂)CH ₂ COCH ₃	
Et	...	88	(72)
2-ClEt	...	86	(72)
iso-Pr	...	76	(72)
sec-Bu	...	99	(72)
4-Me-2-pentyl	...	85	(72)
<i>n</i> -tetradecyl	...	81	(72)
C ₆ H ₅	...	74	(72)
CH ₃ COSH		(CH ₃) ₂ C(SCOCH ₃)CH ₂ COCH ₃	
CH ₃ COSH	Piperidine	64	(150)
CH ₃ COSH	Benzoyl peroxide	92	(23)
EtP(O)(OR)H		(CH ₃) ₂ C(P(O)(OR)Et)CH ₂ COCH ₃	
Et	NaOEt	46	(125)
Bu	NaOBu	36	(125)

Treatment of mesityl oxide with phosphorus trichloride and acetic anhydride (42), followed by hydrolysis, gives 33% of the phosphite XVIIa. The butyl ester XVIIb is obtained in 20% yield when the intermediate is treated with butyl alcohol and sodium rather than hydrolyzed. By direct action of butyldichlorophosphine-acetic anhydride and diphenoxychlorophosphine-acetic anhydride on mesityl oxide, the addition products XVIIc (74%) and XVIId (41%) are obtained.




The Ritter reaction of nitriles and mesityl oxide using a sulfuric acid catalyst (135) results in fair yields of the substituted amides. These reactions along with several other miscellaneous simple additions are listed in Table IV.

TABLE IV


Reagent	Catalyst	Product (CH ₃) ₂ C(R)-CH ₂ COCH ₃ , R =	Yield, %	Reference
C ₆ H ₅ CN	H ₂ SO ₄	NHCOC ₆ H ₅	45–55	(135)
CH ₃ CN	H ₂ SO ₄	NHCOCH ₃	25	(135)
CH(NO ₂) ₂	...	C(NO ₂) ₂	...	(153)
NaN ₃	...	N ₃	38	(18)
CH ₃ NO ₂	Et ₂ NH	CH ₂ NO ₂	65	(88)

The peroxide-catalyzed addition of simple aldehydes to mesityl oxide (116) gives fair to good yields of diketones XVIII, although conversions are rather low.





R	Conversion, %	Yield, %
CH ₃	10	31
<i>n</i> -C ₂ H ₅	41	97
<i>n</i> -C ₆ H ₁₃	..	37–61

In the case of butyraldehyde, a small amount (about one part in ten) of the counter-Markownikoff product XIX is isolated from the reaction.





B. ADDITION ACCOMPANIED BY SECONDARY REACTIONS


Several other addition reactions of mesityl oxide have been described but these are somewhat complicated by side-reactions or reactions at other sites. Potassium cyanide adds to mesityl oxide (99) giving the nitrile XX (mesitonitrile) in good yields. Compound XX may then react, *in situ*, to give the cyanohydrin XXI which cyclizes to the pyrrolidone XXII (mesitylic acid).



A similar situation exists with hydroxylamine. The hydroxamic acid XXIII is formed by addition of mesityl oxide, but this is accompanied by formation of mesityl oxide oxime (XXIV) and the cyclized product XXV, arising from an internal condensation of XXIII (64).




The reaction of elemental sulfur, hydrogen and mesityl oxide, catalyzed by cobalt polysulfide-acetic acid (50), gives 15% of 4-mercapto-2-methyl-2-pentene (XXVI) and 20% of 2,4-dimercapto-2-methylpentane (XXVII), formed by addition and oxygen replacement. The reaction of ethanethiol with mesityl oxide (123) may result in counter-Markownikoff addition yielding 2,2,3-tris-(ethylmercapto)-4-methylpentane (XXVIII), but no evidence exists that the "abnormal" adduct is actually the product.




C. HYDROGEN SUBSTITUTION AND MISCELLANEOUS REACTIONS

Several reactions involving substitution of the hydrogen on the α -ethylenic carbon atom have been reported. Treatment of mesityl oxide with dimethyl sulfate using sodium *tert*-amylate as catalyst (34) yields 41% of 3,4-dimethyl-3-pentene-2-one (XXIX). The similar reaction using benzyl bromide gives an unspecified amount of the 3-benzyl analog of XXIX. The reaction of ethylene with mesityl oxide (75) results in a 7% yield of 3-ethyl-4-methyl-3-pentene-2-one (XXX).



The base-catalyzed cyanoethylation of mesityl oxide with acrylonitrile (25) gives 10–15% of XXXI and 74% of XXXII. The formation of XXXII illustrates the double bond shift which may occur in the presence of strong base and a good hydrogen acceptor.



IV. REACTIONS AT THE CARBONYL GROUP

A. GRIGNARD REACTIONS

The majority of reactions at the carbonyl group of mesityl oxide are Grignard or Grignard-type. As opposed to simple addition reactions at the double bond which are generally based on 1,4-addition, these may be considered as 1,2-additions across the carbonyl double bond. There is a wide range of yields in mesityl oxide-Grignard reactions and a number of low yield syntheses have been included because of the unusual products obtainable from them. Table V summarizes some of these reactions.


The mode of dehydration of the product alcohols can be changed by variation in the reaction conditions. For example, with ethyl Grignard reagents dehydration

TABLE V
GRIGNARD REACTIONS OF MESITYL OXIDE

Reagent	Product	Yield, %	Reference
MeMgI	Me ₂ C=CHC(OH)Me ₂	29	(51)
MeMgI	Me ₂ C=CHCMe=CH ₂	..	(56)
EtMgBr	Me ₂ C=CHC(Et)=CH ₂	12	(83)
EtMgBr	Me ₂ C=CHC(Me)=CHMe	51	(47)
EtMgI	Me ₂ C=CHC(Me)=CHMe	53	(141)
<i>tert</i> -BuMgCl	Me ₂ C=CHC(CMe ₃)=CH ₂ + CH ₂ =CMeCH=CMeCMe ₃	35	(154)
isoamylMgBr	CH ₂ =CMeCH ₂ CMe=CHCH ₂ - CHMe ₂	...	(84)
C ₆ H ₅ MgBr	Me ₂ C=CHC(C ₆ H ₅)=CH ₂	"Low"	(83)
C ₆ H ₅ MgBr	Me ₂ C=C=C(C ₆ H ₅)Me	44	(87)
C ₆ H ₅ CH ₂ MgCl	Me ₂ C=CHCMe(OH)CH ₂ C ₆ H ₅	26	(57)
CH ₂ =CHCH ₂ MgBr	Me ₂ C=CHCMe(OH)CH ₂ CH=CH ₂	90	(109)
Me ₂ C=CHLi	Me ₂ C=CHCMe(OH)CH=CMe ₂	8	(20)
Me ₂ C=CMeLi	Me ₂ C=CHCMe(OH)CMe=CMe ₂	67	(21)
CH ₂ =CMeLi	Me ₂ C=CHCMe(OH)CMe=CH ₂	74	(21)
C ₂ H ₅ -Ca	Me ₂ C=CHCMe(OH)C≡CH	75	(114)




may occur between the hydroxyl and ethyl or methyl groups (47, 83, 141); with phenyl Grignards, dehydration occurs with the α -methyl group or internally with the α -ethylenic carbon giving an allene (83, 87). In the case of *tert*-butyl magnesium chloride (154) one of the products obtained is that from a double-bond shift and internal dehydration, while with isoamylmagnesium bromide (84) the double bond shift also occurs but the product is still formed by external dehydration with the amyl group. No systematic investigation of the various reactions of mesityl oxide with Grignard reagents has as yet been made and this appears a fertile field for investigation.

The reaction of 9-fluorenylmagnesium bromide and mesityl oxide has also been reported (147) and gives 30% of the hydrocarbon XXXIII. Here, the dehydration involves the 9-hydrogen of fluorene.



B. MISCELLANEOUS REACTIONS




In addition to Grignard-type reactions, mesityl oxide undergoes a few other miscellaneous reactions at the carbonyl. Halogenolysis with phosphorus pentachloride (10) gives 4,4-dichloro-2-methyl-2-pentene (XXXIV). The acid-catalyzed reaction with ketene (58) yields 47% of the enol acetate of mesityl oxide (XXXV). This is one of the very few enolic reactions of mesityl oxide described in the literature. Cyanoacetic acid adds to mesityl oxide at the carbonyl group under the influence of piperidine hydrochloride (90) ultimately to give XXXVI. Finally, the reaction of mesityl oxide with potassium cyanide, followed by treatment with ammonia and hydrogen sulfide, results in an over-all yield of 14% of the thioamide XXXVII (1).






V. REACTIONS AT THE α -METHYL GROUP

Aldol-type reactions of mesityl oxide and various carbonyl compounds have been reported and comprise the bulk of the material on simple reactions at the α -methyl group. In the presence of aniline-N-magnesium bromide, mesityl oxide undergoes self-condensation to give 67% of a seven to one mixture of XXXVIII and its dehydration product XXXIX (28). Similar catalysis of the reaction of mesityl oxide and acetophenone yields 31% of XL if mesityl oxide is added to





the catalyst solution first and 33% of XLI if acetophenone is added first.



Aldol condensations of mesityl oxide with benzaldehyde (acid catalyzed) (31), camphenaldehyde (base catalyzed, 43% yield) (5) and 4-(2,3,3-trimethylcyclopentenyl)-acetaldehyde (base catalyzed, 35%) (6) have also been reported and give compounds XLII, XLIII and XLIV, respectively.



Ethyl oxalate undergoes a base-catalyzed Claisen condensation with mesityl oxide (30) to give a 100% yield (70% conversion) of the diketone XLV. When trioxymethylene is condensed with mesityl oxide using an anion exchange resin (160) there is obtained at an early stage of the reaction 10% of the condensation product XLVI and 30% of XLVII, formed by a double bond shift. On prolonged heating with magnesia, the mixture isolated consists of 25% XLVI and 20% XLVII. Some of the bis-hydroxymethyl compound XLVIII is also isolated.



VI. REACTIONS AT SEVERAL SITES—CYCLIZATIONS


The outstanding utility of mesityl oxide lies in its ability to undergo a variety of condensation reactions leading to carbocyclic and heterocyclic compounds. This ability is a direct result of the polyfunctionality of the molecule and the fact that any of five different sites, the α -methyl group, the carbonyl group, the α -ethylenic carbon, the β -ethylenic carbon and either

exchange resin (161) yields 16% of LI along with some 4-isopropylidene-3,5,5-trimethyl-2-cyclohexenone (LIV). Compound LIV is formed through Type 4 condensation of one molecule and an unusual two carbon reaction of the other molecule. This can be formulated as substitution at the α -ethylenic carbon and an aldol reaction and dehydration at the carbonyl group.

B. CONDENSATIONS GIVING CARBOCYCLIC COMPOUNDS


1. Cyclopentane Derivatives

The condensation of mesityl oxide and benzil (base-catalyzed) (82) gives 16% of 3,4-diphenyl-4-hydroxy-5-isopropylidene-2-cyclopentenone (LV). This is the only five-membered carbocyclic compound derived from mesityl oxide (except by self-condensation) and is formed by a Type 3b reaction.




2. Cyclohexane Derivatives

The majority of carbocyclic compounds obtained from mesityl oxide are six-membered and the formation of a number of these materials serve as excellent examples of several condensation types. Type 4 condensation is illustrated by the sodium-catalyzed reaction of ethyl or methyl cyanoacetate with mesityl oxide (133), yielding a mixture of the open-chain adduct LVI and the cyclohexadione LVII. LVI, when separated, is readily cyclized to LVII, establishing the probable mechanism.




A group of similar reactions is summarized below.





Reactant	Catalyst	Product, R =	Yield, %	Reference
$\text{C}_6\text{H}_5\text{CH}_2\text{CO}_2\text{Et}$	NaOEt	C_6H_5	57	(17)
$\text{CH}_2(\text{CO}_2\text{Et})_2$	NaOEt	H	70	(95)
$\text{CH}_2(\text{CO}_2\text{Et})(\text{Me})_2$	Na or KOH	$\text{CO}_2\text{Et}(\text{Me})$..	(152)

Benzyl phenyl ketone (desoxybenzoin) also undergoes a Type 4 reaction with mesityl oxide (78). The final product LVIII (52% yield) arises through dehydration of the intermediate β -keto alcohol. The linear intermediate LIX may also be isolated in 25–50% yield by use of milder conditions. A similar base-catalyzed


reaction of mesityl oxide and ethyl phenyl ketone (27) yields 20% of the cyclohexenone LX.



LIX







The base-catalyzed reaction of N,N-diethylacetamide with mesityl oxide (149) illustrates a Type 3a condensation giving a cyclohexane derivative. The product, N,N-diethyl 4,6,6-trimethyl-3-cyclohexen-2-one carboxamide (LXIa), is obtained in 61% yield. The corresponding reaction with N-ethyl-N-(2-methylphenyl)-acetoacetamide (148) yields the carboxamide LXIb. Mesityl oxide also reacts with methyl ethyl ketone and benzyl methyl ketone (27) in a like manner, the products isolated being LXIc (10% yield) and LXId (73% yield), respectively.




LXIa, R = $\text{CON}(\text{C}_2\text{H}_5)_2$; c, R = CH_3
 b, R = $\text{CON}(\text{C}_2\text{H}_5)\text{C}_6\text{H}_4\text{CH}_3\text{-o}$; d, R = C_6H_5

Mesityl oxide and ethyl acetoacetate condense in the presence of sodium (102) to give a low yield of ethyl 4-hydroxy-4,6,6-trimethyl-2-cyclohexanone carboxylate (LXII). More rigorous conditions (89) result in dehydration and decarboxylation of LXII so that isophorone (LI) is eventually obtained. A similar reaction with ethyl 3-methylacetoacetate yields 44% of the 1-methyl analog of LXII (LXIII), 7% of the dehydrated compound LXIV and 8% of the isophorone analog LXV.




The condensation of mesityl oxide with 2,3-dimethylbutadiene (55) proceeds by a Diels–Alder reaction (Type 2) to give 5-acetyl-1,2,4,4-tetramethylcyclohexene (LXVI). The reaction of mesityl oxide and methyl vinyl ketone, catalyzed by sodium *tert*-amylate, has been surrounded by some controversy as to the actual

products (15, 16, 35, 108). It now seems that the material isolated from this condensation is a mixture of 3-methyl-6-isopropylidene-2-cyclohexenone (LXVII, piperetinone) and isoxylitone (LV). Compound LXVII arises from a Type 3b reaction, while LV arises from self-condensation of two molecules of mesityl oxide. The combined yield is about 40%.




3. Polycyclic Derivatives

The ring-fusion of mesityl oxide with pseudocumene (1,2,4-trimethylbenzene) (140) is catalyzed by aluminum chloride and yields 25% of 1,1,3,4,5,7-hexamethylindene (LXVIIIa). The reactions of *m*- and *p*-xylene (32) give 27% of 1,1,3,4,6-pentamethylindene (LXVIIIb) and 28% of 1,1,3,4,7-pentamethylindene (LXVIIIc), respectively.




LXVIIIa, $R_1 = R_2 = R_3 = R_4 = R_5 = R_7 = \text{CH}_3$; $R_6 = \text{H}$
 b, $R_1 = R_2 = R_3 = R_4 = R_5 = \text{CH}_3$; $R_6 = R_7 = \text{H}$
 c, $R_1 = R_2 = R_3 = R_4 = R_7 = \text{CH}_3$; $R_5 = R_6 = \text{H}$


The probable mechanism of these Type 3a condensations is demonstrated by isolation of 4-(3,5-dimethylphenyl)-4-methyl-2-pentanone (LXIX) from the reaction giving LXVIIIb.



Mesityl oxide also reacts with catechol (1,2-dihydroxybenzene) (11). In this case, however, a rather complicated secondary reaction ensues and the product finally isolated is the bis-spirohydrindene LXX (17% yield).




The base-catalyzed reaction of mesityl oxide with 2-hydroxy-cyclohexanone (33) proceeds by a Type 4 condensation and dehydration of the intermediate alcohol to yield 46% of LXXIa. A similar reaction with 4-methyl-2-hydroxycyclohexanone gives a 35-40% yield of LXXIb.




C. CONDENSATIONS GIVING HETEROCYCLIC COMPOUNDS

1. Five-membered or Smaller Rings—One Hetero Atom

The methoxide-catalyzed reaction of mesityl oxide with ethyl chloroacetate (69) yields 55% of the oxetane LXXII ($R = \text{CH}_2\text{CH}_3$). A small amount of the methyl analog LXXIII ($R = \text{CH}_3$) is also formed by preliminary alcoholysis of ethyl chloroacetate by methoxide. The reaction appears to proceed by an initial 1,2-addition and subsequent cyclization by dehydrohalogenation. Formaldehyde and ethyl oxalate condense with mesityl oxide in the presence of methoxide (126) to give 53% of 2,3-diketo-4-isobutenyltetrahydrofuran (LXXIII). The reaction involves a rather unusual donation of the carbonyl carbon atom and α -methyl group as a two-carbon fragment by mesityl oxide.



Low yields of both 2-(2-hydroxyisobutyl)-2-methyl-3,4,5-triketopyrrole (LXXIV) and 5-methyl-5-(2-methyl-1-propenyl)-hydantoin (LXXV) may be obtained by the reaction of mesityl oxide with potassium cyanide and ammonium carbonate (68). The formation of both of these materials illustrates Type 1 condensation except that LXXIV also contains a hydroxylic function arising from the addition of the elements of water to the carbon-carbon double bond.



2. Five-membered or Smaller Rings—Two Hetero Atoms

The reaction of mesityl oxide with 1,2-glycols and similar compounds leads to formation of 1,3-dioxolanes by condensation at the carbonyl (Type 1). The general reaction and some specific examples are listed in Table VI.

The condensation of mesityl oxide with 2,3-epoxy-3-methyl-6-heptene-4-yne (120) also yields 23% of a substituted dioxolane LXXVI.





TABLE VI
FORMATION OF 1,3-DIOXOLANES


Reagent	Product	Yield, %	Reference
$\begin{array}{c} R_1R_2C(OH) \\ \\ R_3R_4C(OH) \end{array}$	$\begin{array}{c} (CH_3)_2C=CH-C-CH_3 \\ \quad \\ O \quad O \\ \quad \\ R_1R_2C-CR_3R_4 \end{array}$		
$R_1 = R_2 = R_3 = R_4 = H$		70-90	(132)
$R_1 = R_2 = R_3 = H; R_4 = CH_3$		70-90	(132)
$R_1 = R_3 = H; R_2 = R_4 = C_6H_5$...	(85)
$R_1 = R_2 = R_3 = R_4 = CH_3$...	(85)
$R_1 = R_3 = CH_3; R_2 = R_4 = C_6H_5$...	(85)

The condensation of mesityl oxide with some of the common "carbonyl" reagents provides an excellent route to the formation of the pyrazole nucleus. The initially formed carbonyl derivative readily cyclizes, with the over-all reaction being of Type 3a. However, as contrasted to previously mentioned condensations of this type, the primary reaction is a 1,2-addition rather than a 1,4-addition.


The reaction of mesityl oxide and hydrazine (38) gives 48% of 3,5,5-trimethyl- Δ^2 -pyrazoline (LXXVIIa). In the corresponding reaction with phenylhydrazine (8), the phenylhydrazone may be isolated or the reaction may be carried directly to 1-phenyl-3,5,5-trimethyl- Δ^2 -pyrazoline (LXXVIIb). The over-all yield is 60%. A similar reaction series has been described for *p*-nitrophenylhydrazine; however, by use of *p*-nitrophenylhydrazine hydrochloride, a direct quantitative yield of the pyrazoline LXXVIIc may be obtained. The analogous reaction with semicarbazide (65) can give either mesityl oxide semicarbazone or the pyrazoline carboxamide LXXVIIId.




The condensation of mesityl oxide with diazomethane and diazoethane (2) also leads to pyrazolines, 3-acetyl-4,4-dimethyl- Δ^1 -pyrazoline (LXXVIII, R = H, 44% yield) and 3-acetyl-4,4,5-trimethyl- Δ^1 -pyrazoline (LXXVIII, R = CH₃), respectively. In these cases, ring formation is of Type 2.



A recent report (76) describes the reaction of mesityl oxide with diazomethane in the presence of boron trifluoride etherate. The products isolated are the ketones LXXIX and LXXX, rather than the pyrazoline.




Mesityl oxide condenses with phosphorus trichloride-acetic anhydride to give 22% of the oxaphosphole oxide LXXXI (4).




3. Six-membered Rings—One Hetero Atom


The reaction of diethyl oxalate and two moles of mesityl oxide (137) gives an 18% yield of compound LXXXII formed by two Claisen-type condensations. This product, in the presence of acid, undergoes cyclization to the bis-pyrone LXXXIII in 80% yield. The over-all process illustrates a Type 4 reaction. A similar reaction of two moles of ethyl oxalate with one mole of mesityl oxide and one mole of acetone yields 17% of the adduct LXXXIV which also undergoes acid-catalyzed cyclization (50%) to the pyrone LXXXV.



A pyrone, LXXXVI, is also formed in the boron trifluoride-catalyzed Type 3a condensation of mesityl oxide with ketene (164). The yield is 80%.





The single six-membered mononitrogen heterocycle reported (127) from mesityl oxide is 6-hydroxy-4,4,6-trimethyl-2-piperidone-3-carbonitrile (LXXXVII), prepared in quantitative yield by a base-catalyzed reaction with cyanoacetamide. This is also a Type 3a condensation, but here, dehydration of the alcohol has not occurred.




When mesityl oxide is treated with sulfuric acid hydrate and acetic anhydride (138), a 22% yield of

the pyrrilium salt (LXXXVIII, as the perchlorate) is obtained. By treatment of mesityl oxide with thionyl chloride (121) a compound tentatively identified as the thiopyrrilium salt LXXXIX is obtained. This is the one example of possible Type 5 condensation in the literature.




4. Six-membered Rings—Two Hetero Atoms


The condensation of mesityl oxide and ammonium thiocyanate (70) gives 84% of 2-thio-4,4,6-trimethyl-dihydropyrimidine (XC_a), while the base-catalyzed condensation of guanidine and mesityl oxide (146) yields 85–91% of 2-amino-4,4,6-trimethyl-dihydropyrimidine (XC_b). Both these reactions illustrate Type 3a condensation with dehydration of the intermediate. Initial 1,4-addition of guanidine is shown by isolation of an 80% yield of the cyclizable linear intermediate XCI using guanidine thiocyanate. A similar reaction with benzamidine gives the 2-phenyl analog (XC_c).




Tetrahydropyrimidines may be obtained by the action of ammonia and acetone on mesityl oxide (19). If mesityl oxide is treated with ammonia and acetone and additional ammonia then added, a 66% yield of 2,2,4,4,6-pentamethyltetrahydropyrimidine (XCII) is isolated. By the initial reaction of ammonia and mesityl oxide, followed by treatment with more mesityl oxide and ammonia (omitting acetone), 11% of pyrimidine XCIII is obtained.



The reaction of mesityl oxide and dithiocarbamic acid, using an acid catalyst (81), yields 91% of 2-mercapto-4,6,6-trimethylthiazine (XCIV). Less rigorous reaction conditions allow isolation of the 4-hydroxy-dihydro compound XCV. This is another example of Type 3a condensation.



The sulfone of mesityl oxide (XCVI) is formed in 41% yield by the action of chlorosulfonic acid-acetic anhydride (43) (Type 4). An equivalent yield is reported (105) by the use of sulfuric acid-acetic anhydride.



5. Seven-membered Rings

Mesityl oxide undergoes Type 3a condensations with four-membered diamines and aminothiols to give diazepines and thiazepines. These reactions are summarized in Table VII. As can be seen, dehydration of the intermediate cyclic alcohol occurs preferentially with the ring nitrogen, but can occur with the α -ethylenic carbon when the nitrogen is substituted.


TABLE VII
FORMATION OF DIAZEPINES AND THIAZEPINES

Reagent	X	R	Yield, %	Reference
NH ₂ CH ₂ CH ₂ NH ₂	NH	H	56	(107)
NH ₂ CH ₂ CH ₂ SH	S	H	68	(106)
NH ₂ CH ₂ CHSHCH ₃	S	CH ₃	41	(107)
C ₆ H ₅ NHCH ₂ CH ₂ SH	S	C ₆ H ₅	..	(107)


6. Polycyclic Compounds

A considerable amount of work has been devoted to the reaction of mesityl oxide with different mole ratios of phenol. The acid-catalyzed condensation of mesityl oxide with two or four moles of phenol (39) was originally claimed to give 2-(*p*-hydroxyphenyl)-2,4,4-trimethylchromane (XCVII). However, it has recently been shown (12) that this Type 3a reaction proceeds counter-wise and the product is 4-(*p*-hydroxyphenyl)-2,2,4-trimethylchromane (XCVIII). The equimolar, acid-catalyzed reaction of mesityl oxide and phenol (or substituted phenols such as *m*- and *p*-nitrophenol, the cresols, trinitro-*m*-cresol, 6-chloro-*m*-cresol, carvacrol and thymol) is described (110) as giving mixtures of the chromanol XCIX and the dichromanyl ether C, but this mode of condensation must also be open to some question.


A similar condensation of mesityl oxide and 4-hydroxycoumarin (77) gives about 13% of 4-methyl-4-(3-(4-hydroxy-2-chromenone))-2-pentanone (CI) and 32% of 2-hydroxy-2,4,4-trimethyl-3H,5H-pyrano(3,2c)-(1)benzopyran (CII).



The fusion of mesityl oxide and aniline using an iodine catalyst (91) and subsequent addition of acid yields 65% of 2,2,4-trimethyldihydroquinoline (CIII) by a Type 3a condensation. The acid-catalyzed reaction of mesityl oxide and *o*-phenylenediamine (44) was originally thought to give a 1,4-dihydroquinoxaline. This material has now been proved (122) to be 2-isobutenyl-2-methylbenzimidazoline (CIV), formed in a Type 1 reaction. This is confirmed by pyrolysis to the known 2-methylbenzimidazole. The over-all yield is 33%.




1,3-Dimethylindole also undergoes an acid-catalyzed fusion with mesityl oxide (130) to give 57% of the carbazole CV. This ring closure involves a Type 4 condensation.




This work has been extended to other simple mono- and dimethylindoles and to indole itself (111, 112). By use of equimolar quantities of mesityl oxide and the indole or excess mesityl oxide, the 4-methyl-4-(3-indolyl)-2-pentanone intermediates (CVI) are isolated. Isolation of CVI demonstrates an initial 1,4-addition in the reaction. These reactions are listed below.

With a molar excess of four parts of 2-methylindole to three parts of mesityl oxide, a 67% yield of the indolylcarbazole CVII is isolated. With a two-to-one excess of 1-methylindole over mesityl oxide, 36% of



R ₁	R ₂	Yield, %
H	H	6
CH ₃	H	28
H	CH ₃	70
CH ₃	CH ₃	5

1,1'-dimethyl-3,3'-isopropylidene-bis-indole (CVIII) is the product.



VII. REDUCTION, OXIDATION, AND PYROLYSIS

The literature contains a considerable body of material relating to these processes as applied to mesityl oxide. However, the discussion in this section will be limited to direct synthetic applications rather than a generalized survey of the area. The references cited have been chosen to give a combination of the best yields and procedures.

A. REDUCTION


Both the double bond and the carbonyl group of mesityl oxide are readily reduced. By proper choice of reducing agent and/or catalyst, reduction to the saturated ketone, alcohol or hydrocarbon or reduction to the unsaturated alcohol may be accomplished. A few typical procedures are summarized in Table VIII.

TABLE VIII
 REDUCTIONS OF MESITYL OXIDE


Product	Reducing agent	Catalyst	Yield, %	Reference
Me ₂ CHCH ₂ COMe	H ₂	Ni	95-100	(36)
	H ₂	Pd	94-97	(96)
Me ₂ CHCH ₂ CHOHMe	H ₂	Copper chromite	100	(3)
	H ₂	Raney Cu	95	(80)
Me ₂ CHCH ₂ CH ₂ CH ₃	H ₂	WS ₂ -(NH ₄) ₂ WS ₄	...	(98)
Me ₂ C=CHCHOHMe	H ₂	Al(iso-Pr) ₃	80	(79)
	NaBH ₄	...	77	(26)

A number of reductive condensations of mesityl oxide leading to cyclic compounds has also been reported. The reduction of mesityl oxide with aluminum amalgam (151) or electrolytically (93) gives 1,3,3,4,4-pentamethyl-5-acetylcyclopentanol (CIX). A direct yield of 60-70% of the dehydrated cyclopentene CX is obtained using sodium amalgam or aluminum amalgam (63). The postulated intermediate of this reaction CXI has not been isolated.

A second cyclization to the tetrahydrofuran CXII or its dehydration product CXIII can occur using




sodium amalgam (159), magnesium- or zinc-acetic acid (156), or electrolytic reduction (157). The intermediate CXIV has also been isolated using electrolytic techniques. The best reported yields are 55% for CXII and 35% for CXIII. By use of magnesium-acetic acid at -25 to -30° "good" yields of both CXII and CXIII are simultaneously obtained (92).



B. OXIDATION

The oxidation of mesityl oxide may proceed in a variety of ways depending upon the oxidizing agent and reaction conditions. The action of dilute permanganate in acetone (66) or hydrogen peroxide-osmium tetroxide (104) gives low yields of the glycol CXV. The only other oxidation of mesityl oxide proceeding without cleavage of the molecule is that with cobalt acetate-cupric acetate-acetic acid (45) which yields the acid CXVI.




A number of oxidative cleavages are listed in Table IX.


TABLE IX
OXIDATIONS OF MESITYL OXIDE

Reagent	Products	Yield, %	Reference
Dil. HNO_3	$\text{MeCOOH} + (\text{COOH})_2$...	(29)
Dil. KMnO_4	$\text{MeCOOH} + \text{Me}_2\text{COHCOOH}$...	(122)
2.5% KMnO_4	$\text{Me}_2\text{CO} + \text{MeCOCOHOH}$...	(143)
NaOCl	$\text{CHCl}_3 + \text{Me}_2\text{C}=\text{CHCOOH}$	75 (acid)	(14)
NaOBr	$\text{CHBr}_3 + \text{Me}_2\text{C}=\text{CHCOOH}$	44 (acid)	(94)
NaOI	$\text{CHI}_3 + \text{Me}_2\text{C}=\text{CHCOOH}$	75 (acid)	(37)


Epoxidation of mesityl oxide to give 3,4-epoxy-4-methyl-2-pentanone (CXVII) can be performed with *tert*-butyl peroxide-Triton B (163) (44% yield), *N*-bromosuccinimide (57) (56%) or hydrogen peroxide-sodium hydroxide-magnesium sulfate (162) (87%). When peroxyacetic acid (119) is used for epoxidation, only 20% of CXVII is obtained but this is accompanied by a 53% yield of the epoxy ester CXVIII.



In the presence of sulfuric acid (129), mesityl oxide and hydrogen peroxide afford 46% of bis-(3,5,5-trimethyl-1,2-dioxolanyl)-3-peroxide (CXIX). The sodium hydroxide-catalyzed reaction of mesityl oxide and hydrogen peroxide at 0 to -5° gives 45% of CXVII, but also 34% of 3-hydroxy-3,5,5-trimethyl-1,2-dioxolane (CXX). The dioxolane CXX on treatment with excess peroxide gives CXIX.



Treatment of mesityl oxide with ozone (62) gives an almost quantitative yield of the explosive, unstable ozonide CXXI. However, when the reaction is run as a heterogeneous system with water (60), the ozonide decomposes, *in situ*, into acetone and pyruvaldehyde (CXXII).



C. PYROLYSIS

The direct pyrolysis of mesityl oxide on activated alumina (100) or phosphoric acid-silica (103) gives acetic acid and isobutylene as major products. The reaction is, however, not clean-cut and a number of other materials are obtained.

In addition to this homo-pyrolysis, a number of mixed pyrolyses have been described and are briefly summarized in Table X.

TABLE X
PYROLYSES OF MESITYL OXIDE

Reagent	Catalyst	Products	Yield, %	Reference
$\text{CO}-\text{H}_2\text{O}$	H_3PO_4	$\text{MeCOOH} + \text{Me}_2\text{CCOOH}$..	(59)
Benzene	$\text{ZnCl}_2-\text{Al}_2\text{O}_3$	$\text{C}_6\text{H}_5\text{CMe}_3$	26	(136)
iso-PrOH	$\text{ZnO}-\text{Al}_2\text{O}_3$	Methylpentadienes	62	(13)
iso-PrOH	$\text{CrO}_3-\text{Al}_2\text{O}_3$	Methylpentadienes	75	(113)

VIII. REFERENCES

- (1) Abe, K., *Sci. Repts. Tokyo Kyoiku Daigaku Sect. A*, **3**, 17 (1935); *Chem. Zentr.*, **107**, I, 3691 (1936).
- (2) Adamson, D. W., and Kenner, J., *J. Chem. Soc.*, 1551 (1937).
- (3) Adkins, H., and Connor, R., *J. Am. Chem. Soc.*, **53**, 1091 (1931).
- (4) Anschütz, L., Klein, E., and Cermak, G., *Ber.*, **77**, 726 (1944).
- (5) Arbuzov, B. A., and Mukhamedova, L. A., *Izvest. Akad. Nauk S.S.S.R.*, 820 (1953); *Chem. Abstr.*, **49**, 963 (1955).
- (6) Arbuzov, B. A., and Mukhamedova, L. A., *Izvest. Akad. Nauk S.S.S.R.*, 448 (1954); *Chem. Abstr.*, **49**, 9568 (1955).
- (7) Arndt, F., Schwarz, R., Martius, C., and Aron, E., *Rev. Fac. sci. univ. Istanbul Ser. A*, **13**, 57 (1948); *Chem. Abstr.*, **42**, 4177 (1948).

- (8) von Auwers, K., and Kreuder, A., *Ber.*, **58**, 1980 (1925).
(9) Bacon, N., Brewis, S., Usher, G. E., and Waight, E. S., *J. Chem. Soc.*, 2255 (1961).
(10) Baeyer, A., *Ann.*, **140**, 298 (1866).
(11) Baker, W., *J. Chem. Soc.*, 1678 (1934).
(12) Baker, W., and McOmie, J. F. W., *Chemistry & Industry*, 256 (1955).
(13) Ballard, S. A., De V. Finch, H., and Peterson, E. A., U.S. Patent 2,693,497; *Chem. Abstr.*, **49**, 14793 (1955).
(14) Barbier, P., and Leser, G., *Bull. soc. chim. France*, (3) **33**, 815 (1905).
(15) Bergmann, E., and Bracha, P., *J. Org. Chem.*, **24**, 994 (1959).
(16) Bergmann, E., and Bracha, P., *J. Org. Chem.*, **26**, 4685 (1961).
(17) Borsche, W., *Ber.*, **42**, 4496 (1909).
(18) Boyer, J. H., *J. Am. Chem. Soc.*, **73**, 5248 (1951).
(19) Bradbury, R. B., Hancox, N. C., and Hatt, H. H., *J. Chem. Soc.*, 1397 (1947).
(20) Braude, E. A., Bruun, T., Weedon, B. C. L., and Woods, R. K., *J. Chem. Soc.*, 1425 (1952).
(21) Braude, E. A., and Evans, E. A., *J. Chem. Soc.*, 3331 (1955).
(22) Braude, E. A., Gofton, B. F., Lowe, G., and Waight, E. S., *J. Chem. Soc.*, 4057 (1956).
(23) Brown, R. W., Jones, W. E., and Pinder, A. R., *J. Chem. Soc.*, 2123 (1951).
(24) Bruson, H. A., U.S. Patent 2,395,453; *Chem. Abstr.*, **40**, 3467 (1946).
(25) Bruson, H. A., and Reiner, T. W., *J. Am. Chem. Soc.*, **64**, 2850 (1942).
(26) Chaikin, S. W., and Brown, W. G., *J. Am. Chem. Soc.*, **71**, 122 (1949).
(27) Chapurlat, R., and Dreux, J., *Compt. rend.*, **253**, 2361 (1961).
(28) Chelinysev, V. V., and Pataraya, A. V., *Zhur. Obschei Khim.*, **11**, 461 (1941); *Chem. Abstr.*, **35**, 6571 (1941).
(29) Claisen, L., *Ann.*, **180**, 6 (1876).
(30) Claisen, L., *Ann.*, **291**, 119 (1896).
(31) Claisen, L., and Claparede, A., *Ber.*, **14**, 439 (1881).
(32) Cologne, J., and Pichat, L., *Bull. soc. chim. France*, 421 (1949).
(33) Cologne, J., and Brisson, P., *Bull. soc. chim. France*, 98 (1962).
(34) Conia, J. M., *Bull. soc. chim. France*, 1392 (1956).
(35) Conia, J. M., and Naves, Y. R., *Compt. rend.*, **250**, 356 (1960).
(36) Covert, L. W., Connor, R., and Adkins, H., *J. Am. Chem. Soc.*, **54**, 1658 (1932).
(37) Cuculescu, V., *Bul. Fac. Stiinte Cernauti*, **1**, 53 (1927); *Chem. Zentr.*, **102**, I, 589 (1931).
(38) Curtius, T., and Wirsing, F., *J. prakt. Chem.*, [2] **50**, 546 (1894).
(39) Dianin, A., *Zhur. Russ. Fiz.-Khim. Obschestva*, **46**, 1310 (1914); *Chem. Zentr.*, **86**, I, 1063 (1915).
(40) Dippy, J. F. J., and Palluel, A. L. L., *J. Chem. Soc.*, 1418 (1951).
(41) Doevre, J., *Bull. soc. chim. France*, (4) **39**, 1594 (1926).
(42) Drake, L. R., and Marvel, C. S., *J. Org. Chem.*, **2**, 393 (1937-8).
(43) Eastman, R. H., and Gallup, D., *J. Am. Chem. Soc.*, **70**, 864 (1948).
(44) Ekely, J. B., and Wells, R. J., *Ber.*, **38**, 2259 (1905).
(45) Elce, A., Hall, R. H., and Tuerck, K. H. W., U.S. Patent 2,443,818; *Chem. Abstr.*, **42**, 7319 (1948).
(46) Elderfield, R. C., and McCarthy, J. R., *J. Am. Chem. Soc.*, **73**, 982 (1951).
(47) Esafov, V. I., *Zhur. Obschei Khim.*, **14**, 84 (1944); *Chem. Abstr.*, **39**, 905 (1945).
(48) Esafov, V. I., *Zhur. Obschei Khim.*, **17**, 1465 (1947); *Chem. Abstr.*, **43**, 110 (1949).
(49) Esafov, V. I., *Zhur. Obschei Khim.*, **19**, 1115 (1949); *Chem. Abstr.*, **44**, 5795 (1950).
(50) Farlow, M. W., and Signaigo, F. K., U.S. Patent 2,402,613; *Chem. Abstr.*, **40**, 5759 (1946).
(51) von Fellenberg, T., *Ber.*, **37**, 3578 (1904).
(52) von Fellenberg, T., *Ber.*, **39**, 2064 (1906).
(53) Foldi, Z., and Kollonitsch, J., *J. Chem. Soc.*, 1683 (1948).
(54) France, H., Maitland, P., and Tucker, S. H., *J. Chem. Soc.*, 1741 (1937).
(55) Gault, H., and Laloi, L., *Bull. soc. chim. France*, 1079 (1952).
(56) Grignard, V., *Compt. rend.*, **130**, 1322 (1900).
(57) Guss, C. O., and Rosenthal, R., *J. Am. Chem. Soc.*, **77**, 2549 (1955).
(58) Gwynn, B. H., and Degering, E. F., *J. Am. Chem. Soc.*, **64**, 2216 (1942).
(59) Hardy, D. V. N., *J. Chem. Soc.*, 464 (1938).
(60) Harries, C., *Ber.*, **36**, 1933 (1903).
(61) Harries, C., *Ann.*, **330**, 188 (1904).
(62) Harries, C., *Ann.*, **374**, 340 (1910).
(63) Harries, C., and Hubner, F., *Ann.*, **296**, 308 (1897).
(64) Harries, C., and Jablonski, L., *Ber.*, **31**, 1377 (1898).
(65) Harries, C., and Kaiser, F., *Ber.*, **32**, 1338 (1899).
(66) Harries, C., and Pappos, *Ber.*, **34**, 2979 (1901).
(67) Haurry, V. E., British Patent 578,635; *Chem. Abstr.*, **41**, 3480 (1947).
(68) Henze, H. R., Thompson, T. R., and Speer, R. J., *J. Org. Chem.*, **8**, 18 (1943).
(69) Heilbron, I. M., Johnson, A. W., Jones, E. R. H., and Spinks, A., *J. Chem. Soc.*, 732 (1942).
(70) Hill, W. H., British Patent 633,353; *Chem. Abstr.*, **44**, 5924 (1950).
(71) Hochstetter, A., and Kohn, M., *Montash.*, **24**, 773 (1903).
(72) Hoegberg, E. I., U.S. Patent 2,632,020; *Chem. Abstr.*, **48**, 2759 (1954).
(73) Hoffman, A., *J. Am. Chem. Soc.*, **49**, 530 (1927).
(74) Hoffman, A., *J. Am. Chem. Soc.*, **51**, 2542 (1929).
(75) Hogsed, M. J., and Lindsey, R. V., *J. Am. Chem. Soc.*, **76**, 2305 (1954).
(76) House, H. O., Grubbs, E. J., and Gannon, W. F., *J. Am. Chem. Soc.*, **82**, 4099 (1960).
(77) Ikawa, M., Stahmann, M. A., and Link, K. P., *J. Am. Chem. Soc.*, **66**, 902 (1944).
(78) Ionescu, V., and Popescu, O. G., *Bull. soc. chim. France*, [4] **51**, 1224 (1932).
(79) Jackman, L. M., MacBeth, A. K., and Mills, J. A., *J. Chem. Soc.*, 2646 (1949).
(80) Jadot, J., and Braine, R., *Bull. soc. roy. sci. Liege*, **25**, 62 (1956); *Chem. Abstr.*, **50**, 16653 (1956).
(81) Jansen, J. E., U.S. Patent 2,568,633; *Chem. Abstr.*, **46**, 3574 (1952).
(82) Japp, F. R., and Knox, J., *J. Chem. Soc.*, 674 (1905).
(83) Jessafow, W. I., *Zhur. Obschei Khim.*, **9**, 467 (1939); *Chem. Zentr.*, **111**, I, 527 (1940).
(84) Jessafow, W. I., and Smirnow, M. W., *Zhur. Obschei Khim.*, **10**, 1535 (1940); *Chem. Zentr.*, **112**, II, 600 (1941).
(85) Joos, B., U.S. Patent 2,554,947; *Chem. Abstr.*, **46**, 3069 (1952).
(86) Kerp, W., and Muller, F., *Ann.*, **299**, 217 (1898).
(87) Klages, A., *Ber.*, **37**, 2305 (1904).
(88) Kloetzel, M. C., *J. Am. Chem. Soc.*, **69**, 2271 (1947).
(89) Knoevenagel, E., *Ann.*, **297**, 185 (1897).

- (90) Knoevenagel, E., German Patent 162,281; *Chem. Zentr.*, **76**, I, 726 (1905).
- (91) Knoevenagel, E., *Ber.*, **56**, 2414 (1923).
- (92) Kolobielski, M., *Ann. chim.* (Paris), **10**, 271 (1955); *Chem. Abstr.*, **50**, 7768 (1956).
- (93) Kolobielski, M., and Wiemann, J., *Compt. rend.*, **238**, 1039 (1954).
- (94) Kologsivova, N. E., and Belov, V. N., *Zhur. Obshechi Khim.*, **28**, 1269 (1958); *Chem. Abstr.*, **52**, 19929 (1958).
- (95) Komppa, G., *Ber.*, **32**, 1421 (1899).
- (96) Lacey, R. N., British Patent 748,239; *Chem. Abstr.*, **51**, 460 (1957).
- (97) Lambiotte and Co., French Patent 972,652; *Chem. Abstr.*, **47**, 1744 (1953).
- (98) Landa, S., and Mostecky, J., *Freiberger Forschungsh.*, **A36**, 58 (1955); *Chem. Abstr.*, **50**, 10666 (1956).
- (99) Lapworth, A., *J. Chem. Soc.*, 1214 (1904).
- (100) Larin, A. Y., and Frost, A. V., *Doklady Akad. Nauk S.S.S.R.*, **59**, 1297 (1948); *Chem. Abstr.*, **42**, 7244 (1948).
- (101) Lorette, N. B., *J. Org. Chem.*, **23**, 973 (1958).
- (102) Mayuranthan, P. S., *J. Indian Inst. Sci. Sect. A*, **16**, 123 (1933); *Chem. Zentr.*, **105**, II, 240 (1934).
- (103) McAllister, S. H., Bailey W. A., and Bouton, C. M., *J. Am. Chem. Soc.*, **62**, 3210 (1940).
- (104) Milas, N. A., Sussman, S., and Mason, H. S., *J. Am. Chem. Soc.*, **61**, 1845 (1939).
- (105) Morel, T., and Verkade, P. E., *Rec. trav. chim.*, **68**, 626 (1949).
- (106) Mushkalo, L. K., and Lanovaya, Z. I., *Ukrain. Khim. Zhur.*, **21**, 631 (1955); *Chem. Abstr.*, **50**, 14783 (1956).
- (107) Mushkalo, L. K., and Shokol, Z. I., *Zhur. Obshechi Khim.*, **30**, 1023 (1960).
- (108) Naves, Y. R., and Conia, J. M., *Compt. rend.*, **251**, 1130 (1960).
- (109) Nazarov, I. N., and Kakhniashvili, A. I., *Sbornik Statei Obshechi Khim.*, **2**, 919 (1954); *Chem. Abstr.*, **49**, 6848 (1955).
- (110) Niederl, J. B., *J. Am. Chem. Soc.*, **51**, 2426 (1929).
- (111) Noland, W. E., Venkiteswaran, M. R., and Richards, C. G., *J. Org. Chem.*, **26**, 4241 (1961).
- (112) Noland, W. E., Richards, C. G., Desai, H. S., and Venkiteswaran, M. R., *J. Org. Chem.*, **26**, 4254 (1961).
- (113) N. V. de Bataafsche Petroleum Maatschappij, Dutch Patent 70,917; *Chem. Abstr.*, **48**, 1405 (1954).
- (114) Oroshnik, W., and McBane, R. D., *J. Am. Chem. Soc.*, **71**, 2062 (1949).
- (115) Pastureau, H., and Bernard, H., *Compt. rend.*, **174**, 1555 (1922).
- (116) Patrick, T. M., *J. Org. Chem.*, **17**, 1269 (1952).
- (117) Pauly, H., and Lieck, H., *Ber.*, **33**, 500 (1900).
- (118) Pawlow, D., *Ann.*, **188**, 138 (1877).
- (119) Payne, G. B., and Williams, P. H., *J. Org. Chem.*, **24**, 284 (1959).
- (120) Perveev, F. Y., and Kudryashova, N. I., *Zhur. Obshechi Khim.*, **22**, 1580 (1952); *Chem. Abstr.*, **47**, 9251 (1953).
- (121) Philippi, E., Moser, H., and Moser, H., *Ber.*, **68**, 1810 (1935).
- (122) Pinner, A., *Ber.*, **15**, 591 (1882).
- (123) Posner, T., *Ber.*, **34**, 1398 (1901).
- (124) Pudovik, A. N., *Zhur. Obshechi Khim.*, **22**, 1371 (1952); *Chem. Abstr.*, **47**, 4837 (1953).
- (125) Pudovik, A. N., Sabirova, R. D., and Tener, T. A., *Zhur. Obshechi Khim.*, **24**, 1026 (1954); *Chem. Abstr.*, **49**, 8790 (1955).
- (126) Putezer, B., Nield, C. H., and Barry, R. H., *J. Am. Chem. Soc.*, **67**, 835 (1945).
- (127) Qudrat-I-Khuda, M., *J. Chem. Soc.*, 205 (1929).
- (128) Richard, G., Mirjolet, M., and Gschwind, P., *Compt. rend.*, **223**, 1007 (1946).
- (129) Rieche, A., Schmitz, E., and Grundeman, E., *Ber.*, **93**, 2443 (1960).
- (130) Robinson, B., and Smith, G. F., *J. Chem. Soc.*, 4574 (1960).
- (131) Rupe, H., and Kessler, S., *Ber.*, **42**, 4715 (1909).
- (132) Salmi, E. J., and Rannikko, V., *Ber.*, **72**, 600 (1939).
- (133) Scheiber, J., and Meisel, F., *Ber.*, **48**, 261 (1915).
- (134) Scheiber, J., and Wolf, H., *Ann.*, **357**, 25 (1902).
- (135) Scheuer, P. J., Bothelo, H. C., and Pauling, C., *J. Org. Chem.*, **22**, 674 (1957).
- (136) Schmerling, L., and Ipatieff, V. N., U.S. Patent 2,410,553; *Chem. Abstr.*, **41**, 1242 (1947).
- (137) Schmitt, J., *Ann.*, **569**, 36 (1950).
- (138) Schneider, W., and Sack, A., *Ber.*, **56**, 1786 (1923).
- (139) Slawinski, K., *Chemik Polski*, **15**, 106 (1918); *Chem. Zentr.*, **89**, I, 9 (1918).
- (140) Smith, L. I., and Prichard, W. W., *J. Am. Chem. Soc.*, **62**, 776 (1940).
- (141) Staudinger, H., Muntwyler, O., Ruzicka, L., and Seibt, S., *Helv. Chim. Acta*, **7**, 398 (1924).
- (142) Stross, F. H., Monger, J. M., and de V. Finch, H., *J. Am. Chem. Soc.*, **69**, 1627 (1947).
- (143) Suida, H., and Poll, H., *Monatsh.*, **48**, 182 (1927).
- (144) Tindall, J. B., U.S. Patent 2,430,436; *Chem. Abstr.*, **42**, 1964 (1948).
- (145) Tischtschenko, D. W., *Zhur. Obshechi Khim.*, **11**, 402 (1941); *Chem. Zentr.*, **113**, I, 1739 (1942).
- (146) Traube, W., and Schwarz, R., *Ber.*, **32**, 3163 (1899).
- (147) Tucker, S. H., and Whalley, M., *J. Chem. Soc.*, 53 (1949).
- (148) Utzinger, G. E., U.S. Patent 2,677,705; *Chem. Abstr.*, **49**, 6303 (1955).
- (149) Utzinger, G. E., and Hoelle, A., *Helv. Chim. Acta*, **35**, 1375 (1952).
- (150) Vinton, W. H., U.S. Patent 2,427,582; *Chem. Abstr.*, **42**, 212 (1948).
- (151) Vogel, I., *J. Chem. Soc.*, 594 (1927).
- (152) Vorlander, D., *Ann.*, **294**, 300 (1875).
- (153) Wetterholm, G. A., German Patent 934,694; *Chem. Abstr.*, **53**, 17513 (1959).
- (154) Whitmore, F. C., Goldsmith, D. P. J., Cook, N. C., Yarze, J. C., and Ecke, G. G., *J. Am. Chem. Soc.*, **72**, 53 (1950).
- (155) Weimann, J., and Dupayrat, J., *Bull. soc. chim. France*, 209 (1961).
- (156) Wiemann, J., and Glacet, C., *Compt. rend.*, **226**, 923 (1948).
- (157) Wiemann, J., and Maitte, P., *Bull. soc. chim. France*, 430 (1952).
- (158) Weimann, J., Martineau, L., and Tiquet, J., *Compt. rend.*, **241**, 807 (1955).
- (159) Weimann, J., and Nahum, R., *Compt. rend.*, **238**, 2091 (1954).
- (160) Wiemann, J., Thuan, L-T., and Rame, D., *Compt. rend.*, **249**, 1529 (1959).
- (161) Wiemann, J., Thuan, L-T., and Sanderer, F., *Bull. soc. chim. France*, 697 (1957).
- (162) Wilder, R., Park, E., and Dolnick, A. A., U.S. Patent 2,431,718; *Chem. Abstr.*, **42**, 3430 (1948).
- (163) Yang, N. C., and Finnegan, R. A., *J. Am. Chem.*, **80**, 5845 (1958).
- (164) Young, F. G., *J. Am. Chem. Soc.*, **71**, 1346 (1949).
- (165) Yur'ev, Y. K., Zeifrov, N. S., Shteinman, A. A., and Gurevich, V. M., *Zhur. Obshechi Khim.*, **30**, 411 (1960).