

# THE CHEMISTRY OF CHLOROFORMATES

MARKUS MATZNER, RAYMOND P. KURKJY, AND ROBERT J. COTTER

Research and Development Department, Union Carbide Corporation, Plastics Division, Bound Brook, New Jersey

Received June 22, 1964


## CONTENTS

I. Introduction.....	645
II. Methods of Synthesis of Chloroformates.....	645
A. Aliphatic Chloroformates.....	645
B. Aromatic Chloroformates.....	649
III. Physical Properties of Chloroformates.....	649
A. Structure.....	649
B. Spectral Studies.....	650
IV. Chemical Properties of Chloroformates.....	651
A. Reactions of Chloroformates with Hydroxylic Compounds.....	651
B. Reaction of Chloroformates with Carboxylic Acids.....	654
C. Reactions of Carboxylic-Carbonic Anhydrides .....	655
D. Reactions of Chloroformates with Nitrogen Compounds.....	656
E. Miscellaneous Reactions and Syntheses.....	666
V. Polymer Syntheses from Chloroformates.....	674
A. Polycarbonates.....	674
B. Polyesters.....	675
C. Polyurethans .....	675
D. Copolyurethans .....	676
E. Polymerization of Chloroformates with Diepoxides.....	676
F. Polymerization of Chloroformates with Bis(hydrazides) .....	676
G. Uses of Chloroformates in Addition-Type Polymers .....	676
H. Uses of Chloroformates in Thermoset Polymers.....	676
I. Miscellaneous Uses of Chloroformates in Polymeric Systems.....	677
VI. References.....	677

## I. INTRODUCTION

The chemistry of chloroformates has not been reviewed in depth previously, but has only received brief treatment in conjunction with reviews on phosgene. Dyson's review on phosgene which appeared in 1927 included four pages on the preparation and properties of chloroformates (193). A review on phosgene and its derivatives which appeared in 1938 by Criscuoli dealt with some of the lower alkyl and chloroalkyl chloroformates, particularly with their use as war gases (160). The chapter on carbonic acid derivatives by Petersen and Pieperbrink in Houben-Weyl devoted five pages to the preparation and properties of chloroformates (334).

Chloroformates have also been referred to in the literature as chlorocarbonates. The formulas shown below indicate the interrelationship.



Throughout this review, the name of chloroformates has been used in order to be consistent with *Chemical Abstracts'* current practice of indexing these compounds under "formic acid, chloro-, esters of."


In view of the voluminous, published literature relating to chloroformates, the authors do not claim that this review contains every reference pertaining to this

field. However, it can be stated that complete coverage through 1962 was our goal. In those instances where we have failed to attain this goal, it is our hope that we have included at least one example of the known uses and chemistry of chloroformates.

## II. METHODS OF SYNTHESIS OF CHLOROFORMATES

### A. ALIPHATIC CHLOROFORMATES

Phosgene reacts with aliphatic hydroxy compounds at room temperature or below to yield chloroformates in high yield (334). Low temperatures are preferred in many instances to minimize formation of dialkyl carbonates.



This synthesis can be carried out advantageously in the presence of a tertiary amine (51) as the hydrogen chloride acceptor, especially with tertiary alcohols. Removal of the hydrogen chloride in this way minimized deleterious acid-catalyzed side reactions in cases where acid-sensitive groups were present, and also increased the rate of reaction. As can be seen from Table I, chloroformates with the following additional functional groups have been synthesized: halogen (fluorine, chlorine, bromine), cyano, nitro, hydroxyl, amino, imino, amido, N-containing heterocycles, ether, ester

TABLE I  
ALIPHATIC CHLOROFORMATES  
1. Alkyl and Alicyclic Monochloroformates, ROCOCl

R	Method <sup>a</sup>	Yield, %	Reference
C <sub>1</sub>			
Cl <sub>3</sub> C			53, 160, 173, 267, 269, 306, 308, 327, 389
			391, 745
Cl <sub>2</sub> CH			160, 173, 268, 269, 308, 327, 389, 391, 745
ClCH <sub>2</sub>			41, 53, 160, 173, 267, 269, 306-309, 327,
			389, 391, 745
CH <sub>3</sub>			53, 98, 104, 140, 187, 189, 305, 307, 325,
			388, 392, 409, 459, 521, 594, 602
C <sub>2</sub>			
Cl <sub>4</sub> CCCl <sub>2</sub>	C		10, 140, 141, 489
Cl <sub>3</sub> CHCCl <sub>2</sub>	D		489
Cl <sub>2</sub> CCHCl			50
Br <sub>2</sub> CCH <sub>2</sub>	A		101
ClCH <sub>2</sub> CCl <sub>2</sub>	C		498
NCCCH <sub>2</sub>	B		300
CH <sub>2</sub> =CH	E		417
ClCH <sub>2</sub> CHCl	C		489
Cl(CH <sub>2</sub> ) <sub>2</sub>			154, 155, 172, 212, 360, 447, 453, 507, 519,
			521, 626
CH <sub>3</sub> CHCl	C		489
F(CH <sub>2</sub> ) <sub>2</sub>	A	81	396, 532
O <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>	A		88
CH <sub>3</sub> CH <sub>2</sub>	B		43, 45, 46, 98, 104, 140, 183, 187, 289, 317,
			404, 405, 464, 543, 594, 602, 659
C <sub>3</sub>			
CH≡CCH <sub>2</sub>	A	50	231
CH <sub>2</sub> =CCICH <sub>2</sub>			505
ClCH=CHCH <sub>2</sub>	A	50	130
(ClCH <sub>2</sub> )(Cl <sub>2</sub> CH)CH	C	48	446
CH <sub>2</sub> =CHCH <sub>2</sub>			85, 227, 491, 493, 495, 500, 505, 630, 676
CH <sub>2</sub> =C(CH <sub>3</sub> )	D		451
(ClCH <sub>2</sub> ) <sub>2</sub> CH	C	54	446, 536
ClCH <sub>2</sub> CHClCH <sub>2</sub>	A		536
Cl(CH <sub>2</sub> ) <sub>3</sub>	A	>80	185, 557
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>2</sub>			43, 287, 593, 594, 602
(CH <sub>3</sub> ) <sub>2</sub> CH			287, 301, 325, 404, 405, 451, 460, 461, 491,
			602, 647, 672, 676
CH <sub>3</sub> O(CH <sub>2</sub> ) <sub>2</sub>	A		199
C <sub>4</sub>			
ClCH <sub>2</sub> C≡CCH <sub>2</sub>	A	70	330
CH <sub>2</sub> =C=CHCH <sub>2</sub>	A	90	105, 106
OCOOCH <sub>2</sub> CHCH <sub>2</sub>			658
CH <sub>2</sub> =CHCH(CH <sub>2</sub> Cl)	A	67	65
CH <sub>2</sub> CH=CHCH <sub>2</sub>	A		491, 495
CH <sub>2</sub> =C(CH <sub>3</sub> )CH <sub>2</sub>	A		491, 493, 495, 505
CH <sub>2</sub> OCH <sub>2</sub> CH <sub>2</sub> CH	A	93.3	544, 578
Cl(CH <sub>2</sub> ) <sub>4</sub>	A		298
CH <sub>3</sub> CH <sub>2</sub> CH(NO <sub>2</sub> )CH <sub>2</sub>	A		66
O <sub>2</sub> NC(CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub>	A		66
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>3</sub>			120, 351, 404, 405, 594
(CH <sub>3</sub> ) <sub>2</sub> CHCH <sub>2</sub>			287, 317, 404, 405, 594
(CH <sub>3</sub> ) <sub>3</sub> C	B	20.4	57, 133
CH <sub>3</sub> CH <sub>2</sub> CH(CH <sub>3</sub> )			6, 48, 367, 404, 405, 695
C <sub>5</sub>			
CH <sub>2</sub> =CHCOO(CH <sub>2</sub> ) <sub>2</sub>	A		291
CH <sub>2</sub> =CHCH <sub>2</sub> OCOCH <sub>2</sub>	A	60	341
CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH	B	61	512
OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CHCH <sub>2</sub>	A	87	589
CH <sub>3</sub> CH(COOC <sub>2</sub> H <sub>5</sub> )	A		248, 676
Cl(CH <sub>2</sub> ) <sub>5</sub>	A		298
Cl(CH <sub>2</sub> ) <sub>2</sub> CH(C <sub>2</sub> H <sub>5</sub> )			547
ClCH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub>			33
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>4</sub>	A	80	6, 602
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>2</sub> CH(CH <sub>3</sub> )	B		43, 48
(CH <sub>3</sub> ) <sub>2</sub> CH(CH <sub>2</sub> ) <sub>2</sub>	B		404, 405, 456, 594
C <sub>2</sub> H <sub>5</sub> C(CH <sub>3</sub> ) <sub>2</sub>	B		465
(CH <sub>3</sub> ) <sub>3</sub> CCH <sub>2</sub>	A	83	660
(CH <sub>3</sub> ) <sub>2</sub> CHCH(CH <sub>3</sub> )			48
(C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub> CH			48
H <sub>2</sub> N(CH <sub>2</sub> ) <sub>6</sub>			344

TABLE I (Continued)

R	Method <sup>a</sup>	Yield, %	Reference
C <sub>6</sub> CH <sub>2</sub> =C(CH <sub>3</sub> )COO(CH <sub>2</sub> ) <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH	A A	291 82	351, 404, 405, 594
Cl(CH <sub>2</sub> ) <sub>6</sub>	A	298	
Cl(CH <sub>2</sub> ) <sub>2</sub> CH(C <sub>3</sub> H <sub>7</sub> )	A	547	
C <sub>4</sub> H <sub>9</sub> N=C(CH <sub>3</sub> )	A	95	299
(CH <sub>3</sub> ) <sub>2</sub> CHCH <sub>2</sub> CH(CH <sub>3</sub> )	A	71	695
CH <sub>3</sub> CH <sub>2</sub> CH(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub>	A	56	695
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>2</sub> CH(CH <sub>3</sub> )		48	
(CH <sub>3</sub> ) <sub>2</sub> CHCH(C <sub>2</sub> H <sub>5</sub> )		48	
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>2</sub> CH(C <sub>2</sub> H <sub>5</sub> )		48	
(C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub> ·HCl	A	89	
C <sub>7</sub> C <sub>6</sub> H <sub>5</sub> CHCl	D	50	
4-(O <sub>2</sub> N)C <sub>6</sub> H <sub>4</sub> CH <sub>2</sub>	A	95	107
C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub>	A	91	45, 46, 58, 80, 113, 223, 404, 405, 676
2-(CH <sub>3</sub> )C <sub>6</sub> H <sub>10</sub>	B	404, 405	
C <sub>6</sub> H <sub>5</sub> CH(CH <sub>3</sub> )CH(C <sub>2</sub> H <sub>5</sub> )		49	
(n-C <sub>3</sub> H <sub>7</sub> ) <sub>2</sub> CH		48	
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>2</sub> CH(C <sub>2</sub> H <sub>5</sub> )	A	695	
i-C <sub>3</sub> H <sub>7</sub> CH <sub>2</sub> CH(C <sub>2</sub> H <sub>5</sub> )		49	
(i-C <sub>3</sub> H <sub>7</sub> ) <sub>2</sub> CH	A	669	
C <sub>8</sub> CF <sub>3</sub> (CF <sub>2</sub> ) <sub>6</sub> CH <sub>2</sub>	B	520	
C <sub>6</sub> H <sub>5</sub> CH(CH <sub>3</sub> )	B	338	
Cl(CH <sub>2</sub> ) <sub>8</sub>	A	298	
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>6</sub> CH(CH <sub>3</sub> )	B	335, 337, 339	
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>7</sub>	A	110, 404, 405, 602	
C <sub>9</sub> 7-(β-Hydroxyethyl)theophylline	B	96	
3,3,5-Trimethylcyclohexyl	A	81	695
Cl(CH <sub>2</sub> ) <sub>9</sub>	A	298	
(i-C <sub>4</sub> H <sub>9</sub> ) <sub>2</sub> CH	A	31	695
(t-C <sub>4</sub> H <sub>9</sub> ) <sub>2</sub> CH	B	93	379
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>8</sub>	B	87	345
C <sub>10</sub> C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> CONH(CH <sub>2</sub> ) <sub>2</sub>	A	55	
C <sub>6</sub> H <sub>5</sub> CH(i-C <sub>4</sub> H <sub>7</sub> )	B	662, 663	
l-Menthyl	B	404, 405	
Cl(CH <sub>2</sub> ) <sub>10</sub>	A	298	
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>9</sub>	A	508	
C <sub>11</sub> CF <sub>2</sub> Cl(CF <sub>2</sub> ) <sub>6</sub> CH <sub>2</sub>	B	520	
CHF <sub>2</sub> (CF <sub>2</sub> ) <sub>6</sub> CH <sub>2</sub>	B	520	
C <sub>12</sub> Diacetone glucosyl	B	50-60	237
C <sub>13</sub> CHF <sub>2</sub> (CF <sub>2</sub> ) <sub>11</sub> CH <sub>2</sub>	B	520	
C <sub>2</sub> H <sub>5</sub> OCOC(CH <sub>2</sub> ) <sub>2</sub> CH(C <sub>6</sub> H <sub>5</sub> )	B	8	
C <sub>15</sub> (C <sub>6</sub> H <sub>5</sub> ) <sub>2</sub> C(CH <sub>2</sub> OH)CH <sub>2</sub>	B	56	
C <sub>n</sub> Complex unsaturated			492, 500, 501, 503
Cl(CH <sub>2</sub> ) <sub>n</sub>		1	
Various β-alkoxyethyl radicals	A	70-93	20
-(CH <sub>2</sub> CH <sub>2</sub> O) <sub>n</sub> (CH <sub>2</sub> ) <sub>2</sub> (CH <sub>2</sub> CH <sub>2</sub> O) <sub>n</sub> (CH <sub>2</sub> ) <sub>2</sub> -		127	
2. Alkyl and Alicyclic Polychloroformates: R(OCOCl) <sub>n</sub>			
C <sub>2</sub> -(CH <sub>2</sub> ) <sub>2</sub> -	B	>75	354, 373, 397, 398, 411, 529, 565, 648, 657
C <sub>3</sub> -(CH <sub>2</sub> ) <sub>3</sub> -	B	>75	354, 373, 397, 439, 529, 657
-CH <sub>2</sub> CH(CH <sub>3</sub> )-	A	65	657
C <sub>4</sub> -CH <sub>2</sub> C≡CCH <sub>2</sub> -	B		275, 566
-CH <sub>2</sub> C(Cl)=C(Cl)CH <sub>2</sub> -	C	96.5	275
-CH <sub>2</sub> CCl <sub>2</sub> CCl <sub>2</sub> CH <sub>2</sub> -	C	~95	275
-(CH <sub>2</sub> ) <sub>4</sub> -	B	80	354, 397, 411, 565
-CH <sub>2</sub> CH(CH <sub>3</sub> )CH <sub>2</sub> -	A		566
-(CH <sub>2</sub> ) <sub>2</sub> CH(CH <sub>3</sub> )-	A		566
-(CH <sub>2</sub> ) <sub>2</sub> O(CH <sub>2</sub> ) <sub>2</sub> -			297
-(CH <sub>2</sub> ) <sub>2</sub> S(CH <sub>2</sub> ) <sub>2</sub> -	A		380
C <sub>5</sub> -(CH <sub>2</sub> ) <sub>5</sub> -	B	75	354
-CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub> -	A	90	246, 373, 439, 566



TABLE I (Continued)

R	Method <sup>a</sup>	Yield, %	Reference
C <sub>6</sub>			
1,4-Cyclohexylene	A	~96	527, 566, 727
1,4-Cyclohexylene ( <i>cis</i> )	B		354
1,4-Cyclohexylene ( <i>trans</i> )	B	85	354
-(CH <sub>2</sub> ) <sub>2</sub> CHCH <sub>2</sub> CH(CH <sub>3</sub> )-	A		411
(			
-CH <sub>2</sub> C(CH <sub>3</sub> )(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> -	B		439
-(CH <sub>2</sub> ) <sub>6</sub> -	A	88	332, 354
-CH <sub>2</sub> CH(CH <sub>3</sub> )OCH <sub>2</sub> CH(CH <sub>3</sub> )-	A		494, 561
-(CH <sub>2</sub> ) <sub>2</sub> O(CH <sub>2</sub> ) <sub>2</sub> O(CH <sub>2</sub> ) <sub>2</sub> -	A		494, 561
C <sub>7</sub>			
-(CH <sub>2</sub> ) <sub>7</sub> -	A	90	354
-(CH <sub>2</sub> ) <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> -	A		566
-(CH <sub>3</sub> )CHC(CH <sub>3</sub> ) <sub>2</sub> CH(CH <sub>3</sub> )-	B		128
-CH <sub>2</sub> C(i-C <sub>3</sub> H <sub>7</sub> )(CH <sub>3</sub> )CH <sub>2</sub> -	B		57, 439
-CH <sub>2</sub> C(C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub> CH <sub>2</sub> -	B	75	439
-CH <sub>2</sub> C(CH <sub>3</sub> )(n-C <sub>8</sub> H <sub>7</sub> )CH <sub>2</sub> -	B		57, 439
C <sub>8</sub>			
1,1-Cyclohexylidene, dimethyl	A		373
1,4-Cyclohexylene, dimethyl			398
2,2,4,4-Tetramethyl-1,3-cyclobutylene	A		100
-(CH <sub>2</sub> ) <sub>8</sub> -	A		354
C <sub>9</sub>			
-(CH <sub>2</sub> ) <sub>9</sub> -	A	80	354
-CH <sub>2</sub> C(n-C <sub>8</sub> H <sub>7</sub> ) <sub>2</sub> CH <sub>2</sub> -	B		439
-CH <sub>2</sub> C(C <sub>2</sub> H <sub>5</sub> )(n-C <sub>4</sub> H <sub>9</sub> )CH <sub>2</sub> -	B		439
C <sub>10</sub>			
-CH <sub>2</sub> C(CH <sub>3</sub> )(C <sub>6</sub> H <sub>5</sub> )CH <sub>2</sub> -	B		57, 439
-(CH <sub>2</sub> ) <sub>10</sub> -	B	85	354, 468, 724
C <sub>11</sub>			
-CH <sub>2</sub> C(C <sub>2</sub> H <sub>5</sub> )(C <sub>6</sub> H <sub>5</sub> )CH <sub>2</sub> -	B		439
C <sub>12</sub>			
1,4-[-(CH <sub>2</sub> ) <sub>2</sub> OCO] <sub>2</sub> C <sub>6</sub> H <sub>4</sub> -	A		468
C <sub>14</sub>			
-(CH <sub>2</sub> ) <sub>14</sub> -	A		468
C <sub>15</sub>			
(CH <sub>3</sub> ) <sub>2</sub> C(4-C <sub>6</sub> H <sub>10</sub> ) <sub>2</sub>			727
C <sub>18</sub>			
-(CH <sub>2</sub> ) <sub>18</sub> -	A		468
C <sub>n</sub>			
-(CH <sub>2</sub> CH <sub>2</sub> O) <sub>n</sub> -	A		468
-[(CH <sub>2</sub> ) <sub>4</sub> O] <sub>n</sub> -	A		468
-(CH <sub>2</sub> ) <sub>2</sub> [OCO(CH <sub>2</sub> ) <sub>4</sub> COO(CH <sub>2</sub> ) <sub>2</sub> ] <sub>n</sub> -	A		468
-(CH <sub>2</sub> ) <sub>2</sub> [OCONH(1,3-C <sub>6</sub> H <sub>4</sub> )NHCOO(CH <sub>2</sub> ) <sub>2</sub> ] <sub>n</sub> -	A		468
Poly(ether-urethanes)	A		468, 654


<sup>a</sup> A, alcohol and phosgene, no catalyst; B, alcohol and phosgene, added base; C, photolytic chlorination; D, carbonyl compound and phosgene; E, pyrolysis of dichloroformate.

(also cyclic carbonates), sulfide, and carbon-to-carbon unsaturation of various types. Chloroformates of partially protected sugar molecules have been prepared in 60% yield (237, 707). Conversion of the hydroxyl groups of cholesterol (393, 413, 711), deoxycholic acid (711), methyl and ethyl cholate (433, 711), and follicular hormones (180) to chloroformates has been reported. Bis- and polychloroformates have also been prepared.

Reaction of aliphatic hydroxy compounds with di(I) and triphosgenes (II) yielded alkyl chloroformates (98, 140). This method has not been used extensively




and possessed no advantage over direct reaction of the hydroxy compound with phosgene.



Formates have been chlorinated to yield chloroform-


ates, but this synthesis is not very useful since all of the possible chlorinated chloroformates are usually produced in varying amounts (10, 53, 140, 141, 267, 269, 307, 389, 391, 745). In this reaction, methyl formate yielded a complex mixture of all the possible chlorinated chloroformates.



Alkyl chloroformates have been chlorinated to yield the same type of product mixture (10, 41, 53, 86, 140, 141, 173, 267, 269, 306, 308, 389, 391, 446, 489).


Chlorination of an acetylenic bis(chloroformate) has been effected without destruction of the dichloroformate groups (275).

$\beta$ -Chlorochloroformates can be readily prepared in 50–100% yields from the reaction of epoxides with phosgene (360, 447).




This reaction, though it proceeds uncatalyzed, is more efficient when catalyzed with either acids or bases.

$\alpha$ -Chlorochloroformates have been prepared in some cases by reaction of phosgene with carbonyl compounds, especially chloral (50).




Where enolizable carbonyl groups are present, e.g., acetone, unsaturated chloroformates were obtained (451).




The reaction of phosgene with carbonyl compounds has not been extensively investigated.


Alkyl chloroformates have been prepared by reaction of dialkyl and cyclic carbonates with phosphorus pentachloride (272, 594).



Reaction of diethyl pyrocarbonate with thionyl chloride has yielded ethyl chloroformate (543).




Chlorination of xanthates and thionoesters has yielded alkyl chloroformates (183).




Even though these routes may be general, they offer no advantage for the preparation of alkyl chloroformates.

An interesting example of "chloroformate ester exchange" has also been reported (382).




### B. AROMATIC CHLOROFORMATES

In contrast to the facile reactivity of aliphatic alcohols toward phosgene, phenols are inert toward phosgene at temperatures below about 75°. However, aryl chloroformates are prepared easily in the presence of an acid acceptor (215, 322). Alkali metal hydroxides which yield the metal phenates are suitable. Early preparations of aryl chloroformates used these reagents. Conducting this reaction in the presence of an



organic solvent for the chloroformate has been widely used (729).

Many tertiary amines can be used as acid acceptors for the preparation of aryl chloroformates (45, 529). Amines of the dimethylaniline type gave the best results with this procedure, probably because of their low reactivity toward the chloroformates. Pyridines are also useful. The effectiveness of a tertiary amine in the synthesis of aryl chloroformates is probably not due solely to its ability to bind the hydrogen chloride that is formed. In addition, it probably forms an intermediate complex salt (7), *viz.*



which is a very reactive acylating agent for the phenol (45). The complex formed from 2 moles of pyridine and phosgene is well known to be a very active acylating agent (623). Aryl chloroformates (Table II) have been prepared which contained halogen (bromine, chlorine), nitro, methoxy, aldehyde, ester, sulfone, and double bond groupings.

### III. PHYSICAL PROPERTIES OF CHLOROFORMATES

#### A. STRUCTURE


Electric dipole moments for various chloroformates are recorded throughout the literature (356, 476, 481, 639). The experimental values are similar to those for simple acyclic carboxylic esters. For methyl chloroformate, values of 1.62 (639) and 2.22 (356) have been

TABLE II  
AROMATIC MONO- AND POLYCHLOROFORMATES:  $R(OCOCl)_n$

R	Base type <sup>a</sup>	Yield, %	Reference
2,4,6-(Br) <sub>3</sub> C <sub>6</sub> H <sub>3</sub>	O	74	348, 567
2,4,6-(Cl) <sub>3</sub> C <sub>6</sub> H <sub>3</sub>	I		348
2,4-(Cl) <sub>2</sub> C <sub>6</sub> H <sub>3</sub>	O	64	567
Various halogenated radicals	I		348
1,3-C <sub>6</sub> H <sub>4</sub>	O	>75	529
1,4-C <sub>6</sub> H <sub>4</sub>	O		566
2-(Cl)C <sub>6</sub> H <sub>4</sub>	O	>75	529
4-(Cl)C <sub>6</sub> H <sub>4</sub>	O	>75	351, 529, 568
2-(NO <sub>2</sub> )C <sub>6</sub> H <sub>4</sub>	O	24	349, 567
3-(NO <sub>2</sub> )C <sub>6</sub> H <sub>4</sub>	O	>75	323, 529
4-(NO <sub>2</sub> )C <sub>6</sub> H <sub>4</sub>	O	80	351
C <sub>6</sub> H <sub>5</sub>	O	70	25, 37, 38, 45, 197, 290, 304 308, 323 351, 377, 421, 568 324 38, 449, 556 556 82 156, 304, 351, 556 421 438 438 65 351 448 I 421 448 119, 421, 434, 435, 529, 555 52, 197, 198, 434, 435, 438, 570, 684, 734 438 438 80 567 742 421 425 75.2 621 87 621 93 146, 354, 621, 724, 726 78.3 621 79.4 621 468
2-(CHO)C <sub>6</sub> H <sub>4</sub>	I		
2-(CH <sub>3</sub> )C <sub>6</sub> H <sub>4</sub>	I		
3-(CH <sub>3</sub> )C <sub>6</sub> H <sub>4</sub>	I		
4-(CH <sub>3</sub> )C <sub>6</sub> H <sub>4</sub>	O		
4-(CH <sub>3</sub> CO)C <sub>6</sub> H <sub>4</sub>	I		
2-(CH <sub>3</sub> OCO)C <sub>6</sub> H <sub>4</sub>			
4-(CH <sub>3</sub> OCO)C <sub>6</sub> H <sub>4</sub>			
4-(C <sub>2</sub> H <sub>5</sub> O)C <sub>6</sub> H <sub>4</sub>	O		
2,6-(CH <sub>3</sub> ) <sub>2</sub> C <sub>6</sub> H <sub>3</sub>	O		
2,3,5-(CH <sub>3</sub> ) <sub>3</sub> C <sub>6</sub> H <sub>2</sub>	I		
2,4,6-(CH <sub>3</sub> ) <sub>3</sub> C <sub>6</sub> H <sub>2</sub>	O		
1-C <sub>10</sub> H <sub>7</sub>	I		
2-C <sub>10</sub> H <sub>7</sub>	O		
2,4-(CH <sub>3</sub> O)(CH <sub>2</sub> CH=CH)C <sub>6</sub> H <sub>4</sub>			
2,4-(CH <sub>3</sub> O)(CH <sub>2</sub> =CHCH <sub>2</sub> )C <sub>6</sub> H <sub>4</sub>			
2,4,5-(i-C <sub>3</sub> H <sub>7</sub> )(Cl)(CH <sub>2</sub> )C <sub>6</sub> H <sub>2</sub>	O		
2,5-(i-C <sub>3</sub> H <sub>7</sub> )(CH <sub>2</sub> )C <sub>6</sub> H <sub>3</sub>			
4-(t-C <sub>4</sub> H <sub>9</sub> )C <sub>6</sub> H <sub>4</sub>	I		
4-(C <sub>4</sub> H <sub>9</sub> OCO)C <sub>6</sub> H <sub>4</sub>	O		
S-(4-C <sub>6</sub> H <sub>4</sub> ) <sub>2</sub>	O		
(CH <sub>3</sub> ) <sub>2</sub> C[4-(2,6-Cl <sub>2</sub> C <sub>6</sub> H <sub>4</sub> ) <sub>2</sub> ]	O		
(CH <sub>3</sub> ) <sub>2</sub> C(4-C <sub>6</sub> H <sub>4</sub> ) <sub>2</sub>	O		
(CH <sub>3</sub> ) <sub>2</sub> C[4-(3-CH <sub>3</sub> C <sub>6</sub> H <sub>4</sub> ) <sub>2</sub> ]	O		
(CH <sub>2</sub> ) <sub>4</sub> C(4-C <sub>6</sub> H <sub>4</sub> ) <sub>2</sub>	O		
└─CH <sub>2</sub> ┘			
Diradical from a hydroxyl-terminated polystyrene			
Diradical from a low molecular weight hydroxyl-terminated bisphenol-A polycarbonate	I		148, 149

<sup>a</sup> I = inorganic base; O = organic base.

published. The fact that chloroformates possessed dipole moments of this magnitude led to the postulate that they possessed a *cis*-planar structure (III) similar to that of acyclic carboxylic esters (IV). Confirmation



of this structure in the case of methyl chloroformate was obtained from electron diffraction studies in the gas phase (530). The results of this study are tabulated in Table III. That this *cis*-planar con-

TABLE III  
BOND DISTANCES AND ANGLES OF CHLOROFORMATES

Property	CH <sub>3</sub> OCCl	ClCH <sub>2</sub> OCCl
C=O	1.19 ± 0.03 Å	1.20 ± 0.03 Å
C—O (carboxyl)	1.36 ± 0.04 Å	1.40 ± 0.04 Å
C—O (methoxyl)	1.47 ± 0.04 Å	1.40 ± 0.04 Å
C—Cl	1.75 ± 0.02 Å	1.74 ± 0.02 Å (both)
∠O=C—O	126 ± 4°	126 ± 4°
∠C—O—C	111 ± 4°	110 ± 4°
∠Cl—C—O	112 ± 3°	112 ± 4°
∠O—C—Cl (chloromethyl)		109.5 ± 3°

figuration is due at least in part to resonance was supported by the marked shortening found for the carboxyl C—O bond as well as the widening of the C—O—C angle. The greater than tetrahedral value of the Cl—C—O angle in methyl chloroformate was also suggestive of double bond character of the C—O bond; the O—C=O angle being close to the value (125°) of the classical model. A similar study (372) of the structure of chloromethyl chloroformate indicated that it possessed the *trans* (V) configuration (Table III). The chlorine of the chloromethyl group was *trans* to the carboxyl carbon atom, and the carboxyl C—O bond showed about 5% double bond character. Since  $\alpha$ -chlorochloroformates show differences in other properties, this effect on structure may be specific.

Other physical properties including densities (26, 182, 234, 355, 371, 546, 680), refractive indices (26, 182, 234, 355, 371, 546, 680), molar refractions (26), and parachors (490) are available for many chloroformates.

#### B. SPECTRAL STUDIES

##### 1. Infrared

The infrared absorptions of a series of chloroformates have been measured and reported (277, 366, 559). For simple aliphatic chloroformates, the carbonyl absorption occurred at 1770–1777 cm.<sup>-1</sup>. Substitution of a chlorine atom on the  $\beta$ -carbon of an aliphatic chloroformate caused no change in the position of the carbonyl absorption, whereas a similar substitution on the  $\alpha$ -carbon raised the carbonyl absorption to the 1793–

1806 cm.<sup>-1</sup> range. Similarly, aryl chloroformates possess carbonyl absorption at higher frequencies than aliphatic chloroformates. The C—O stretching frequency for chloroformates occurred in the 1050–1200 cm.<sup>-1</sup> range.

### 2. Raman

Raman spectra have been obtained for many chloroformates and are discussed in the literature (164, 358, 399, 400).

### 3. Nuclear Quadrupole Resonance


For ethyl chloroformate at liquid nitrogen temperatures, a resonance frequency of 33.858 Mc./sec. was obtained for Cl<sup>35</sup> (704).

## IV. CHEMICAL PROPERTIES OF CHLOROFORMATES

### A. REACTIONS OF CHLOROFORMATES WITH HYDROXYLIC COMPOUNDS

#### 1. Water




The products from the reaction of a chloroformate with water are the hydroxy compound from which it was prepared, hydrogen chloride, and carbon dioxide (519, 594). High yields are generally obtainable. The symmetrical carbonate is also formed, presumably by reaction of the hydroxy compound with starting chloroformate (457, 483).




Ethylene has been obtained from hydrolysis of ethyl chloroformate (186, 188, 676).



Chlorine-substituted chloroformates have yielded different products on hydrolysis (140, 141, 268, 389–391, 489). A few cases are depicted below.




The kinetics of the hydrolysis of chloroformates have been studied. An S<sub>N</sub>2 mechanism was found to be operative for the hydrolysis of chloroformates derived from primary and secondary alcohols (74, 75, 279, 687). Some recent work found an increase in the hydrolysis rate of chloroformates derived from secondary alcohols; hence a change in mechanism to S<sub>N</sub>1 has been postulated (161). Two postulated mechanisms for S<sub>N</sub>1 hydrolysis are shown. Path 1 was favored, al-



though not conclusively proved. Chloroformates of tertiary alcohols are known to be unstable at room temperature, with their facile decomposition probably proceeding through a carbonium ion.

The relative reactivity of ethyl chloroformate for S<sub>N</sub>2 hydrolysis, as compared to other acyl halides, is shown (278).




Hydrolysis of dimethylcarbamyl chloride proceeds via an S<sub>N</sub>1 mechanism accounting for the large increase in its rate.

#### 2. Alkali Metal Hydroxides

Addition of base to aqueous chloroformates can merely catalyze hydrolysis to yield the parent hydroxy compound (521). However, the use of a stoichiometric amount can yield the symmetrical carbonate, especially from aryl chloroformates (107, 483). Quantitative yields are readily obtained.



Chloroformates derived from tertiary alkyl radicals yield olefins on treatment with base (465).



An analytical method for chloromethyl and dichloromethyl chloroformates has been developed, based on their hydrolysis by sodium hydroxide (273).


#### 3. Aliphatic Alcohols and Thiols

Aliphatic alcohols react with chloroformates to yield carbonates and hydrogen chloride (Table IV). In many cases, the reaction proceeds at room temperature without a catalyst or hydrogen chloride acceptor (140, 141). However, faster reactions and better yields are usually ensured if alkali metals or their hydroxides, or tertiary amines, are used to react with the hydrogen chloride that is formed.


Thiols yield the monothiolocarbonates on reaction with chloroformates (599). Reaction of 2-mercaptopropanol with ethyl chloroformate in bulk gave the mercaptocarbonate as the only isolable product (586, 587).

TABLE IV  
REACTIONS OF CHLOROFORMATES,  $\text{ROCOCl}$ , WITH ALCOHOLS,  $\text{R}'(\text{OH})_m$


R	R'	Yield, %	Reference
<b>1. Reactions of Monochloroformates with Monoalcohols</b>			
$\text{Cl}_3\text{C}$	3-Cholesteryl		393
$\text{Cl}_2\text{CH}$	$\text{CH}_3$		392
$\text{Cl}_2\text{CH}$	3-Cholesteryl		393
$\text{ClCH}_2$	3-Cholesteryl		393
$\text{CH}_2=\text{CH}$	$\text{CH}_2=\text{CHCH}_2$		416
$\text{ClCH}_2\text{CHCl}$	$\text{C}_2\text{H}_5$		489
$\text{Cl}(\text{CH}_2)_2$	$\text{Cl}(\text{CH}_2)_2$	70	519
$\text{CH}_3\text{CHCl}$	$\text{C}_2\text{H}_5$		489
$\text{F}(\text{CH}_2)_2$	$\text{F}(\text{CH}_2)_2$	45	396, 532
$\text{F}(\text{CH}_2)_2$	$\text{C}_2\text{H}_5$	80	382
$\text{C}_6\text{H}_5$	$\text{C}_6\text{H}_5\text{O}(\text{CH}_2)_2$	86	715
$\text{C}_2\text{H}_5$	$1-(\text{C}\equiv\text{CH})-\text{C}_6\text{H}_{10}$		184
$\text{C}_2\text{H}_5$	$\text{C}_6\text{H}_5(\text{CH}_2)_3$	83	715
$\text{C}_2\text{H}_5$	$4-(\text{CH}_3)-1-(\text{C}\equiv\text{CH})-1-\text{C}_6\text{H}_{10}$		184
$\text{C}_2\text{H}_5$	Various radicals		628
$\text{CH}_2=\text{CHCH}_2$	$\text{CH}_2\text{OCHCH}_2$		487
$\text{CH}_2=\text{C}=\text{CHCH}_2$	Various radicals		105
$\text{C}_6\text{H}_5$	Various radicals		120
$\text{CH}_2=\text{CHCH}_2\text{OCOCH}_2$	$\text{CH}_2=\text{CHCH}_2\text{OCOCH}_2$	67	443
$\text{C}_6\text{H}_5$	$\text{C}_2\text{H}_5$		483
$\text{C}_6\text{H}_5$	$\text{HC}\equiv\text{CC}(\text{R}'')(\text{R}''')$ ( $\text{R}''$ and $\text{R}'''$ are various aliphatic radicals)		455
$\text{C}_6\text{H}_5$	$\text{C}_6\text{H}_5\text{CH}(\text{CH}_2\text{R}'')$ ( $\text{R}''$ are N-pyrrolidinyl, N-piperidinyl, N-morpholinyl, and N-(N'-methyl)piperazinyl)	~70	294
$\text{C}_6\text{H}_5$	Various radicals		483, 422
$4-(\text{NO}_2)\text{C}_6\text{H}_4\text{CH}_2$	$\text{C}_2\text{H}_5$	107	
$4-(\text{NO}_2)\text{C}_6\text{H}_4\text{CH}_2$	$4-(\text{NO}_2)\text{C}_6\text{H}_4\text{CH}_2$	52	107
Various aliphatic radicals	$\text{R}'\text{NCOCCH}_2$ ( $\text{R}''$ are various aliphatic and aromatic radicals)	20-96	632
Various aliphatic radicals	$\text{CH}_3\text{CH}(\text{COOR}'')$ ( $\text{R}''$ are various aliphatic radicals)	30-85	576
Various aliphatic radicals	$\text{CH}_3\text{CH}(\text{CONHR}'')$ ( $\text{R}''$ are various aliphatic and aromatic radicals)	40-90	632
Various aliphatic radicals	$(\text{CH}_3)_2\text{C}(\text{CONHR}'')$ ( $\text{R}''$ are various aliphatic and aromatic radicals)	30-90	632
Various aliphatic radicals	$\text{C}_6\text{H}_5\text{CH}(\text{CH}_2\text{R}'')$ ( $\text{R}''$ are N-pyrrolidinyl, N-piperidinyl, N-morpholinyl, and N-(N'-methyl)piperazinyl)	~85	294
Various aliphatic radicals	$\text{C}_6\text{H}_5\text{CH}(\text{CONHR}'')$ ( $\text{R}''$ are various aliphatic and aromatic radicals)	30-95	632
Various aliphatic radicals	$\text{CH}_3$		426, 540
Various aromatic radicals	Various radicals		197
Various aliphatic and aromatic radicals	Various radicals		339, 594, 630
Various steroidal radicals	$\text{CH}_3$	180	
Various steroidal radicals	$\text{C}_2\text{H}_5$	180	
<b>2. Reactions of Monochloroformates with Diols</b>			
$\text{CH}_3$	Radical from cellulose		595
$\text{C}_2\text{H}_5$	$-\text{CH}_2\text{CH}=\text{CHCH}_2-$		54
$\text{C}_2\text{H}_5$	$1,4-\text{CH}_2\text{C}_6\text{H}_4\text{CH}_2$		583
$\text{C}_2\text{H}_5$	$1,4-(\text{CH}_2)_2\text{C}_6\text{H}_4(\text{CH}_2)_2$	80	585
$\text{C}_2\text{H}_5$	$1,4-(\text{CH}_2)_2\text{OC}_6\text{H}_4\text{O}(\text{CH}_2)_2$	>54	584
$\text{CH}_2=\text{CHCH}_2$	$\text{O}[\text{Si}(\text{CH}_3)_2\text{CH}_2]_2$	30	9
$\text{CH}_3(\text{CH}_2)_4$	$1,4-\text{CH}_2\text{C}_6\text{H}_4\text{CH}_2$		583
$\text{C}_6\text{H}_5$	$1,4-\text{CH}_2\text{C}_6\text{H}_4\text{CH}_2$		583



However, the greater reactivity of the anion from a thiol as compared to that from an alcohol was shown by the formation of the monothiolcarbonate from the same reaction when carried out with added sodium hydroxide (506, 586, 587).




$\alpha$ -Hydroxy esters (676) and  $\alpha$ -hydroxy amides (632) reacted with chloroformates to yield the expected carbonates. The reported yields were variable.



$\text{R}' = -\text{NHR}''$  or  $-\text{OR}''$

A carbonate has been prepared in low yield from a cyanohydrin by reaction of ethyl chloroformate with acetone in the presence of potassium cyanide (168).



#### 4. Phenols

Phenols are unreactive toward chloroformates at room temperature, and at elevated temperatures only

small yields (<10%) of the expected carbonates are produced. However, phenols react readily with chloroformates when bases are used (224), and high yields of the expected carbonates are obtainable (Table V).





TABLE V  
REACTIONS OF CHLOROFORMATES, ROCOCl, WITH PHENOLS, Ar(OH)<sub>m</sub>

R	Ar	Yield, %	Reference
<b>1. In the Presence of a Stoichiometric Amount of Base</b>			
CH <sub>3</sub>	2,4,5,6-(Cl) <sub>3</sub> (NO <sub>2</sub> )C <sub>6</sub> H		635
CH <sub>3</sub>	5-Acetyl-8-quinolyl		318, 319
CH <sub>3</sub>	5-Butyryl-8-quinolyl		318, 319
CH <sub>3</sub>	5-Valeryl-8-quinolyl		318, 319
CH <sub>3</sub>	(CH <sub>3</sub> ) <sub>2</sub> C(4-C <sub>6</sub> H <sub>4</sub> ) <sub>2</sub>		59
CH <sub>3</sub>	Various radicals		312, 313
C <sub>2</sub> H <sub>5</sub>	2,4,5,6-(Cl) <sub>3</sub> (NO <sub>2</sub> )C <sub>6</sub> H		635
C <sub>2</sub> H <sub>5</sub>	C <sub>6</sub> H <sub>5</sub>	71	138, 195, 225, 718
C <sub>2</sub> H <sub>5</sub>	2-(COOH)C <sub>6</sub> H <sub>4</sub>		395
C <sub>2</sub> H <sub>5</sub>	4-(COOH)C <sub>6</sub> H <sub>4</sub>		682
C <sub>2</sub> H <sub>5</sub>	5-Acetyl-8-quinolyl		318, 319
C <sub>2</sub> H <sub>5</sub>	5-Butyryl-8-quinolyl		318, 319
C <sub>2</sub> H <sub>5</sub>	5-Valeryl-8-quinolyl		318, 319
C <sub>2</sub> H <sub>5</sub>	2-Formyl-5-acetyl-8-quinolyl		319, 318
C <sub>2</sub> H <sub>5</sub>	Various radicals		312, 313
CH <sub>2</sub> =CHCH <sub>2</sub>	4-Cl-1,3-C <sub>6</sub> H <sub>4</sub>		84
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>2</sub>	2,4,5,6-(Cl) <sub>3</sub> (NO <sub>2</sub> )C <sub>6</sub> H		635
(CH <sub>3</sub> ) <sub>2</sub> CH	2,4,5,6-(Cl) <sub>3</sub> (NO <sub>2</sub> )C <sub>6</sub> H		635
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>2</sub>	5-Acetyl-8-quinolyl		318, 319
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>2</sub>	5-Butyryl-8-quinolyl		318, 319
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>2</sub>	5-Valeryl-8-quinolyl		318, 319
CH <sub>2</sub> =C=CHCH <sub>2</sub>	Various radicals		105
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>3</sub>	2,4,5,6-(Cl) <sub>3</sub> (NO <sub>2</sub> )C <sub>6</sub> H		635
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>3</sub>	5-Acetyl-8-quinolyl		318, 319
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>3</sub>	5-Butyryl-8-quinolyl		318, 319
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>3</sub>	5-Valeryl-8-quinolyl		318, 319
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>4</sub>	2,4,5,6-(Cl) <sub>3</sub> (NO <sub>2</sub> )C <sub>6</sub> H		635
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>4</sub>	5-Acetyl-8-quinolyl		318, 319
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>4</sub>	5-Butyryl-8-quinolyl		318, 319
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>4</sub>	5-Valeryl-8-quinolyl		318, 319
4-(Cl)C <sub>6</sub> H <sub>4</sub>	1,4-C <sub>6</sub> H <sub>4</sub>	58	568
C <sub>6</sub> H <sub>5</sub>	C <sub>6</sub> H <sub>5</sub>		314
C <sub>6</sub> H <sub>5</sub>	5-Acetyl-8-quinolyl		318, 319
C <sub>6</sub> H <sub>5</sub>	5-Butyryl-8-quinolyl		318, 319
C <sub>6</sub> H <sub>5</sub>	5-Valeryl-8-quinolyl		318, 319
C <sub>6</sub> H <sub>5</sub>	(CH <sub>3</sub> ) <sub>2</sub> C(4-C <sub>6</sub> H <sub>4</sub> ) <sub>2</sub>		79
Various halogenated aromatic radicals	5-Acetyl-8-quinolyl		318, 319
Various halogenated aromatic radicals	5-Butyryl-8-quinolyl		318, 319
Various halogenated aromatic radicals	5-Valeryl-8-quinolyl		318, 319
Various aromatic radicals	2-Formyl-5-acetyl-8-quinolyl		318, 319
Various aromatic radicals	Various radicals		314
Various aliphatic and aromatic radicals	Various radicals	~90	229, 230, 667
Various aliphatic radicals	2-(NO <sub>2</sub> )C <sub>6</sub> H <sub>4</sub>	70-95	381
Various aliphatic radicals	Various diradicals	~90	229, 230, 667
Various aliphatic radicals	Various radicals		121
3-Cholesteryl	Various di- and polyyradicals		604
3-Cholesteryl	Various radicals		604
<b>2. Catalytic</b>			
C <sub>2</sub> H <sub>5</sub> <sup>a</sup>	C <sub>6</sub> H <sub>5</sub>		545
C <sub>9</sub> H <sub>19</sub> <sup>b</sup>	C <sub>6</sub> H <sub>5</sub>		625

<sup>a</sup> With AlCl<sub>3</sub> as catalyst. <sup>b</sup> Activated carbon catalyst.

Sodium hydroxide and amines, such as pyridine and dimethylaniline, are particularly useful. Activated carbon has been claimed to be a very good catalyst for the reaction of phenol with an aliphatic chloroformate (345).

The cyclizations shown in eq. a (737), and b (364) have been reported.




### 5. Heterocyclic Alcohols


Alcohols of nitrogen-containing heterocycles have been converted to carbonates by reaction with chloroformates (96, 171, 316, 514, 515, 647). Furan- and tetrahydrofuran-derived alcohols reacted similarly to give carbonates in 75% yield (715). As with aliphatic alcohols, inorganic bases and tertiary amines increased reactivity and the yields.

### B. REACTION OF CHLOROFORMATES WITH CARBOXYLIC ACIDS


Most of the early literature pertaining to the reaction of a chloroformate with a carboxylic acid described the formation of esters as the main products (3, 29, 196, 310, 335, 336, 395, 538). The conditions that were used



generally employed heating and/or acidic or basic catalysts. Good yields of esters were obtained. Esters of fluorinated carboxylic acids can also be advantageously prepared in 75–80% yields by this method (266). A variation from the usual result is shown in the following equation (174, 409).




For many years, the expected intermediates in this reaction, the mixed carboxylic-carbonic anhydrides (VI), i.e.




were thought to be incapable of isolation and purification because of their thermal instability. However, several groups (216, 670, 721) have recently synthesized and isolated many examples of this class of compounds (VI). The mechanism of the thermal decom-

position of these mixed carboxylic-carbonic anhydrides has been the subject of recent publications. Both Tarbell and Longosz (437, 671) and Windholz (722, 723) have observed paths 1 and 2 below. Reaction 3




was also demonstrated to occur, but temperatures much higher than those used for the mechanism studies were required to effect it. The preferred route (1 or 2) of decomposition was found to be dependent on the particular groups (R, R') that were present, the temperature of decomposition, and the presence or absence of catalysts. For benzoic n-butylcarbonic anhydride (VI, R = C<sub>6</sub>H<sub>5</sub>, R' = C<sub>6</sub>H<sub>5</sub>), Tarbell and Longosz found that their results agreed with the idea that the products were formed by a series of ionic chain reactions which were initiated by catalysts acting as nucleophiles (437). Cyclic transition states of types VII and VIII were ruled out on the basis of a study of the thermal decomposition of benzoic sec-butylcarbonic




anhydride containing O<sup>18</sup> in the alcohol fragment (475). No alkyl-oxygen cleavage was observed, and hence transition states of types VII and VIII could not be operative. It was pointed out, however, that when R is benzyl or a tertiary radical, alkyl-oxygen cleavage might occur to yield the more stable carbonium ions. Early workers (335) had shown that the reaction of potassium benzoate with (−)-2-octyl chloroformate resulted in the recovery, after hydrolysis, of only (−)-2-octanol. This retention of configuration was reconfirmed by Tarbell, *et al.*, by the preparation and decomposition of pure benzoic (−)-2-octylcarbonic anhydride (671). It was found that the resulting 2-octyl benzoate gave pure (−)-2-octanol on saponification. This stereochemical result is compatible with the ionic chain mechanism proposed above.

Pyrocarbonates (anhydrides of carbonic acids) have been prepared by the following routes (169, 406, 677).



The two pyrocarbonates shown are fairly stable compounds but do decompose at temperatures over 100° to yield the corresponding ester and carbon dioxide.

Mixed anhydrides of peroxy carboxylic acids have been prepared and isolated from chloroformates using pyridine as a hydrogen chloride acceptor (541). They




have been claimed to have utility as initiators for vinyl polymerization.

### C. REACTIONS OF CARBOXYLIC-CARBONIC ANHYDRIDES


Carboxylic-carbonic anhydrides are very active acylating agents and were widely used for this purpose even before their isolation. Their reaction with nucleophiles proceeds readily under mild conditions to give good yields of the expected products.

#### 1. With Alcohols (39, 335, 359)


The preparation of esters by the route shown in the following equation is especially useful in cases where





the acids are thermally or otherwise unstable. The use of chloroformates as esterification catalysts has been reported (238). Their utility may stem from the formation of the mixed anhydride which then reacts rapidly with the alcohol.




#### 2. With Thiols (39, 335, 359)



#### 3. With Phenols (669) and Thiophenols (714)




#### 4. With Acids (611, 669)



#### 5. With Amines


The reaction of amines (39, 73, 315, 335, 359, 581, 609, 611, 694-698, 702, 713) with carboxylic-carbonic

anhydrides to yield amides has been used widely for the synthesis of peptide bonds because of the requisite mild reaction conditions. Excellent yields are obtainable. In a recent study, Tarbell and Leister (427) have shown that this reaction can proceed by the three routes shown below, depending on the particular structure of the




carboxylic-carbonic anhydride. In many cases when benzoic-carbonic anhydrides were treated with amines, urethans (path 2) were the major products. Paths 1 and 2 are merely the result of attack at either the carboxylic or carbonic carbonyl groups, whereas path 3 is a disproportionation of the anhydride.


#### 6. With Organometallic Reagents (668, 669)



#### 7. With Diazoalkanes (34, 669)




#### 8. With Hydrocarbons (129, 167, 191)



D. REACTIONS OF CHLOROFORMATES WITH  
NITROGEN COMPOUNDS


1. Ammonia

The reaction of chloroformates with ammonia is the classical method for preparing primary urethans and, therefore, has been used extensively in the past for this purpose. In most cases, ammonia is used in excess to remove the hydrogen chloride. The utility of primary urethans, and urethans in general, in diverse applications within the general areas of biological and agricultural chemicals has contributed to the widespread use of this reaction. As can be seen from the examples listed in Table VI, this reaction has been effected successfully in the presence of many types of reactive groups, even some of which are normally considered to be reactive towards ammonia. These include double and triple bonds, alenic groups, alkyl chlorides and fluorides, hydroxyl groups, cyclic and acyclic ethers, and esters of carbonic and carboxylic acids. Bis(urethans) have also been prepared by this method. The reaction of the chloroformate of pentachloroethanol with ammonia is a special case (140, 141, 489).




2. Primary and Secondary Aliphatic Amines

Primary and secondary aliphatic amines react rapidly with chloroformates to yield the urethans. Organic (excess amine) or inorganic bases can be used to remove the hydrogen chloride that is formed. For many of the same reasons mentioned above with respect to the reaction of ammonia with chloroformates, this reaction has been used extensively throughout the literature (Table VII). Reaction of ethylenimine with ethyl chloroformate in the presence of a tertiary amine as the hydrogen chloride acceptor gave the expected urethan (351) (Table VIII).




However, when no acid acceptor was present, ring opening occurred to give the  $\beta$ -chlorourethan (60).



An interesting synthesis of oxazolidines and 1,3-oxazines resulted from the reaction of aminovinyl esters with alkyl or aryl chloroformates (462).


TABLE VI  
REACTIONS OF CHLOROFORMATES,  $\text{R}(\text{OCOCl})_n$ ,  
WITH AMMONIA

R	Yield, %	Reference
$-(\text{CH}_2)_4-$		354
$\text{Cl}(\text{CH}_2)_2$		155
$\text{F}(\text{CH}_2)_2$	85	382, 532
$\text{C}_2\text{H}_6$		186, 188
$-(\text{CH}_2)_3-$		354
$\text{Cl}(\text{CH}_2)_3$	71	185
$\text{C}_2\text{H}_7$		676
$\text{HC}\equiv\text{CCH}(\text{CH}_3)$		553, 554
$\text{CH}_2=\text{C}=\text{CHCH}_2$		105, 106
$\text{OCOOCH}_2\text{CHCH}_2$		658
$\boxed{\quad}$		
$-\text{CH}_2\text{CH}=\text{CHCH}_2-$	90	276
$\text{CH}_2\text{OCH}_2\text{CH}_2\text{CH}$	$\sim 90$	578
$-(\text{CH}_2)_4\text{C}$		354
$\text{CH}_3\text{CH}(\text{CH}_3)\text{CH}_2$		676
$-(\text{CH}_2)_5-$		354
$\text{CH}_3\text{C}(\text{CH}_3)(\text{C}_2\text{H}_5)$		466
$\text{C}_6\text{H}_5$		377, 489
$-(\text{CH}_2)_6-$		354
$\text{Cl}(\text{CH}_2)_6$		298
$\text{C}_6\text{H}_5\text{CH}_2$	90	112
$-(\text{CH}_2)_7-$		354
$\begin{array}{c}   \\ \text{CH}_3\text{CHC}(\text{CH}_3)_2\text{CHCH}_3 \end{array}$		128
$(i\text{-C}_3\text{H}_7)_2\text{CH}$		669
$\text{CF}_3(\text{CF}_2)_8\text{CH}_2$	$\sim 70$	520
$-(\text{CH}_2)_8-$		354
$\text{Cl}(\text{CH}_2)_8$		298
1-(1-Cyclohexenyl)propargyl	40	553, 554
$-(\text{CH}_2)_9-$		354
$\text{Cl}(\text{CH}_2)_9$		298
$[(\text{CH}_3)_3\text{C}]_2\text{CH}$		379
1-(2-Bicyclo[2.2.1]heptenyl-2)-propargyl	53	553, 554
$(\text{C}_6\text{H}_5)(i\text{-C}_3\text{H}_7)\text{CH}$		662-664
$-(\text{CH}_2)_{10}-$		354
$\text{Cl}(\text{CH}_2)_{10}$		298
$\text{ClCF}_2(\text{CF}_2)_9\text{CH}_2$	$\sim 70$	520
$\text{CHF}_2(\text{CF}_2)_9\text{CH}_2$	$\sim 70$	520
$4-(\text{C}_4\text{H}_9\text{OCO})\text{C}_6\text{H}_4$		425
$\text{CHF}_2(\text{CF}_2)_{11}\text{CH}_2$	$\sim 70$	510
$\text{C}_2\text{H}_5\text{OCOC}(\text{CH}_3)_2\text{CH}(\text{C}_6\text{H}_5)$		8
$(\text{C}_6\text{H}_5)_2\text{C}(\text{CH}_2\text{OH})\text{CH}_2$		56
3-Ethylcholatetyl		701
$\text{HC}\equiv\text{CC}(\text{R}')(\text{R}'')$ ( $\text{R}'$ and $\text{R}''$ are various aliphatic, cycloaliphatic, and/or aromatic radicals)		516
$\text{HC}\equiv\text{CCH}_2\text{C}(\text{R}')(\text{R}'')$ ( $\text{R}'$ and $\text{R}''$ are various aliphatic and/or aromatic radicals; together they may form an alicyclic radical)	60-70	422, 423
Various aromatic radicals		311
Various chlorinated aliphatic radicals		743
$\text{R}'\text{O}(\text{CH}_2)_2$ ( $\text{R}'$ are various aliphatic radicals)	13-66	20
$-\text{CH}_2\text{C}(\text{R}')(\text{R}'')\text{CH}_2-$ ( $\text{R}'$ and $\text{R}''$ are various aliphatic or aromatic radicals)	60-90	439
Various aliphatic radicals	55-80	48, 49, 440




Recent kinetics studies have shown that the reaction of primary and secondary amines is an extremely fast one which proceeds by an  $S_N2$  mechanism (280, 282). This is due to the high nucleophilicity of the amines and the ease of displacing a chloride ion attached to a carbonyl group. Rate constants ( $k_2$ ) of the order of  $10^4$  were found for piperidine;  $10^2$  for morpholine, 1-carboethoxypiperazine, 2-methylpiperidine, *n*-butylamine, and neopentylamine; 10 for *sec*-butylamine; and 1 for *t*-butylamine. The reaction is markedly subject to steric hindrance although it was surprising that neopentylamine did not exhibit a conspicuous degree of hindrance.


Reaction of hydroxylamine with a chloroformate leads to N-acylation preferentially, *viz.* (292)



However, the use of 3 equiv. of a chloroformate can lead to the replacement of the three active hydrogens of hydroxylamine (85). Other examples of this reaction have also been reported (749, 750).



When N-acylation is not possible as with dialkyl hydroxylamines, aldoximes, and ketoximes, O-acylation occurs readily, *viz.* (81, 82).




### 3. Primary and Secondary Aromatic Amines


Aromatic primary and secondary amines react with chloroformates in the same manner as do aliphatic amines. However, since aromatic amines are weaker bases (and weaker nucleophiles), the reactions are slower than with aliphatic amines. For reasons of economy, it is more advantageous to use inorganic bases as the hydrogen chloride acceptor instead of excess amine. The numerous examples shown in Table IX are indicative of the generality of this reaction, even with many types of other functional groups present.

An interesting use of an intermediate urethan pre-

pared by this route in the synthesis of 3-aryl-substituted 2-oxazolidones has been reported (5, 453).




A similar cyclization has yielded cyclic ureas (738).



### 4. Tertiary Amines


Tertiary amines react with chloroformates to yield quaternary ammonium compounds of the general structure IX (42, 44, 47, 122-124, 329, 394, 543, 590).



The properties of these salts vary depending on the specific R groups that are present.

#### a. Thermal Reactions of Tertiary Amine-Chloroformate Complexes

Complexes formed from aliphatic chloroformates with pyridine are unstable at room temperature (329). Reaction of pyridine with ethyl chloroformate at  $0^\circ$  gave ethyl chloride, carbon dioxide, pyridine, and pyridine hydrochloride. Chloroformates derived from secondary alcohols react with pyridine quite cleanly to yield the secondary chlorides (544). This reaction has been shown to proceed with inversion in the case of 1,2-octyl and 1,1-phenethyl chloroformates (335-338). Inversion probably occurs by backside attack of the chloride ion on the cationic complex ion.



Catalytic amounts of triethylamine have been used

TABLE VII  
REACTIONS OF CHLOROFORMATES,  $R'(OCOCl)_n$ , WITH ALIPHATIC AMINES

R	R'	1. Primary Amines $R'(NH_2)_m$	
		Yield, %	Reference
$-(CH_2)_2-$	$C_4H_9$		354
$Cl(CH_2)_2$	$CH_3$		626
$Cl(CH_2)_2$	Various radicals	~70	558, 626
$F(CH_2)_2$	$CH_3$	81	532
$C_2H_5$	HO		292
$C_2H_5$	$-(CH_2)_2-$		582
$C_2H_5$	$-(CH_2)_3-$		582
$C_2H_5$	$-(CH_2)_6-$		582
$C_2H_5$	$C_6H_5CH(CH_3)$		705
$C_2H_5$	$-(CH_2)_8-$		582
$C_2H_5$	Various radicals		282
$ClCH=CHCH_2$	$Cl(CH_2)_2$	46	130
$CH_2=CHCH_2$	HO		85
$-(CH_2)_3-$	$C_4H_9$		354
$Cl(CH_2)_3$	Various radicals	~70	558
$ClCH_2C\equiv CCH_2$	Various radicals	10-90	31, 330, 685
$CH_2OCOOCHCH_2$	Various radicals		658
$[-]$			
$-CH_2CH=CHCH_2-$	$CH_3$		276
$CH_2O(CH_2)_2CH$	$-(CH_2)_5-$		578
$-(CH_2)_4-$	$C_4H_9$		354
$+(CH_2)_2+\_O$	2-(12-Hydroxyoleoyloxy)ethyl		170
$+(CH_2)_2+\_O$	Various radicals	64-99	661
$-(CH_2)_5-$	$C_4H_9$		354
$C_2H_5C(CH_3)_2$	Various radicals		466
$1,4-C_6H_4$	$C_4H_9$		354
$C_6H_5$	Various radicals		484
1,4-Cyclohexylene ( <i>cis</i> )	$C_4H_9$		354
1,4-Cyclohexylene ( <i>trans</i> )	$C_4H_9$		354
$-(CH_2)_6-$	$C_4H_9$		354
2-Chlorobenzyl	$(C_2H_5)_2N(CH_2)_2$		295
$-(CH_2)_7-$	$C_4H_9$		354
$4-(CH_3CO)C_6H_4$	$CH_3$		421
$2,6-(CH_3)_2C_6H_3$	Various radicals	70-80	448
$-(CH_2)_8-$	$C_4H_9$		354
2,6-Dimethylbenzyl	$(C_2H_5)_2N(CH_2)_2$		295
$2,4,6-(CH_3)_3C_6H_2$	Various radicals	70-80	448
$-(CH_2)_9-$	$C_4H_9$		354
1-Naphthyl	$CH_3$		420
1-Naphthyl	$C_4H_9$	~70	434, 435
2-Naphthyl	$C_4H_9$	~70	434, 435
1-Naphthyl	Various radicals		421
5,6,7,8-Tetrahydro-2-naphthyl	$C_4H_9$	~70	434, 435
$1,4-[-(CH_2)_2OCO]_2C_6H_4$	$C_4H_9$		354
$(CH_3)_2C(4-C_6H_4)_2$	$C_4H_9$		354
3-Cholesteryl	$C_6H_5CH_2$	89	413, 454
Various aromatic radicals	1-Methyl		556
Various aliphatic and aromatic radicals	$(CH_3)_2N(CH_2)_3$	50-55	404, 405
Various aliphatic and aromatic radicals	$(C_2H_5)_2N(CH_2)_3$	25-85	404, 405
Various aliphatic and aromatic radicals	$(CH_2(CH_2)_4N(CH_2)_2$	50-80	404, 405
$[-]$	Various radicals		
1-Propargyl alicyclic radicals			423
$(R_1)(R_2)(HC\equiv C)C$ ( $R_1$ and $R_2$ are various aliphatic radicals)	Various radicals		516
$-CH_2C(C_2H_5)_2CH_2-$	Various radicals	50-60	439
Various aliphatic radicals	Camphoryl	~90-95	227
Various tertiary aliphatic radicals	Various radicals		466
Various aliphatic, alicyclic, and heterocyclic radicals	Various bivalent radicals		346


TABLE VII (Continued)  
2. Secondary Amines, R'R''NH

R	R'	R''	Yield, %	Reference
F(CH <sub>2</sub> ) <sub>2</sub>	CH <sub>3</sub>	CH <sub>3</sub>	85	532
C <sub>2</sub> H <sub>5</sub>	Cyclopropyl	C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub>		2
CH <sub>2</sub> =C=CHCH <sub>2</sub>	CH <sub>3</sub>	CH <sub>3</sub>		105, 106
+-(CH <sub>2</sub> ) <sub>2</sub> -O-	C <sub>4</sub> H <sub>9</sub>	C <sub>4</sub> H <sub>9</sub>	70	
C <sub>6</sub> H <sub>5</sub>	CH <sub>3</sub>	CH <sub>3</sub>		421
C <sub>6</sub> H <sub>5</sub>	Various radicals	Various radicals		484
1-Naphthyl	CH <sub>3</sub>	CH <sub>3</sub>		421, 434, 435
2-Naphthyl	CH <sub>3</sub>	CH <sub>3</sub>		434, 435
1-Naphthyl	Cl(CH <sub>2</sub> ) <sub>2</sub>	Cl(CH <sub>2</sub> ) <sub>2</sub>	80	434, 435
2-Naphthyl	Cl(CH <sub>2</sub> ) <sub>2</sub>	Cl(CH <sub>2</sub> ) <sub>2</sub>	60	434, 435
1-Naphthyl	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>		434, 435
2-Naphthyl	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>		434, 435
1-Naphthyl	C <sub>3</sub> H <sub>7</sub>	C <sub>3</sub> H <sub>7</sub>		434, 435
1-Naphthyl	C <sub>4</sub> H <sub>9</sub>	C <sub>4</sub> H <sub>9</sub>		421, 434, 435
1-Naphthyl	(CH <sub>3</sub> ) <sub>2</sub> CH(CH <sub>2</sub> ) <sub>2</sub>	(CH <sub>3</sub> ) <sub>2</sub> CH(CH <sub>2</sub> ) <sub>2</sub>	70	434, 435
1-Naphthyl	C <sub>6</sub> H <sub>11</sub>	C <sub>6</sub> H <sub>11</sub>	60	434, 435
5,6,7,8-Tetrahydro-1-naphthyl	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>	70	434, 435
5,6,7,8-Tetrahydro-2-naphthyl	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>		434, 435
5,6,7,8-Tetrahydro-1-naphthyl	C <sub>4</sub> H <sub>9</sub>	C <sub>4</sub> H <sub>9</sub>		434, 435
4-[(CH <sub>3</sub> ) <sub>3</sub> C]C <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	CH <sub>3</sub>		421
Various aromatic radicals	Various radicals	Various radicals		691
(R <sub>1</sub> )(R <sub>2</sub> )(HC≡C)C (R <sub>1</sub> and R <sub>2</sub> are various aliphatic radicals)	Various radicals	Various radicals		516
Various tertiary aliphatic radicals	Various radicals	Various radicals		466
-(CH <sub>2</sub> CH <sub>2</sub> NH) <sub>n</sub> -	Various radicals	Various radicals		688


TABLE VIII  
REACTION OF CHLOROFORMATES, R(OCOCl)<sub>n</sub>,  
WITH ETHYLENIMINE

R	Yield, %	Reference
-(CH <sub>2</sub> ) <sub>2</sub> -		354
C <sub>2</sub> H <sub>5</sub>	60-65	60
-(CH <sub>2</sub> ) <sub>3</sub> -	25	354
-(CH <sub>2</sub> ) <sub>4</sub> -	50-60	354
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>3</sub>	78	351
-(CH <sub>2</sub> ) <sub>5</sub> -	38	354
1,4-C <sub>6</sub> H <sub>4</sub>	58	354
4-(Cl)C <sub>6</sub> H <sub>4</sub>	70-80	351
4-(NO <sub>2</sub> )C <sub>6</sub> H <sub>4</sub>	70-80	351
C <sub>6</sub> H <sub>5</sub>	70-80	351
1,4-Cyclohexylene ( <i>cis</i> )		354
1,4-Cyclohexylene ( <i>trans</i> )	53	354
Cyclohexyl	72	351
-(CH <sub>2</sub> ) <sub>6</sub> -	60	354
4-(CH <sub>3</sub> )C <sub>6</sub> H <sub>4</sub>	70-80	351
-(CH <sub>2</sub> ) <sub>7</sub> -		354
4-(CH <sub>3</sub> CH <sub>2</sub> O)C <sub>6</sub> H <sub>4</sub>	70-80	351
-(CH <sub>2</sub> ) <sub>8</sub> -		354
-(CH <sub>2</sub> ) <sub>9</sub> -	35	354
-(CH <sub>2</sub> ) <sub>10</sub> -	77	354
1,4-[-(CH <sub>2</sub> ) <sub>2</sub> OCO] <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	76	354
(CH <sub>3</sub> ) <sub>2</sub> C(4-C <sub>6</sub> H <sub>4</sub> ) <sub>2</sub>		354
Various bivalent radicals		480
Polymeric polyoxyethylene radicals	88-91	352, 353


for the preparation of acetylenic chlorides from their respective chloroformates (28, 231).



With chloroformates derived from tertiary alcohols, elimination occurs on reaction with pyridine (466). Extensive studies on the decomposition of chloroformates in the presence of organic bases have been made (108, 109, 111, 249).



Cleavage of the pyridine ring has been reported when 2- and 4-alkyl-substituted pyridines reacted with chloroformates (692, 693). This ring opening reaction



has been used for the degradation of some natural products (90, 232, 233, 242, 708).


Reaction of pyridine with ethyl chloroformate in the presence of zinc dust has been investigated (692, 693). The major products from this reaction are shown below.

TABLE IX  
REACTIONS OF CHLOROFORMATES,  $R(OCOCl)_n$ , WITH AROMATIC AMINES  
1. Primary Amines,  $R'(NH_2)_m$


$R$	$R'$	Yield, %	Reference
$CH_3$	4-( $H_2O_3Sb$ ) $C_6H_4$	73	288
$-(CH_2)_2-$	$C_6H_5$		354
$Cl(CH_2)_2$	$C_6H_5$		521
$Cl(CH_2)_2$	2-Methyl-1-anthraquinonyl	70	507
$Cl(CH_2)_2$	Various radicals	>95	453
$C_2H_5$	$C_6H_5$	75	138, 719
$C_2H_5$	4-( $H_2O_3Sb$ ) $C_6H_4$	85	288
$C_2H_5$	4-( $NaO_3SCH_2O$ ) $C_6H_4$	77	36
$C_2H_5$	2-( $NaOCOCO$ ) $C_6H_4$		293
$C_2H_5$	2,5-[ $CH_3COO(CH_2)_8$ ]( $NO_2$ ) $C_6H_3$	74	539
$C_2H_5$	$CH_2(4-C_6H_4)_2$		99
$ClCH=CHCH_2$	$C_6H_5$		130
$-(CH_2)_3-$	$C_6H_5$		354
$C_3H_7$	4-( $H_2O_3Sb$ ) $C_6H_4$	87	288
$(CH_3)_2CH$	4-( $H_2O_3Sb$ ) $C_6H_4$	89	288
$CH_2OCOOCHCH_2$	Various radicals		658
$[-]$			
$-CH_2CH=CHCH_2-$	$C_6H_5$		276
$BrCH_2C\equiv CCH_2$	Various radicals	>50	330, 331
$ClCH_2C\equiv CCH_2$	Various radicals	30-90	330, 331
$CH_2O(CH_2)_2CH$	$C_6H_5$		578
$[-]$			
$-(CH_2)_4-$	$C_6H_5$		354
$+-(CH_2)_2+-O$	3-( $Cl$ ) $C_6H_4$	64	661
$+-(CH_2)_2+-O$	$C_6H_5$	78	661
$C_4H_9$	4-( $H_2O_3Sb$ ) $C_6H_4$	70	288
$C_4H_9$	Various radicals		120
$(CH_3)_2CHCH_2$	4-( $H_2O_3Sb$ ) $C_6H_4$	85	288
$-(CH_2)_5-$	$C_6H_5$		354
$1,4-C_6H_4$	$C_6H_5$		354
$4-(Cl)C_6H_4$	1,4- $C_6H_4$	91	568
$C_6H_5$	1,4- $C_6H_4$	57	568
$C_6H_5$	2-( $Cl$ ) $C_6H_4$		568
$C_6H_5$	$C_6H_5$	95	270, 484, 568
$C_6H_5$	Various radicals		484
$-(CH_2)_6-$	$C_6H_5$		354
$CH_3(CH_2)_5$	4-[ $(C_2H_5)_2N(CH_2)_2OCO$ ] $C_6H_4$		508
$CH_3(CH_2)_5$	Various radicals		508
$-(CH_2)_7-$	$C_6H_5$		354
$-CH_2C(C_2H_5)_2CH_2-$	$C_6H_5$	67	439
$-CH_2C(CH_3)(C_3H_7)CH_2-$	$C_6H_5$		257
$-CH_2C(CH_3)(C_3H_7)CH_2-$	4-( $CH_3O$ ) $C_6H_4$		257
$-CH_2C(CH_3)(C_3H_7)CH_2-$	4-( $C_2H_5O$ ) $C_6H_4$		257
$CH_3(CH_2)_6$	4-[ $(C_2H_5)_2N(CH_2)_2OCO$ ] $C_6H_4$		508
$CH_3(CH_2)_6$	Various radicals		508
$-(CH_2)_8-$	$C_6H_5$		354
$CH_3(CH_2)_7$	4-[ $(C_2H_5)_2N(CH_2)_2OCO$ ] $C_6H_4$		508
$CH_3(CH_2)_7$	Various radicals		508
$2,3,5-(CH_3)_3C_6H_2$	$C_6H_5$		421
$-(CH_2)_9-$	$C_6H_5$		354
1-Naphthyl	1-Naphthyl		421
$-(CH_2)_{10}-$	$C_6H_5$		354
$CH_3(CH_2)_9$	4-[ $(C_2H_5)_2N(CH_2)_2OCO$ ] $C_6H_4$		508
$CH_3(CH_2)_9$	Various radicals		508
4-[ $CH_3(CH_2)_8OCO$ ] $C_6H_4$	2-( $CH_3O$ ) $C_6H_4$	71	425
4-[ $CH_3(CH_2)_8OCO$ ] $C_6H_4$	Various radicals	72-98	425
1,4-[ $(CH_2)_8OCO$ ] $_2C_6H_4$	$C_6H_5$		354
3-Cholesteryl	$C_6H_5$	85.2	413
3-Cholesteryl	4-( $H_2O_3As$ ) $C_6H_4$		433
Various aliphatic radicals	4-( $H_2O_3Sb$ ) $C_6H_4$		288
Various aliphatic radicals	1-Naphthyl		683
Various aliphatic radicals	2-Naphthyl		683

TABLE IX (Continued)


R	R'	Yield, %	Reference
Various aliphatic and cycloaliphatic radicals	4-[ $(C_2H_5)_2N(CH_2)_2OCO]C_6H_4$	80-99	566
Various radicals	Various radicals containing the $AsO_3H_2$ substituent		1
	2. Secondary Amines $R'R''NH$		
	R'	R''	Reference
$Cl(CH_2)_2$	$C_6H_5$	$CH_3$	155
$Cl(CH_2)_2$	$C_6H_5$	$C_6H_5$	509
$CH_3O(CH_2)_2$	$2-(CH_3)C_6H_4$	$C_2H_5$	199
$CH_2=C=CHCH_2$	$C_6H_5$	$CH_3$	105, 106
3-Cholesteryl	$C_6H_5$	$CH_3$	454



Trialkylamines react with chloroformates at higher temperatures to yield urethans; presumably the complex salts are intermediates. This reaction proceeds



in high yields with trimethyl- and triethylamines (52, 103, 335). Suitably activated benzyl tertiary amines have given a slightly different reaction with ethyl chloroformate (409). Dimethylbenzylamine did not react




under the same conditions which indicates the need for an electron-donating group on the ring of the benzylamine. However, more recent work has uncovered other examples of the debenzylation of tertiary amines with ethyl chloroformate (739).

Complexes of chloroformates with dimethylaniline have not been studied extensively (590). It has been reported that the complex between phenyl chloroformate and dimethylaniline decomposes at 100° to yield phenol, carbon dioxide, and dimethylaniline.


### b. Reaction of Tertiary Amine-Chloroformate Complexes with Hydroxylic Compounds

It has been reported in the literature that the thermal


reaction of pyridine with an aryl chloroformate leads directly to a diaryl carbonate (449, 484). Work has shown that water is a necessary reactant for this reaction to proceed (47). The complex of pyridine and phenyl chloroformate can be heated at 130° in chlorobenzene with only traces of diphenyl carbonate being formed. Reaction of this complex with water at room



temperature, however, gave quantitative yields of diphenyl carbonate. Similarly, reaction of these pyridine-chloroformate complexes with alcohols and phenols gave the expected carbonates in high yields (123). Although complexes derived from the other types of tertiary amines have not been studied in any detail, they should behave similarly.



One difference that has been noted, however, is the reported (69, 543) preparation of pyrocarbonates by treatment of chloroformates with aqueous potassium hydroxide in the presence of the alkaloid amine, emetine. Other amines were tried in this reaction but none was as useful as emetine.




### 5. Heterocyclic Amines

Chloroformates have been treated with many nitrogen-containing heterocycles to yield urethans. Inorganic base was used as the acid acceptor. Table X is illustrative of the various types of chloroformates and heterocycles that have been used.

TABLE X  
REACTIONS OF CHLOROFORMATES,  $R(OCOCl)_n$ , WITH  
HETEROCYCLED AMINES,  $R'R''NH$


R	$R'R''NH$	Yield, %	Reference
$CH_3$	Benzopyrazole	90	24
$CH_3$	Carbazol		424
$CH_3CH_2$	Imidazol	80-85	649
$CH_3CH_2$	2,4-Diamino-6-methyl-s-triazine	32	192
$CH_3CH_2$	Piperidine		627
$CH_3CH_2$	Benzopyrazol	97	24
$CH_3CH_2$	2,4-Diamino-6-phenyl-s-triazine	77	192
$CH_3CH_2$	Carbazol		424
$CH_3CH_2$	Various aminopyrimidines		259
$ClCH_2C\equiv CCH_2$	2-Aminopyridine	11	330
$ClCH_2C\equiv CCH_2$	Various heterocyclic amines	17-66	330
$-CH_2CH=CHCH_2-$	Piperidine		276
$(CH_3)_2CH(CH_2)_2$	Benzopyrazol	90	24
$C_6H_5$	Benzopyrazol	86	24
$C_6H_5$	Various aminopyrimidines		259
$C_6H_5CH_2$	Benzopyrazol	83	24

Urethans prepared from imidazole and chloroformates are very reactive acylating agents (649). These




compounds have been used in the synthesis of monomers and polymers by their further reaction with compounds containing active hydrogen. The use of N-tri-

methylsilyl derivatives of nitrogen-containing heterocycles for reaction with chloroformates has been reported to give better yields and reaction rates (64).




### 6. Amino Alcohols

Chloroformates react with primary or secondary amino alcohols at the amino group more rapidly than at the hydroxyl group (127, 510, 511). Examples are shown in the following equation and in Table XI. Acid



acceptors can be used in this reaction. Hydroxyurethans of this type can be cyclized with base to yield 2-oxazolidones (5). Further reaction with additional chloroformate leads to acylation of the hydroxyl group.



When the amino group of the amino alcohol is tertiary, the hydrochloride of the aminocarbonate is isolated




TABLE XI  
REACTIONS OF CHLOROFORMATES,  $ROCOCl$ , WITH AMINO ALCOHOLS,  $R_1R_2NR_3OH$

R	$R_1R_2NR_3OH$	Acylation site	Yield, %	Reference
$Br_3CCH_2$	$(CH_3)_2N(CH_2)_2OH$	O		101
$CH_3CH_2$	$H_2N(CH_2)_2OH$	N	79	597
$CH_3CH_2$	2-Hydroxycyclohexylamine	N	75	597
$CH_3CH_2$	$H_2NCHR'CHR''OH$ ( $R'$ and $R''$ are various alkyl radicals)	N	Varied	510, 511
$CH_3CH_2O(CH_2)_2$	$H_2N(CH_2)_2OH$	O and N		525
6-Bromo-2-naphthyl	$(CH_3)_2N(CH_2)_2OH$	O	82	570
2-Naphthyl	$(CH_3)_2N(CH_2)_2OH$	O	80	570
3-Methyl-6-isopropylphenyl	$(C_2H_5)_2N(CH_2)_2OH$	O		742
Various aliphatic and aromatic radicals	$CH_2(CH_2)_3NCH_2CH(C_6H_5)OH$	O	Varied	294
Various aliphatic and aromatic radicals	$(CH_2)_2O(CH_2)_2NCH_2CH(C_6H_5)OH$	O	Varied	294
Various aliphatic and aromatic radicals	$CH_2(CH_2)_4NCH_2CH(C_6H_5)OH$	O	Varied	294
Various aliphatic and aromatic radicals	$(CH_2)_2N(CH_3)(CH_2)_2NCH_2CH(C_6H_5)OH$	O	Varied	294
$-(CH_2CH_2O)_n-$	$CH_3(CH_2)_{11}N(CH_3)(CH_2CH_2O)_nH$	O		127

(101, 294, 570, 742). Generation of the free amine can be accomplished by reaction of the hydrochloride with base. Nonionic detergents have been prepared from polyethylene glycol bis(chloroformates) and long-chain tertiary amino alcohols (127).

### 7. Aminophenols

As with amino alcohols, the amino group of aminophenols is more reactive toward a chloroformate (568). An example is shown and others are tabulated in




Table XII. With *o*-aminophenol, benzoxazolones are produced via cyclization of the intermediate phenolic urethan (567).




TABLE XII

REACTIONS OF CHLOROFORMATES,  $ROCOCl$ , WITH AMINOPHENOLS,  $H_2NArOH$

R	ArOH	Yield, %	Reference
$CH_3$	4-Hydroxyphenyl	572	
$CH_3CH_2$	4-Hydroxyphenyl	572	
$CH_3CH_2$	2-Hydroxyphenyl	572	
$CH_3(CH_2)_2$	4-Hydroxyphenyl	572	
$CH_3(CH_2)_3$	4-Hydroxyphenyl	572	
$CH_3(CH_2)_3$	2-Hydroxyphenyl	572	
$CH_3CH_2CH(CH_3)$	4-Hydroxyphenyl	572	
$CH_3(CH_2)_4$	4-Hydroxyphenyl	572	
$C_6H_5$	3,5-Dichloro-4-hydroxyphenyl	64	572
$CH_3(CH_2)_6$	2-Hydroxyphenyl	572	
$C_6H_5CH_2$	4-Hydroxyphenyl	572	
$C_6H_5CH_2$	2-Hydroxyphenyl	572	
$CH_3(CH_2)_6$	4-Hydroxyphenyl	93	573
$CH_3(CH_2)_6$	2-Hydroxyphenyl	573	
$C_6H_5(CH_2)_2$	4-Hydroxyphenyl	95	573
$CH_3(CH_2)_7$	4-Hydroxyphenyl	83	573
$CH_3(CH_2)_7$	2-Hydroxyphenyl	573	
$CH_3(CH_2)_9$	4-Hydroxyphenyl	86	573
$CH_3(CH_2)_{13}$	4-Hydroxyphenyl	51	573
Various aryl radicals	2-Hydroxyphenyl	88	567


### 8. Amino Acids

The reaction of chloroformates with amino acids has been used for the protection of the amine group during peptide synthesis (Table XIII). In particular, benzyl

TABLE XIII  
REACTIONS OF CHLOROFORMATES,  $R(OCOCl)_n$ , WITH AMINO ACIDS,  $NH_2R'H$

R	R'H	Yield, %	Reference
$CH_3$	Various aminoaryl arsonic acids	65-91	287, 460, 461
$C_2H_5$	Various aminoaryl arsonic acid	59-90	287, 460, 461
$C_3H_7$	Various aminoaryl arsonic acids	35-90	287, 460, 461
$(CH_3)_2CH$	Various aminoaryl arsonic acids	45-85	287, 460, 461
$XCH_2C\equiv CCH_3$ (X is a halogen)	$CH_2COOH$	26	330
$XCH_2C\equiv CCH_2$ (X is a halogen)	Various radicals	11-96	330
$-CH_2CH=CHCH_2-$	$CH_2COOH$	276	
$-CH_2CH=CHCH_2-$	$(CH_3)_2COOH$	276	
$-CH_2CH=CHCH_2-$	$4-(HOCO)C_6H_4$	276	
$CH_3(CH_2)_3$	Various radicals	40-90	287, 460, 461
$(CH_3)_2CHCH_2$	Various radicals	50-90	287, 460, 461
$2-(NO_2)C_6H_4$	$CH(CH_3)COOH$	849	
$4-(Br)C_6H_4CH_2$	$CH_2COOH$	119	
$4-(NO_2)C_6H_4CH_2$	$CH_2COOH$	78	107
$C_6H_5CH_2$	$CH_2COOH$	90	112
$C_6H_5CH_3$	$CH(CH_3)(CH_2)_{10}COOH$	350	
$C_6H_5CH_3$	$CH(CH_3)(CH_2)_{11}COOH$	38	350
$C_6H_5CH_3$	Various radicals	Varied	58, 63, 223, 542
$4-(CH_3)C_6H_4$	$-CH_2COOH$	119	
$4-(CH_3)C_6H_4$	$-CH(CH_3)COOH$	849	
1-Naphthyl	$-CH_2COOH$	119	
2-Naphthyl	Various radicals	40-90	734
Various aliphatic and aromatic radicals	Various degraded peptides		646

chloroformate has been preferred for this purpose. The



derived urethan can be hydrogenolyzed easily to free the amine after the carboxyl group has reacted further (58, 223, 350, 542). The above equation illustrates the greater reactivity of the amine group toward the chloroformate than the acid group in amino acids. A similar reactivity was found with *p*-aminophenyl-arsinic acid (217).


This reaction has found utility for the resolution of optically active alcohols (6). A racemic alcohol is converted to its chloroformate ester which is then treated with an optically active amino acid. The resulting urethan-acid was then separated into its antipodes, followed by regeneration of the original alcohol.

No examples have been found in the literature where the urethan acid has been reacted further with another mole of chloroformate to yield the ester or the mixed carboxylic-carbonic anhydride. This reaction should be possible in view of the recent synthesis and isolation of mixed carboxylic-carbonic anhydrides. A patent


describing the preparation of useful surface active agents by the reaction of chloroformates with amino acids or partially degraded peptides has been issued (646).

### 9. Amides


Only a few examples of the reaction of a chloroformate with an amide have appeared in the literature. Apparently, either O- or N-acylation can occur depending on the specific reactants. Formamide reacted with ethyl chloroformate to yield tris(formylamino)methane (X) in 31% yield along with other products (87). The initial step was postulated to be O-acylation followed by reaction along the paths shown in the diagram.




Hall has found that chloroformates form conducting solutions in dimethylformamide (281). This observation was attributed to the following reaction for which he has obtained additional evidence based on the reactivity of these complexes.




An intramolecular assist by an amide group has been postulated to aid the decomposition of a chloroformate prepared from a  $\beta$ -hydroxyethylamide (55).



Trichloromethyl chloroformate has been used to dehydrate amides to nitriles on simple heating in an inert solvent (303).



Examples of the acylation of a sulfonamide have appeared recently (22).



Additional examples are listed in Table XIV.

TABLE XIV  
REACTIONS OF CHLOROFORMATES,  $\text{ROCOCl}$ , WITH  
SULFONAMIDES,  $\text{R}'\text{SO}_2\text{NH}_2$

R	$\text{R}'$	Yield, %	Reference
$\text{CH}_3$	Cyclohexyl		23
$\text{C}_2\text{H}_5$	$\text{CH}_3$		114
$\text{C}_2\text{H}_5$	2-Thienyl		114
$\text{C}_2\text{H}_5$	$\text{CH}_3(\text{CH}_2)_3$		114
$\text{C}_2\text{H}_5$	Cyclopentyl		114
$\text{C}_2\text{H}_5$	$\text{CH}_3(\text{CH}_2)_4$		114
$\text{C}_2\text{H}_5$	Cyclohexyl		114
$\text{C}_2\text{H}_5$	$\text{CH}_3(\text{CH}_2)_5$		114
$\text{C}_2\text{H}_5$	$4-(\text{CF}_3)\text{C}_6\text{H}_4$		552
$\text{C}_2\text{H}_5$	$2,4-(\text{NO}_2)(\text{CF}_3)\text{C}_6\text{H}_3$		552
$\text{C}_2\text{H}_5$	$4-(\text{COH})\text{C}_6\text{H}_4$		482
$\text{C}_2\text{H}_5$	$\text{C}_6\text{H}_5\text{CH}_2$		114
$\text{C}_2\text{H}_5$	$4-(\text{CH}_3\text{CONHCH}_2)\text{C}_6\text{H}_4$		482
$\text{C}_2\text{H}_5$	$4-[CH_3\text{CONH}(\text{CH}_2)_2]\text{C}_6\text{H}_4$		482
$\text{C}_2\text{H}_5$	$4-[CH_3\text{CONHCH}(\text{CH}_3)]\text{C}_6\text{H}_4$		482
$\text{C}_2\text{H}_5$	$\text{CH}_3(\text{CH}_2)_9$		114
$\text{C}_2\text{H}_7$	$\text{CH}_3(\text{CH}_2)_3$	60-80	{ 114
$\text{CH}_3\text{O}(\text{CH}_2)_2$	$4-(\text{CH}_3)\text{C}_6\text{H}_4$	22	
$\text{CH}_3(\text{CH}_2)_3$	$\text{CH}_3(\text{CH}_2)_3$		114
$\text{CH}_3(\text{CH}_2)_3$	$\text{CH}_3(\text{CH}_2)_4$		114
$\text{CH}_3\text{CH}(\text{C}_2\text{H}_5)$	$\text{CH}_3(\text{CH}_2)_4$		114
$\text{CH}_3(\text{CH}_2)_4$	$\text{CH}_3(\text{CH}_2)_3$		114
$\text{CH}_3(\text{CH}_2)_5$	$\text{CH}_3(\text{CH}_2)_3$		114
$\text{CH}_3(\text{CH}_2)_5$	Cyclohexyl		23


### 10. Amidoximes

Amidoximes are O-acylated by chloroformates (Table XV) in high yields. An equivalent of the amidoxime

TABLE XV  
REACTIONS OF ETHYL CHLOROFORMATE WITH AMIDOXIMES,  
 $\text{R}'\text{C}(\text{NH}_2)=\text{NOH}$

$\text{R}'$	Reference
3-Pyridyl	474
$4-(\text{NO}_2)\text{C}_6\text{H}_4$	706
$3-(\text{NO}_2)\text{C}_6\text{H}_4$	625
$\text{C}_6\text{H}_5$	4, 203, 204
$2-(\text{HO})\text{C}_6\text{H}_4$	477
$4-(\text{CH}_3)\text{C}_6\text{H}_4$	629
$4-(\text{CH}_3\text{O})\text{C}_6\text{H}_4$	624
$\text{C}_6\text{H}_5\text{CH}(\text{OH})$	271
$2,4-(\text{CH}_3)_2\text{C}_6\text{H}_3$	535
6-Quinolyl	62
1-Naphthyl	588
2-Naphthyl	588
Various radicals	678, 679


can act as the hydrogen chloride acceptor although aqueous base or organic amines can serve the same purpose (203, 563, 669, 735). Amidoximes derived from




tertiary amides and bis(amidoximes) react similarly (83, 748). The products are very useful intermediates for the preparation of various types of heterocyclic ring systems.

### 11. Urethans

Heating chloroformates prepared from  $\beta$ -hydroxyethylurethans gives 2-oxazolidones in excellent yields (55).




Acylation of ammonia with a chloroformate was described earlier. Further acylation with additional chloroformate is readily accomplished by using the sodium salt of the urethan as shown in the following equations (176, 407, 741). The use of different chloro-




formates at the different steps has provided mixed urethans. This reaction has also been carried out on the sodio derivatives of phenylurethan and the ethylurethan of glycine (177, 608, 636, 651).

The reaction of N-chloroethylurethan with ethyl chloroformate in the presence of potassium hydroxide has been reported to yield N,N-dichloroethylurethan and diethyl azamalonate (681). A proposed reaction scheme is shown.




Hydrazine reacts with chloroformates to yield the expected hydrazides, depending on whether 1 or 2 moles of chloroformate is employed (162, 163, 274).




Cyclization of the hydrazide formed from the chloro-

formate of 2-chloroethanol with sodium ethoxide offers a route to N-amino-2-oxazolidones (172).




1,1-Bis(2-cyanoethyl) hydrazine reacted with ethyl chloroformate to yield a pyrazolidine (XI) in addition to low yields of the expected carbazate and carbohydrazide (442).




### 13. Hydrazides


A series of bis(hydrazodicarboxylates) has been prepared from aliphatic and aromatic bis(chloroformates) by reaction with hydrazides (565, 724).




An acid acceptor was used. The formyl hydrazide shown below was acylated with methyl chloroformate (717).



O-Acylation occurred by reaction of maleic hydrazide with ethyl chloroformate to yield XII (652). N-Acy-




ation has been observed with the cyclic hydrazide XIII to yield the depicted products, which have utility as color couplers in color photography (736).




#### 14. Cyanamid

Both hydrogens of cyanamid have been acylated by chloroformates in high yields (40, 179, 368, 463). Polymerizable derivatives of cyanamid have been prepared by using allyl chloroformate in this reaction (85, 258).




Dicyandiamide has been monoacylated (70–95% yields) under similar conditions (7, 560).



These products have been described as possessing a variety of uses.


#### 15. Urea

Urea has been monoacylated by a chloroformate (165, 610, 718).




#### 16. Guanidine

Guanidine has been diacylated with ethyl chloroformate (522).




#### 17. Thioureas

Thiourea has been reported to react with chloroformates as shown (181).



In the presence of an acid acceptor, substituted thio-

ureas have yielded carbodiimides in fair yields on reaction with a chloroformate (144).




#### 18. Miscellaneous Nitrogen-Containing Compounds

A series of Swiss patents describe several uses for the water-soluble products obtained from reaction of various chloroformates with imidazolsulfonic acids (*i.e.*, sodium hendecylbenzimidazolsulfonic acid). Foaming agents and textile-dyeing assistants are among the uses claimed (640–645).

#### E. MISCELLANEOUS REACTIONS AND SYNTHESES

##### 1. Sulfinic Acids

Very little research has been done on this reaction (537). The reaction of sodium *p*-toluenesulfinate with *d*-2-octyl chloroformate yielded alkyl *p*-toluenesulfinates (335, 336). This result is in contrast to the fact that sulfones are obtained by reaction of alkyl halides with sodium *p*-toluenesulfinate. Hydrolysis of the sulfinic ester obtained from this reaction gave *d*-2-



octanol of retained configuration. This fact was interpreted as evidence that an intermediate anhydride was formed which decomposed without alkyl-O cleavage, resulting in retention of configuration.

##### 2. Active Methylene Compounds

There are many reported examples (Table XVI) of the acylation of various types of active methylene compounds by chloroformates. Apparently much of this work was done quite early, so that the poor yields that have been reported for these reactions might be increased by using modern techniques and condensing agents. Dimethyl malonate has been converted to the tetracarboxylate by this method (135, 136, 153, 517, 518, 622).







TABLE XVI  
REACTIONS OF CHLOROFORMATES,  $\text{ROCOCl}$ , WITH ACTIVE  
METHYLENIC COMPOUNDS,  $\text{R}'\text{H}$

$\text{R}$	$\text{R}'\text{H}$	Site of carboxylation	Yield, %	Reference
$\text{CH}_3$	5,5-Dimethyl-1,3-cyclohexanedione	O		633
$\text{CH}_3$	Dibenzoylmethane	C and O	Varies	473
$\text{C}_2\text{H}_5$	Ethyl acetate	C		606
$\text{C}_2\text{H}_5$	Ethyl cyanoacetate	C		283, 284, 302
$\text{C}_2\text{H}_5$	Dimethyl malonate	C		136, 153
$\text{C}_2\text{H}_5$	Cyclohexanone	O		285, 286
$\text{C}_2\text{H}_5$	Ethyl acetoacetate	C and O		153, 470, 471, 472, 606, 725
$\text{C}_2\text{H}_5$	Ethyl bis(methoxycarbonyl)acetate	C		153, 517, 518, 622
$\text{C}_2\text{H}_5$	Propiophenone	O		285, 286
$\text{C}_2\text{H}_5$	Butyrophenone	O		285, 286
$\text{C}_2\text{H}_5$	Isobutyrophenone	O		285, 286
$\text{C}_2\text{H}_5$	Ethyl phenylacetate	C		607
$\text{C}_2\text{H}_5$	Ethyl 2-(phenyl)butyrate	C		703

Acetoacetates are converted to acetyl malonates (137, 139, 153, 470, 471, 725).




Cyanoacetates have been converted to cyanomalonates (283, 284, 302).  $\alpha$ -Aryl acetates have also been acylated by chloroformates using potassium or sodamide as catalysts (473, 607, 703). Mixtures of products




and low yields have been reported.

O-Acylation has been observed in this reaction. Dibenzoylmethane gave a mixture of the C- and O-acylated products on treatment with methyl chloroformate in the presence of sodamide (473). In some cases, O-acylation is the main reaction, as with the  $\beta$ -diketone, dimedon (633).





Cyclohexanone reacted similarly in the presence of sodamide (285, 286).



A series of phenyl alkyl ketones gave O-acylated products on reaction with ethyl chloroformate (285, 286).

The copper(II) chelate of ethyl acetoacetate has been treated with ethyl chloroformate to yield the analogous chelate of acetylmalonic ester (472). Similar results were obtained with the Cu(II) chelate of benzoylacetic ester which yielded the expected benzoyl malonate. The following two reactions illustrate cyclizations of  $\beta$ -dicarbonyl compounds through C- and O-acylations.





$\text{R} = \text{C}_6\text{H}_5; 29\%$   
 $\text{R} = \alpha\text{-C}_10\text{H}_7; 51\%$

### 3. Organometallic Compounds


Many diverse types of organometallic compounds react with chloroformates. They are shown in Table XVII.

Organometallic compounds of the Grignard type react with chloroformates to yield esters (333, 410, 452, 571). However, with the more reactive organometallic compounds, the initially formed ester reacts further to yield the tertiary alcohol. Usually both products are



formed although their relative amounts can be controlled by the molar amounts of the starting materials that are used and by the reaction conditions employed.

Acetylenic Grignard reagents or alkali metal acetylides yield the carboxylic ester with little or no tertiary carbinol being formed (190, 485, 486, 534). However,




the reaction of ethyl chloroformate with the lithium derivative of ethoxyacetylene gave only a small yield of the expected ester. Surprisingly, the lithium derivatives of the sulfur analog of ethoxyacetylene gave a 56% yield of the expected ester in the same reaction (13).

3-Thianaphenylmagnesium chloride reacted


TABLE XVII  
REACTIONS OF CHLOROFORMATES, ROCOCl, WITH ORGANOMETALLIC DERIVATIVES

R	Organometallic Derivative	Product(s)	Yield, %	Reference
CH <sub>3</sub>	(CH <sub>3</sub> ) <sub>2</sub> Zn			94
CH <sub>3</sub>	(CH <sub>3</sub> ) <sub>2</sub> CHMgCl			571
C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub> MgBr			410, 452
C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub> MgI			452
C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub> C≡CMgBr			190, 485, 486
C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub> C≡CNa			190, 485, 486
C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> =CHCH <sub>2</sub> MgBr			410
C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> =CHCH <sub>2</sub> ZnI	(CH <sub>2</sub> =CHCH <sub>2</sub> ).COH	40	92, 574, 575
C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub> SC≡CLi	C <sub>2</sub> H <sub>5</sub> SC≡CCOOC <sub>2</sub> H <sub>5</sub>	56	13
C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>3</sub> MgBr			410
C <sub>2</sub> H <sub>5</sub>	(CH <sub>3</sub> ) <sub>2</sub> NCH <sub>2</sub> C≡CNa	(CH <sub>3</sub> ) <sub>2</sub> NCH <sub>2</sub> C≡CCOOC <sub>2</sub> H <sub>5</sub>	24	534
C <sub>2</sub> H <sub>5</sub>	C <sub>6</sub> H <sub>5</sub> MgBr	C <sub>6</sub> H <sub>5</sub> COOC <sub>2</sub> H <sub>5</sub>	16-75	
		(C <sub>6</sub> H <sub>5</sub> ) <sub>3</sub> COH	9-34	333
C <sub>2</sub> H <sub>5</sub>	[C <sub>6</sub> H <sub>5</sub> Br + NaHg]			102, 740
C <sub>2</sub> H <sub>5</sub>	C <sub>6</sub> H <sub>5</sub> P(MgBr) <sub>2</sub>			357
C <sub>2</sub> H <sub>5</sub>	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>3</sub> C≡CNa	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub> C≡CCOOC <sub>2</sub> H <sub>5</sub>	23	534
C <sub>2</sub> H <sub>5</sub>	C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> MgBr	C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> COOC <sub>2</sub> H <sub>5</sub>	43	333
C <sub>2</sub> H <sub>5</sub>	C <sub>6</sub> H <sub>5</sub> CH(MgCl)COONa	(C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> ) <sub>3</sub> COH		650
C <sub>2</sub> H <sub>5</sub>	1,4-(CH <sub>3</sub> )(SO <sub>2</sub> CH <sub>2</sub> MgBr)C <sub>6</sub> H <sub>4</sub>	1,4-(CH <sub>3</sub> )(SO <sub>2</sub> CH <sub>2</sub> )C <sub>6</sub> H <sub>4</sub>	70	226
C <sub>2</sub> H <sub>5</sub>	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>5</sub> C≡CNa	1,4-(CH <sub>3</sub> )(SO <sub>2</sub> CH <sub>2</sub> COOC <sub>2</sub> H <sub>5</sub> )C <sub>6</sub> H <sub>4</sub>	22	534
C <sub>2</sub> H <sub>5</sub>	2-Methyl-2,5-(dimethylmethano)cyclohexyl-magnesium chloride	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>5</sub> C≡CCOOC <sub>2</sub> H <sub>5</sub>	53	591, 700
C <sub>2</sub> H <sub>5</sub>	(C <sub>6</sub> H <sub>5</sub> ) <sub>2</sub> PMgBr			357
C <sub>2</sub> H <sub>5</sub>	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>9</sub> C≡CNa	(CH <sub>3</sub> ) <sub>2</sub> N(CH <sub>2</sub> ) <sub>9</sub> C≡CCOOC <sub>2</sub> H <sub>5</sub>		534
C <sub>2</sub> H <sub>5</sub>	RC≡CMgBr (R are various aliphatic radicals)			190, 485, 486
C <sub>2</sub> H <sub>5</sub>	RC≡CNa (R are various aliphatic radicals)			190, 485, 486


with ethyl chloroformate with rearrangement to yield the 2-carboxylate (243). None of the expected ester was found in the reaction product.




Half-esters of malonic acid have been prepared by the following reaction (650).




Reaction of alkylzinc compounds with chloroformates also gives tertiary alcohols but to a lesser degree (11, 91, 94, 574, 575). Heating ethyl chloroformate with zinc alone resulted in extensive decomposition (250).




A Reformatsky-type reaction has been observed in the reaction of ethyl chloroformate with ethyl acetoacetate in the presence of zinc and reduced copper powder (11).



The phosphorus-containing Grignard reagent in the following equation reacted to yield the expected ester (357).



The reaction of sodium with ethyl iodide and ethyl chloroformate led to the mixture of products shown (250).




#### 4. Inorganic Salts


A comparative rate study of the reaction of various anions with ethyl chloroformate gave the following reactivity order (265): acetoxime pH (6 or 7) > OH<sup>-</sup> > C<sub>6</sub>H<sub>5</sub>OH (pH 7) > NO<sub>2</sub><sup>-</sup> > N<sub>3</sub><sup>-</sup> > S<sub>2</sub>O<sub>3</sub><sup>-2</sup> > F<sup>-</sup>. In the reaction medium used (85:15 water-acetone), chloride, bromide, iodide, cyanate, thiocyanate, chlorate, and nitrate ions had no effect on the rate of reaction. The possible reasons for this entirely different rate order for the reaction of anions with ethyl chloro-

formate (acylation) as compared to their reactivity in alkylations was discussed. The rate of reaction of ethyl chloroformate with sodium iodide in acetone was studied by Conant in his classical study of the reactivity of organic chlorides (151, 152).

Reaction of sodium iodide with chloromethyl chloroformates led to the decomposition reactions shown in the following equations (550, 551).




Alkali metals, especially, sodium, decomposed ethyl



chloroformate to yield diethyl carbonate and carbon monoxide (250, 718).


Two early reports described the conversion of alkyl chloroformates to the corresponding carbonates by lead oxide and water (159, 457). Water was thought to be necessary for the reaction to occur.

Alkylmetal sulfides and polysulfides react with chloroformates to yield bis(carbalkoxy) sulfides (326, 469, 686). This class of compound decomposes on




heating to yield the carbonate, carbonoxysulfide, and sulfur.

The reaction of chloroformates with inorganic fluorides yields the corresponding fluoroformates (512).




Thallous fluoride is a preferred reagent for this conversion, although conditions for the use of potassium fluoride have also been found (382, 531, 707). The facile catalyzed decomposition of fluoroformates to alkyl fluorides represents a convenient synthesis of the latter type of compound.


Silver nitrate has been treated with alkyl chloroformates to yield nitrate esters (68, 76). This reaction can be run in either heterogeneous or homogeneous systems and possesses advantages for the preparation of nitrates




when oxidizable groups are present. In addition, it was shown by mechanism studies that this reaction proceeded mainly with retention of the configuration of the R group. Hence, it was postulated that an SNI-type decomposition of the intermediate was operative. Some reaction was thought to occur by the mechanism shown in the following equation. An attempt to pre-



pare phenyl nitrate from phenyl chloroformate and silver nitrate resulted in a 64% yield of *o*-nitrophenol (118).




The reaction of silver hyponitrite with chloroformates has been described as leading to the formation of hyponitrous esters (631). The products, though not isol-




able, were found to be useful as polymerization catalysts for olefins. Their decomposition products are nitrogen, carbon dioxide, nitrous oxide, and free radicals, the latter functioning as polymerization initiators.


An early attempt to react potassium cyanate with ethyl chloroformate in the presence of ethanol gave a variety of products (718). Somewhat later, it was



found that simple heating of ethyl chloroformate with potassium cyanate gave some carbethoxyethyl isocyanate with recoverable chloroformate (178, 250, 251, 720).




Cyanoformates have been prepared from chloroformates by reaction with sodium cyanide or hydrogen cyanide-pyridine in 5–40% yields. Pyrolysis of the



cyanoformates at 700° yielded the corresponding nitrile (40–60%) in most cases.


Magnesyl selenohydrate (from ethylmagnesium bromide and hydrogen selenide) reacted with ethyl chloroformate to give the following products (478, 479). Pure




compounds were not isolated in high yields in this work.

### 5. In the Presence of Lewis Acids

Aliphatic chloroformates undergo decomposition with Lewis acids according to the following equation (91, 93, 94, 513, 577, 689, 690). Yields of the products



vary according to particular reactants, reagents, and reaction conditions. Chlorinated chloroformates yield the parent chloroalkane in many cases. Some examples are shown below. In some cases, free phosgene




has been detected among the decomposition products. Table XVIII lists other examples of reactions of Lewis acids with chloroformates.

TABLE XVIII  
REACTIONS OF CHLOROFORMATES,  $\text{ROCOCl}$ , WITH LEWIS ACIDS


R	Lewis acid	Products (yield, %)	Reference
$\text{Cl}_2\text{C}$	$\text{AlCl}_3$	$\text{CO}_2, \text{CCl}_4$	269, 306
$\text{ClICH}_2$	$\text{AlCl}_3$	$\text{COCl}_2$ , other products	390
$\text{ClICH}_2$	$\text{FeCl}_3$	$\text{COCl}_2$ , other products	390
$\text{Cl}_2\text{CCCl}_2$	$\text{AlCl}_3$	Hexachloroethane, $\text{CO}_2$	489
$\text{Cl}_2\text{CHCCl}_2$	$\text{AlCl}_3$	Tetrachloroethylene, $\text{CO}_2$ , $\text{HCl}$	489
$\text{ClICH}_2\text{CCl}_2$	$\text{AlCl}_3$	<i>sym</i> -Tetrachloroethane, trichloroethylene, $\text{CO}_2$ , $\text{HCl}$	489
$\text{ClICH}_2\text{CHCl}$	$\text{AlCl}_3$	1,1,2-Trichloroethane, $\text{CO}_2$	489
$\text{CH}_3\text{CHCl}$	$\text{AlCl}_3$	$\text{CH}_3\text{CHCl}_2$ , $\text{CH}_2=\text{CHCl}$ , $\text{CO}_2$	489
$\text{C}_2\text{H}_5$	$\text{AlBr}_3$	$\text{C}_2\text{H}_5\text{Cl}, \text{C}_2\text{H}_5\text{Br}, \text{CO}_2$	403
$\text{C}_2\text{H}_5$	$\text{AlCl}_3$	$\text{C}_2\text{H}_5\text{Cl}, \text{CO}_2$	577
$\text{C}_2\text{H}_5$	$\text{BF}_3\text{-ether}$	$\text{C}_2\text{H}_5\text{Cl}$ (62), $\text{CO}_2$ (64)	513
$\text{C}_2\text{H}_5$	$\text{ZnCl}_2$	$\text{C}_2\text{H}_5\text{Cl}$ (26), ethylene (18), $\text{HCl}, \text{CO}_2$ , 449, 689, 690	91, 98, 94, (18), HCl, CO <sub>2</sub> , 449, 689, 690
$(\text{ClICH}_2)(\text{Cl}_2\text{CH})\text{CH}$	$\text{FeCl}_3$	$\text{CO}_2, \text{HCl}$ , carbon	446
$(\text{Cl}_2\text{CH})\text{CH}$	$\text{AlCl}_3$	$\text{COCl}_2, \text{HCl}$ , carbon	446
$n\text{-C}_2\text{H}_7$	$\text{ZnCl}_2$	$n\text{-C}_2\text{H}_7\text{Cl}$ (23), propylene (55), $\text{HCl}, \text{CO}_2$	690
Various aliphatic radicals	$\text{BF}_3\text{-ether}$	Alkyl chloride, olefins, $\text{HCl}$	513

Reaction of aliphatic chloroformates with Lewis acids in the presence of an aromatic hydrocarbon results in alkylation rather than acylation (95, 239–241, 403, 414, 415, 577, 744). Apparently the decomposition reaction occurs first to give the carbonium ion which then alkylates the aromatic ring. A patent claims to have upgraded isobutane to gasoline with high anti-knock value by treating it with alkyl chloroformates and a Friedel-Crafts catalyst (613),

In contrast to the aliphatic chloroformates, aromatic chloroformates are quite stable to Lewis acids (157). Complexes are undoubtedly formed, but extensive decomposition reactions are not observed up to temperatures of 130°. Because of this fact, aromatic chloroformates are good acylating agents under Friedel-Crafts conditions.




The use of hydrogen fluoride as a catalyst in a reaction of a chloroformate has resulted in halogen exchange to yield the fluoroformate (380).



### 6. Thermal Decomposition of Chloroformates

Aliphatic chloroformates decompose thermally by the two paths shown in the following equation (533, 710) (Table XIX). The temperature required for de-



composition varies widely depending on the structure of the aliphatic radical. For example, *t*-butyl chloroformate is unstable at room temperature (133, 464, 465). The most recent studies on the mechanism of this reaction were carried out in the vapor phase at temperatures near 250° (429–432). Path 1 above is thought to proceed through a rather polar transition state of the S<sub>Ni</sub> type (XIV). This mechanism was





TABLE XIX  
THERMAL DECOMPOSITION OF CHLOROFORMATES, R(OCOCl)<sub>n</sub>

R	Product(s)	Yield, %	Reference
Cl <sub>3</sub> C-(CH <sub>2</sub> ) <sub>2</sub> -	COCl <sub>2</sub> , CH <sub>2</sub> =CHOCOCl, (CH <sub>2</sub> Cl) <sub>2</sub> , CH <sub>2</sub> CHCl <sub>2</sub>	100 417	569 417
Cl(CH <sub>2</sub> ) <sub>2</sub> C <sub>2</sub> H <sub>5</sub>	ClCH <sub>2</sub> CH <sub>2</sub> Cl	360	131, 132,
	C <sub>2</sub> H <sub>5</sub> Cl, CH <sub>2</sub> =CH <sub>2</sub> , HCl, CO <sub>2</sub>	429, 513, 546, 592, 710, 718	
CH <sub>2</sub> CH <sub>2</sub> - (CH <sub>2</sub> ) <sub>2</sub> CH	CH <sub>2</sub> CH=CHOCOCl (CH <sub>2</sub> ) <sub>2</sub> CHCl, CO <sub>2</sub> , CH <sub>2</sub> =CHCH <sub>2</sub> , HCl	417 134, 150	
CH <sub>2</sub> CH=CHCH <sub>2</sub>	CH <sub>2</sub> CH=CHCH <sub>2</sub> Cl, CH <sub>2</sub> CH(CH=CH <sub>2</sub> )Cl, (CH <sub>2</sub> =CH) <sub>2</sub>	533	
CH <sub>2</sub> CH(CH=CH <sub>2</sub> )	CH <sub>2</sub> CH=CHCH <sub>2</sub> Cl, CH <sub>2</sub> CH(CH=CH <sub>2</sub> )Cl, (CH <sub>2</sub> =CH) <sub>2</sub>	533	
FCOO(CH <sub>2</sub> ) <sub>2</sub> S(CH <sub>2</sub> ) <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> CHCH <sub>2</sub> (CH <sub>2</sub> ) <sub>2</sub> C	F(CH <sub>2</sub> ) <sub>2</sub> S(CH <sub>2</sub> ) <sub>2</sub> Cl (CH <sub>2</sub> ) <sub>2</sub> C=CH <sub>2</sub> , HCl, CO <sub>2</sub>	22 >98	380 428 133, 464, 465
C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> C <sub>6</sub> H <sub>5</sub> CH(CH <sub>3</sub> ) CH <sub>3</sub> (CH <sub>2</sub> ) <sub>2</sub> CH(CH <sub>3</sub> ) [(CH <sub>2</sub> ) <sub>2</sub> C]CH	C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> Cl C <sub>6</sub> H <sub>5</sub> CH(CH <sub>3</sub> )Cl CH <sub>3</sub> (CH <sub>2</sub> ) <sub>2</sub> CH(CH <sub>3</sub> )Cl [(CH <sub>2</sub> ) <sub>2</sub> C]CHCl	676, 710 710 296, 337 40	379
(C <sub>6</sub> H <sub>5</sub> ) <sub>2</sub> CH (C <sub>6</sub> H <sub>5</sub> ) <sub>2</sub> C 3-Cholesteryl	(C <sub>6</sub> H <sub>5</sub> ) <sub>2</sub> CCl 3-Cholesteryl chloride	50 100	80 80 454, 711 465
Various tertiary alkyl radicals	Corresponding chlorides	710	
Various alkyl- substituted $\alpha$ - phenethyl radicals	Corresponding chlorides	296	
Various long-chain aliphatic radicals	Corresponding chlorides	454, 711	
Various steroidal radicals	Corresponding chlorides		

supported by the relative reactivities of the different alkyl chloroformates ( $3^\circ > 2^\circ > 1^\circ >> C_6H_5$ ), the large negative entropy of activation, the retention of configuration, and the absence of rearranged alkyl chlorides. The elimination reaction (2) was shown to be a completely stereospecific *cis* elimination and the transition state XV was postulated to be operative. Transition state XV was described as leading to an intermediate of substantial charge separation that can react to give different olefins and rearranged products, but does not dissociate enough to give stereochemically random products. The extent of the importance of ion pairs was also discussed.

Benzyl chloroformates decompose quite readily on heating to yield the respective chlorides (80, 676, 710). Presumably the stabilization of the incipient positive charge by the phenyl group (*i.e.*, a benzyl carbonium ion) facilitates their decomposition (21).


Literature reports that aryl chloroformates can be converted to aryl carbonates and phosgene by simply heating are probably not true (37, 38, 80, 308, 556). Recent research has shown that aryl chloroformates




are very stable thermally when pure (418). Apparently, moisture or other catalysts were present when this reaction was reported to occur.

### 7. Peroxides


Metal peroxides or hydrogen peroxide and base react with chloroformates to give peroxy compounds. At low temperatures, the sodium salt of ethyl monoperoxy-carbonic acid (450) was not obtainable. Instead, the following reaction occurred. However, the symmetrical




diethyl ester of peroxydicarbonic acid has been prepared by the reaction shown below (656, 660, 712). Many di-



alkyl peroxides of this type have been reported. Peroxydicarbonates formed *in situ* have been found to be good initiators for vinyl polymerizations (263). An example that used hydrogen peroxide and sodium hydroxide is shown in the following reaction (66).



The reaction of hydroperoxides with chloroformates in the presence of acid acceptors has yielded esters of monoperoxy-carbonic acid (660). With peroxycar-




boxylic acids, the mixed anhydrides were prepared (541).

Compounds which can be considered to be amide derivatives of monoperoxy-carbonic acid (XVI) have




been prepared by the reaction of hydroxylamine, dialkylhydroxylamines, and oximes with chloroformates (85, 749). An example of each type is shown. These




compounds can be analyzed iodometrically because they yield peroxy structures on hydrolysis.

An analytical method for determination of chloroformates is based on their reaction with isonitrosoacetone. Subsequent hydrolysis liberates cyanide ion (603). Quantitative analysis of the cyanide ion is a measure of the chloroformate content.




#### 8. Synthesis of Isothiocyanates

The synthesis of isothiocyanates based on the early observations of Kaluza is a general method (200–202, 370, 383–386, 709). This route, from which high yields can be obtained, is shown in the following equations. The final step was improved by using triethyl-




amine in methylene chloride (244). The kinetics and mechanism of the decomposition of the intermediate carbethoxy dithiocarbamates have been studied (320, 321).

An analogous reaction of dithiocarbamates derived from secondary amines with aryl chloroformates yielded the expected products (369). The products decomposed slowly at room temperature over a period of weeks.



#### 9. Xanthates


Several patents have described the reaction of chloroformates with alkylxanthates to yield alkyl xanthogen formates (74, 75, 228). These products are claimed to



possess utility in the concentration of minerals by flotation processes.


#### 10. Phosphoranes (Wittig Reaction)

The reaction of a chloroformate with phosphoranes led to a new synthesis of substituted acrylates (61).




#### 11. Phosphites and Phosphines


Sodium diethyl phosphite can react with ethyl chloroformate in both of the ways shown in the following equations (12).




Trialkyl phosphites have been treated with bis(chloroformates) to yield products resulting from an Arbuzov-type rearrangement (297).



Similar results were obtained with phenyl dimethyl phosphonite.





Metal phosphides yielded the expected products on reaction with chloroformates (347).




#### 12. $\alpha$ -Amino Phosphonates

Acylation of the amino group in  $\alpha$ -amino phosphonates by chloroformates occurs readily in the presence of a tertiary amine (458).




### 13. Thiophosphates

The thiophosphates shown in the following equations were acylated in good yield by methyl chloroformate (97).




### 14. Organic Sulfates

Alkyl chloroformates behave as alkylating agents toward alkyl hydrogen sulfates (408). Some examples are




Chloromethyl chloroformate behaved differently, yielding methyl chlorosulfonate in 78% yield. The reaction of chlorosulfonic acid with chloroformates yielded alkyl chlorosulfonates (174, 409).




### 15. Amic Acids


Maleamic acid was simultaneously esterified and dehydrated by chloroformates and triethylamine to yield  $\beta$ -cyanoacrylates in high yields (600).



N-substituted maleamic acids yielded isomaleimides from the same reactants (158, 600, 601).




Similarly, dehydration of N-acylamino acids with ethyl chloroformate and triethylamine gave azlactones in 40–70% yields (528).




### 16. Enols

Phosgene reacts with enols via chloroformates to effect their chlorination (14–19). An example is shown.




The chloroformylation of phenylacetylene probably proceeds through a similar type of intermediate (747).




### 17. Phenylazoformates

Ethyl chloroformate gave a mixture of products on reaction with potassium phenylazoformate (523). A free-radical mechanism was proposed.




### 18. Bromine

Alkyl bromides were obtained from the reaction of bromine with alkyl chloroformates (340, 524).



### 19. Epoxides


As mentioned later in section V, chloroformates react with epoxides to yield  $\beta$ -chlorocarbonates (360).



Acids or bases catalyze this reaction.

### 20. Nitrite Esters

The following reaction has been reported (519). Its



generality is unknown.

### 21. Reducing Agents

Formic acid was obtained in 51% yield by treatment of ethyl chloroformate with sodium amalgam (725). The electrolytic reduction of methyl chloroformate has also been studied (256).


## V. POLYMER SYNTHESSES FROM CHLOROFORMATES

The utility of chloroformates, especially bis(chloroformates), has been greatly extended recently because of the recent research on interfacial polymerization. They have been used for the preparation of many different classes of polymers through their acid chloride type reactivity. Brief summaries of the available literature on these various types are presented below.


### A. POLYCARBONATES

#### 1. Aliphatic Polycarbonates

The condensation of an aliphatic bis(chloroformate) with an aliphatic diol can be effected merely by heating under vacuum (412).




A patent has also described the condensation of diols with trichloromethyl chloroformate to yield aliphatic polycarbonates that were described as being wax-like with good oil solubility (716).



The incorporation of sucrose into polycarbonates by direct reaction with phosgene or ester exchange with diphenyl carbonate has also been reported (673-675). Reaction of the hydroxyl groups of polyvinyl alcohol with monochloroformates was described in a recent patent (253).


#### 2. Aliphatic-Aromatic Polycarbonates

Mixed polycarbonates of this type have been prepared by both possible routes (32, 145, 147, 212, 214, 252, 261, 262, 580, 620). The polycondensation of an aliphatic bis(chloroformate) with a bisphenol has been



effected in interfacial systems as well as by the use of stoichiometric amounts of a tertiary base (*e.g.*, pyridine). The mixed polycarbonate from ethylene glycol and 2,2-bis(4-hydroxyphenyl)propane undergoes an interesting thermal decomposition at 280°. Ethylene carbonate and the polycarbonate of 2,2-bis(4-hydroxyphenyl)propane (665) were obtained in quantitative yield.


The alternate route using an aromatic bis(chloroformate) proceeds well only when stoichiometric amounts



of a tertiary base are employed (619). Attempts at using this route in interfacial systems have yielded the completely aromatic polycarbonate.


#### 3. Aromatic Polycarbonates

Completely aromatic polycarbonates are prepared most economically by the direct phosgenation of bisphenols. They can also be readily prepared by the reaction of a bisphenol with an aryl bis(chloroformate). This latter route can be used advantageously to prepare polycarbonates containing bisphenols that cannot be



phosgenated directly, *i.e.*, the most acidic and hindered ones. This polymerization can be carried out in interfacial systems, by using stoichiometric amounts of a tertiary base, and catalytically in inert solvents using metal-containing catalysts (77, 78, 206-211, 213, 218-220, 222, 579, 614-619). Catalysis of this type of interfacial polymerization with phosphonium, arsonium, and sulfonium salts has been the subject of a recent patent (255).

The self-polymerization of an aryl bis(chloroformate) to an aromatic polycarbonate by hydrolysis in an inter-



facial system has also been accomplished. This process lacks practicality because of poor phosgene efficiency.

Cyclic tetramers of bisphenol-A carbonate have been isolated (21%) from the reaction of bisphenol-A dichlo-

roformate with bisphenol-A under high dilution reaction conditions. Polymerization of these cyclic oligomers to very high molecular weight polymers was also effected (621).


#### 4. Copolycarbonates

The interfacial polymerization of hydroxyl-terminated aromatic polycarbonates with polymeric bis(chloroformates) to yield high molecular weight copolymers has been patented. The polymeric bis(chloroformates) that were described were prepared from polyethylene glycols, polytetramethylene glycol, polystyrene glycol, and hydroxyl-terminated poly(ethylene adipate) (398, 468).

Copolymerization of a bisphenol with a bis(chloroformate) and phenyl phosphonyl dichloride was used for the preparation of poly(carbonate-phosphonates) in interfacial systems (254). Phosphonium salts were used as catalysts.

#### B. POLYESTERS

The reaction of a bis(chloroformate) with a dicar-




boxylic acid in the presence of excess tertiary amine has been patented as a route to polyesters (247).

#### C. POLYURETHANS


##### 1. Aliphatic Polyurethans

Many different aliphatic polyurethans have been prepared in interfacial systems from aliphatic bis(chloroformates) and diamines (67, 115-117, 205, 342, 361, 387, 411, 444, 526, 562, 728, 730). This route, of




course, is an alternate to that using diisocyanates and glycols. Low molecular weight aliphatic polyurethans have been prepared in anhydrous systems using calcium hydroxide and other metalliferous hydrogen chloride acceptors (236, 653). Cross-linked, aliphatic polyurethans have been obtained from the above reactions by using reactants with a functionality of greater than two (30, 166).


Polyurethans have been prepared from bis(chloroformates) and diamines by using an excess of the diamine and heating in an inert solvent (332).




The use of a chloroformate in an intermediate step in a melt process for polyurethane synthesis has been de-



scribed (612). A similar technique, but a variant on an interfacial process, is also shown (27, 344, 402, 605, 638).




Oligomeric urethans were prepared in 30-70% yields by the reaction of excess diamine with bis(chloroformates) in dilute solution (378). Earlier work employed



the conversion of hydroxyl-terminated oligomers from the diisocyanate-glycol reaction to chloroformates, followed by reaction with mono(carbobenzoxy)hexamethylenediamine, for their characterization (746).

##### 2. Aliphatic-Aromatic Polyurethans

Polyurethans prepared from piperazine and aryl bis(chloroformates) have been prepared interfacially (221, 362, 363, 638, 666, 726-728, 730). The polymers ob-



tained possessed softening temperatures of  $\geq 200^\circ$ . Polyurethans of this class have also been prepared by the interfacial polymerization of the bis(chloroformate) of neopentyl glycol with a series of aromatic diamines (376).

The polymerization of bis(chloroformates) with diamines to yield polyurethans directly on the surface of fibers has been described as a method of shrink-proofing fabrics (436).

## D. COPOLYURETHANS

1. *Poly(urethan carbonates)*

Interfacial polymerization of a bis(chloroformate) with a diamine and a bisphenol yielded the title copolymers (146). Various diamines and bisphenols have been used in varying ratios.

Block poly(urethan carbonates) have been prepared from the polymerization of low molecular weight chloroformate-terminated polycarbonates with aliphatic diamines (467). Polymerization of diisocyanates with low molecular weight hydroxyl-terminated polycarbonates has yielded similar products (488). Using amine-dicarboxylic acid salts with low molecular weight, chloroformate-terminated polycarbonates gave poly(urethane amide carbonates) (488).


Polymerization of dichloroformates prepared from ester-diols with diamines yielded poly(urethan esters) (733).

2. *Poly(urethan sulfonamides)*

Poly(urethan sulfonamides) have been reportedly prepared by the interfacial reaction of a bis(chloroformate), a bis(sulfonyl chloride), and a diamine (328, 343).

3. *Poly(urethan amides)*

The subject polymers have been prepared by interfacial polymerization of a diacid chloride, a diamine and a bis(chloroformate) (194, 374, 375, 441, 731, 732). An example using piperazine as the diamine is shown.




Bis(chloroformates) prepared from phosgene and polyalkylene oxide glycols have also been used in this type of polymerization (142, 143, 235, 260, 375, 401).

4. *Poly(urethan ethers)*

Polymerization of a bis(chloroformate) derived from a glycol polyether with diamines yielded elastomeric polyurethans (264).


## E. POLYMERIZATION OF CHLOROFORMATES WITH DIEPOXIDES

The reaction of a bis(chloroformate) with a diepoxyde has led to polymers of the type shown in the following equation (146).



## F. POLYMERIZATION OF CHLOROFORMATES WITH BIS(HYDRAZIDES)

Fiber-forming polymers have been prepared by polymerization of bis(chloroformates) with bis(hydrazides) (125, 126, 146, 175).



## G. USES OF CHLOROFORMATES IN ADDITION-TYPE POLYMERS

Unsaturated chloroformates have found utility because after reaction of the chloroformate group with other compounds, the resulting monomers can be polymerized through the double bonds (84, 291, 491-505, 548). Saturated and/or unsaturated chloroformates were condensed with various unsaturated and/or saturated hydroxy compounds and the resulting carbonates polymerized through their double bonds.

Similarly, reaction of unsaturated chloroformates with nitrogen compounds (*e.g.*, NH<sub>2</sub>OH and NH<sub>2</sub>NH<sub>2</sub>) led to polymerizable derivatives (85).

A recent patent has claimed the preparation of urethans by reaction of compounds of the general structure CH<sub>2</sub>=CH-S-Z-C (R<sub>1</sub>R<sub>2</sub>)NHR (Z = any bivalent radical) with chloroformates (596). These materials were then copolymerized with various acrylic-type compounds.

Allyl chloroformate has been found to be a catalyst in the photopolymerization of vinyl monomers, especially acrylates (598). Peroxy compounds prepared from chloroformates and hydroperoxides (637) and peroxy-carboxylic acids (541) have been described as initiators for vinyl polymerizations.

## H. USES OF CHLOROFORMATES IN THERMOSET POLYMERS

Patents have been issued on the use of alkyl and alkenyl chloroformates for the curing of phenol-formaldehyde resins (419, 549). Ethyl chloroformate is claimed to eliminate shrinkage when it is used to set fluid or semifluid thermosetting resinous mixtures.

The use of bis(chloroformates) or their thio analogs for the tanning of leather has been patented (564). Another patent has described the reaction of chloroformates with proteins to give derivatives which were useful in photographic film manufacture (245).

### I. MISCELLANEOUS USES OF CHLOROFORMATES IN POLYMERIC SYSTEMS

Bis(chloroformates) prepared from polyalkylene oxide glycols have found varied uses in preparing polymers (143, 235, 260). For example, condensation of the bis(chloroformate) of poly(tetramethylene ether) glycol with excesses of various diamines gave poly(ether urethans) with terminal NH<sub>2</sub> groups. Further reaction of these prepolymers with aliphatic bis(chloroformates) yielded useful elastomers.

Conversion of low molecular weight, hydroxyl-terminated polyesters to bis(chloroformates), followed by their reaction with diamines and aromatic bis(chloroformates) have yielded poly(ester urethans) (365). Piperazines and the bis(chloroformates) of resorcinol and hydroquinone were used.

Polymers with utility for tanning, dyeing, and textile water repellents have been prepared from chloroformates of alcohols derived from palm oil fatty acids and polyethylenimines (688).

### VI. REFERENCES

- (1) Abbott Laboratories, Inc., British Patent 255,971; *Chem. Abstr.*, **21**, 2908<sup>3</sup> (1927).
- (2) Abbott Laboratories, Inc., Belgian Patent 596,083 (1961).
- (3) Adam, K., and Trieschmann, H. G., German Patent 890,-792; *Chem. Abstr.*, **50**, 12,099h (1956).
- (4) Adams, P., Kaiser, D. W., and Peters, G. A., *J. Org. Chem.*, **18**, 934 (1953).
- (5) Adams, R., and Segur, J. B., *J. Am. Chem. Soc.*, **45**, 785 (1923).
- (6) Adembri, G., *Ann. Chim. (Rome)*, **46**, 62 (1956); *Chem. Abstr.*, **50**, 13,714e (1956).
- (7) American Cyanamid Co., British Patent 586,633; *Chem. Abstr.*, **41**, 6898a (1947).
- (8) American Cyanamid Co., British Patent 773,011; *Chem. Abstr.*, **51**, 15,605h (1957).
- (9) Andrianov, K. A., and Dabagova, A. K., *Zh. Obshch. Khim.*, **30**, 1968 (1960); *Chem. Abstr.*, **55**, 6363d (1961).
- (10) Anschutz, R., and Emery, W. O., *Ann.*, **273**, 56 (1893).
- (11) Antia, M. B., and Kaushal, R., *Agra Univ. J. Res.*, **3**, 197 (1954); *Chem. Abstr.*, **49**, 6121h (1955).
- (12) Arbuzov, A. E., and Kamai, G., *J. Gen. Chem. USSR*, **17**, 2149 (1947); *Chem. Abstr.*, **42**, 4523g (1948).
- (13) Arens, J. F., Volger, H. C., Dornbos, T., Bonnema, J., Greidanus, J. W., and Van den Hende, J. H., *Rec. trav. chim.*, **75**, 1459 (1956).
- (14) Arnold, Z., and Sórm, F., *Chem. Listy*, **51**, 1082 (1957); *Chem. Abstr.*, **51**, 13,761c (1957).
- (15) Arnold, Z., and Zemlicka, J., *Chem. Listy*, **52**, 458 (1958); *Chem. Abstr.*, **53**, 4164a (1959).
- (16) Arnold, Z., and Zemlicka, J., *Proc. Chem. Soc.*, 227 (1958).
- (17) Arnold, Z., and Zemlicka, J., *Collection Czech. Chem. Commun.*, **24**, 786 (1959); *Chem. Abstr.*, **53**, 11,206b (1959).
- (18) Arnold, Z., and Zemlicka, J., *Collection Czech. Chem. Commun.*, **24**, 2378 (1959); *Chem. Abstr.*, **54**, 1274b (1960).
- (19) Arnold, Z., and Zemlicka, J., *Collection Czech. Chem. Commun.*, **24**, 2385 (1959); *Chem. Abstr.*, **54**, 1274h (1960).
- (20) Ashburn, H. G., Collett, A. R., and Larzzel, C. L., *J. Am. Chem. Soc.*, **60**, 2933 (1938).
- (21) Augustin, E. A., *Dissertation Abstr.*, **22**, 733 (1961); *Chem. Abstr.*, **56**, 371e (1962).
- (22) Aumuller, W., Korger, G., Weyer, R., and Wagner, H. (to Farbwereke Hoechst A.-G.), German Patent 1,048,907; *Chem. Abstr.*, **55**, 4433i (1961).
- (23) Aumuller, W., Korger, G., and Wagner, H. (to Farbwereke Hoechst A.-G.), German Patent 1,075,587; *Chem. Abstr.*, **55**, 13,339e (1961).
- (24) Auwers, K. v., and Schaich, W., *Ber.*, **54**, 1738 (1921).
- (25) Auwers, K. v., and Schaich, W., *Ber.*, **54**, 1768 (1921).
- (26) Auwers, K. v., *Ber.*, **60**, 2122 (1927).
- (27) Badische Anilin- und Soda-Fabrik A.-G., French Patent 961,754; *Chem. Abstr.*, **46**, 6148i (1952).
- (28) Badische Anilin- und Soda-Fabrik A.-G., German Patent 821,207; *Chem. Abstr.*, **47**, 3329e (1953).
- (29) Badische Anilin- und Soda-Fabrik A.-G., British Patent 730,599; *Chem. Abstr.*, **50**, 12,100a (1956).
- (30) Badische Anilin- und Soda-Fabrik A.-G., British Patent 743,917; *Chem. Abstr.*, **50**, 12,548d (1956).
- (31) Badische Anilin- und Soda-Fabrik A.-G., German Patent 1,034,912; *Chem. Abstr.*, **54**, 20,060f (1960).
- (32) Badische Anilin- und Soda-Fabrik A.-G., British Patent 843,314; *Chem. Abstr.*, **55**, 10,962h (1961).
- (33) Badische Anilin- und Soda-Fabrik A.-G., German Patent 1,091,555; *Chem. Abstr.*, **55**, 24,119e (1961).
- (34) Baker, B. R., McEvoy, F. J., Schaub, R. E., Joseph, J. P., and Williams, J. H., *J. Org. Chem.*, **18**, 161 (1953).
- (35) Baltze, K. H., and Heidenbluth, K., *Chem. Ber.*, **92**, 986 (1959).
- (36) Barber, H. J., Cottrell, H. J., and Green, M. B., *J. Appl. Chem.*, **3**, 259 (1953).
- (37) Barral, E., and Morel, A., *Compt. rend.*, **128**, 1578 (1899).
- (38) Barral, E., and Morel, A., *Bull. soc. chim. France*, [3] **21**, 722 (1899).
- (39) Barnden, R. L., Evans, R. M., Hamlet, J. C., Heins, B. A., Jansen, A. B. A., Trevett, M. E., and Webb, G. B., *J. Chem. Soc.*, 3733 (1953).
- (40) Bässler, P., *J. prakt. Chem.*, [2] **16**, 125 (1877).
- (41) Batke, T. L., Dorfman, L. M., and LeRoy, S. J., *J. Chem. Phys.*, **17**, 566 (1949).
- (42) Bayer, F., and Co., German Patent 114,025; *Friedländer*, **6**, 1161 (1900).
- (43) Bayer, F., and Co., German Patent 117,624; *Chem. Zentr.*, **1**, 428 (1901); *Friedländer*, **6**, 1164 (1900).
- (44) Bayer, F., and Co., German Patent 117,625; *Friedländer*, **6**, 1162 (1900).
- (45) Bayer, F., and Co., German Patent 118,536; *Friedländer*, **6**, 1165 (1901).
- (46) Bayer, F., and Co., German Patent 118,537; *Friedländer*, **6**, 1166 (1901).
- (47) Bayer, F., and Co., German Patent 118,566; *Friedländer*, **6**, 1163 (1901).
- (48) Bayer, F., and Co., German Patent 120,863; *Friedländer*, **6**, 1169 (1901).
- (49) Bayer, F., and Co., German Patent 120,865; *Friedländer*, **6**, 1170 (1901).
- (50) Bayer, F., and Co., German Patent 121,223; *Friedländer*, **6**, 1173 (1901).
- (51) Bayer, F., and Co., German Patent 251,805; *Friedländer*, **11**, 82 (1912).
- (52) Bayer, F., and Co., German Patent 255,942; *Friedländer*, **11**, 115 (1912).
- (53) Bayer, F., and Co., German Patent 297,933; *Friedländer*, **13**, 126 (1921).
- (54) Benedict, G. E., and Russel, R. R., *J. Am. Chem. Soc.*, **73**, 5444 (1951).
- (55) Ben Ishai, D., *J. Am. Chem. Soc.*, **78**, 4962 (1956).
- (56) Berger, F. M., and Ludwig, B. J., U. S. Patent 2,656,378; *Chem. Abstr.*, **49**, 1104h (1955).

- (57) Berger, F. M., and Ludwig, B. J. (to Carter Products, Inc.), U. S. Patent 2,724,720; *Chem. Abstr.*, **50**, 12,104b (1956).
- (58) Bergmann, M., and Zervas, L., *Ber.*, **B65**, 1192 (1932).
- (59) Besfamilnil, I. B., Kotrelev, V. N., and Kostryukova, T. D., USSR Patent 138,030; *Chem. Abstr.*, **56**, 4956a (1962).
- (60) Bestian, H., *Ann.*, **566**, 210 (1950).
- (61) Bestmann, H. J., and Schulz, H., *Angew. Chem.*, **73**, 27 (1961).
- (62) Biedermann, J., *Ber.*, **22**, 2762 (1889).
- (63) Birkofe, L., and Barnikel, C. D., *Chem. Ber.*, **92**, 2990 (1959).
- (64) Birkofe, L., Richter, P., and Ritter, A., *Chem. Ber.*, **93**, 2804 (1960).
- (65) Bissinger, W. E., Fredenburg, R. H., Kadesch, R. D., Kung, F., Langstrom, J. H., Stevens, H. C., and Strain, F., *J. Am. Chem. Soc.*, **69**, 2955 (1947).
- (66) Bissinger, W. E. (to Pittsburgh Plate Glass Co.), U. S. Patent 2,517,964; *Chem. Abstr.*, **45**, 2019b (1951).
- (67) Black, W. B. (to Chemstrand Corp.), U. S. Patent 2,987,-494; *Chem. Abstr.*, **55**, 25,361c (1961).
- (68) Boardman, F., "The Sni Reaction: Derivatives of Chloroformic Acid with Silver Nitrate," Seminars, Polytechnic Institute of Brooklyn, Course 1973-4-5-6, fall, 1959, pp. 1-4.
- (69) Boehme, T., and Mehta, D., *Ber.*, **71**, 1797 (1938).
- (70) Bohl, L. E. (to Columbia-Southern Chemical Corp.), U. S. Patent 2,651,658; *Chem. Abstr.*, **48**, 13,716c (1954).
- (71) Böhme, H., *Ber.*, **B74**, 248 (1941).
- (72) Böhme, H., and Shurhoff, W., *Chem. Ber.*, **84**, 28 (1951).
- (73) Boissonnas, R. A., *Helv. Chim. Acta*, **34**, 874 (1951).
- (74) Booth, R. B., and Light, C. E. (to American Cyanamid Co.), British Patent 650,493; *Chem. Abstr.*, **46**, 2573e (1952).
- (75) Booth, R. B., and Hill, J. V. (to American Cyanamid Co.), British Patent 650,498; *Chem. Abstr.*, **45**, 7508g (1951).
- (76) Boschan, R., *J. Am. Chem. Soc.*, **81**, 3341 (1959).
- (77) Bottenbruch, L., and Schnell, H. (to Farbenfabriken Bayer A.-G.), German Patent Application F171671Vb/39c (March 26, 1955).
- (78) Bottenbruch, L., and Schnell, H. (to Farbenfabriken Bayer A.-G.), German Patent 959,497; *Chem. Abstr.*, **53**, 13,665h (1959).
- (79) Bottenbruch, L., and Schnell, H. (to Farbenfabriken Bayer A.-G.), German Patent 1,101,386; *Chem. Abstr.*, **56**, 412d (1962).
- (80) Bowden, S. T., *J. Chem. Soc.*, 310 (1939).
- (81) Brady, O. L., and McHugh, G. P., *J. Chem. Soc.*, **123**, 1190 (1923).
- (82) Brady, O. L., and McHugh, G. P., *J. Chem. Soc.*, **127**, 2414 (1925).
- (83) Brady, O. L., and Peakin, F., *J. Chem. Soc.*, 2267 (1929).
- (84) Bralley, J. A., and Poppe, F. B. (to B. F. Goodrich Co.), British Patent 611,529; *Chem. Abstr.*, **43**, 4298h (1949).
- (85) Bralley, J. A. (to B. F. Goodrich Co.), U. S. Patent 2,568,-608; *Chem. Abstr.*, **46**, 5084c (1952).
- (86) Brandy, J. H., and LeRoy, D. J., *J. Chem. Phys.*, **21**, 1049 (1953).
- (87) Bredereck, H., Gompper, R., Rempfer, H., Klemm, K., and Keck, H., *Chem. Ber.*, **92**, 329 (1959).
- (88) Brintzinger, H., and Janecke, J., *Chem. Ber.*, **83**, 103 (1950).
- (89) Brotherton, T. K., and Lynn, J. W. (to Union Carbide Corp.), U. S. Patent 3,008,978; *Chem. Abstr.*, **57**, 5807e (1962).
- (90) Bruchhausen, F. v., and Knabe, J., *Arch. Pharm.*, **287**, 601 (1954).
- (91) Butlerow, A., *Z. Chem.*, 448 (1863).
- (92) Butlerow, A., *Z. Chem.*, 484 (1863).
- (93) Butlerow, A., *Bull. soc. chim. France*, 582 (1863).
- (94) Butlerow, A., *Jahresber. Fortschr. Chem.*, 474 (1863).
- (95) Buu-Hoi, Ng. Ph., and Janicaud, J., *Bull. soc. chim. France*, **12**, 640 (1945).
- (96) Cacace, F., Fabrizi, G., and Ziffero, M., *Ann. Chim. (Rome)*, **46**, 91 (1956); *Chem. Abstr.*, **50**, 12,071d (1956).
- (97) Cadogan, J. I. G., *J. Chem. Soc.*, 3067 (1961).
- (98) Cahours, A., *Ann.*, **64**, 312 (1847).
- (99) Caldwell, J. R. (to Eastman Kodak Co.), U. S. Patent 2,801,231; *Chem. Abstr.*, **51**, 18,704b (1957).
- (100) Caldwell, J. R., and Jackson, W. J. (to Eastman Kodak Co.), U. S. Patent 3,096,359; *Chem. Abstr.*, **59**, 13,841f (1963).
- (101) Callsen, J. (to I. G. Farbenindustrie A.-G.), German Patent 491,492; *Friedländer*, **16**, 2469; *Chem. Abstr.*, **24**, 2148<sup>e</sup> (1930).
- (102) Calver, H. O., *J. Am. Chem. Soc.*, **48**, 1009 (1926).
- (103) Campbell, J. A., *J. Org. Chem.*, **22**, 1259 (1957).
- (104) Capelli, G., *Gazz. chim. ital.*, **50**, II, 8 (1920); *Chem. Abstr.*, **15**, 524 (1921).
- (105) Carothers, W. H., and Berchet, G. J. (to E. I. du Pont de Nemours and Co., Inc.), U. S. Patent 2,073,363; *Chem. Abstr.*, **31**, 3503<sup>a</sup> (1937).
- (106) Carothers, W. H., and Berchet, G. J. (to E. I. du Pont de Nemours and Co., Inc.), U. S. Patent 2,136,178; *Chem. Abstr.*, **33**, 1345<sup>a</sup> (1939).
- (107) Carpenter, F. H., and Gish, D. T., *J. Am. Chem. Soc.*, **74**, 3818 (1952).
- (108) Carré, P., and Passedouet, H., *Compt. rend.*, **200**, 1767 (1935).
- (109) Carré, P., and Passedouet, H., *Compt. rend.*, **201**, 898 (1935).
- (110) Carré, P., *Bull. soc. chim. France*, [5] **3**, 1064 (1936).
- (111) Carré, P., and Peigné, L., *Compt. rend.*, **202**, 2159 (1936).
- (112) Carter, H. E., Frank, R. L., and Johnston, H. W., *Org. Syn.*, **23**, 13 (1943).
- (113) Carter, H. E., Frank, R. L., and Johnston, H. W., "Organic Syntheses," Coll. Vol. III, John Wiley and Sons, Inc., New York, N. Y., 1955, p. 167.
- (114) Cassady, D. R., Ainsworth, C., Easton, N. R., Livezey, M., Sigal, M. V., Jr., and Van Heyningen, E., *J. Org. Chem.*, **23**, 923 (1958).
- (115) Celanese Corporation of America, Inc., British Patent 683,-504; *Chem. Abstr.*, **47**, 5169a (1953).
- (116) Celanese Corporation of America, Inc., British Patent 685,-729; *Chem. Abstr.*, **47**, 7257g (1953).
- (117) Celanese Corporation of America, Inc., British Patent 743,-918; *Chem. Abstr.*, **50**, 12,548g (1956).
- (118) Chaney, A., and Wolfrom, M. L., *J. Org. Chem.*, **26**, 2998 (1961).
- (119) Channing, D. M., Turner, P. B., and Young, G. F., *Nature*, **167**, 487 (1951); *Chem. Abstr.*, **46**, 902e (1951).
- (120) Chattaway, F. D., and Saerens, E., *J. Chem. Soc.*, **117**, 708 (1920).
- (121) Chemische Fabrik von Heyden A.-G., German Patent 99,057; *Friedländer*, **5**, 727 (1898).
- (122) Chemische Fabrik von Heyden A.-G., German Patent 109,-933; *Friedländer*, **5**, 730 (1900).
- (123) Chemische Fabrik von Heyden A.-G., German Patent 116,-386; *Friedländer*, **6**, 1160 (1900).
- (124) Chemische Fabrik von Heyden A.-G., German Patent 117,-346; *Friedländer*, **5**, 954 (1900).
- (125) Chemstrand Corporation, Belgian Patent 582,348 (1959).

- (126) Chemstrand Corporation, British Patent 880,755; *Chem. Abstr.*, **54**, 23,426f (1960).
- (127) Chenicek, J. A. (to Universal Oil Products Co.), U. S. Patent 2,649,473; *Chem. Abstr.*, **47**, 11,764g (1953).
- (128) Cherbuliez, E., Baehler, B., Sussmann, A. R., and Rabinowitz, J., *Helv. Chim. Acta*, **44**, 1806 (1961).
- (129) Chichibabin, A. E., *Compt. rend.*, **213**, 355 (1941).
- (130) Childs, A. F., Plant, S. G. P., Tompsett, A. L. L., and Weeks, G. A., *J. Chem. Soc.*, 2180 (1948).
- (131) Choppin, A. R., Frediani, H., and Kirby, G. F., Jr., *J. Am. Chem. Soc.*, **61**, 3176 (1939).
- (132) Choppin, A. R., and Kirby, G. F., Jr., *J. Am. Chem. Soc.*, **62**, 1592 (1940).
- (133) Choppin, A. R., and Roger, J. W., *J. Am. Chem. Soc.*, **70**, 2967 (1948).
- (134) Choppin, A. R., and Compere, E. L., *J. Am. Chem. Soc.*, **70**, 3797 (1948).
- (135) Claisen, L., and Zedel, W., *Ber.*, **21**, 3397 (1888).
- (136) Claisen, L., *Ber.*, **21**, 3567 (1888).
- (137) Claisen, L., *Ber.*, **25**, 1760 (1892).
- (138) Claisen, L., *Ber.*, **27**, 3182 (1894).
- (139) Claisen, L., and Haase, E., *Ber.*, **33**, 3778 (1900).
- (140) Cloez, S., *Ann.*, **60**, 259 (1846).
- (141) Cloez, S., *Ann. Chim.*, [3] **17**, 297 (1846).
- (142) Cluff, E. F., Paper presented at the Gordon Research Conference on Polymers, Colby College, New London, N. H., July, 1960.
- (143) Cluff, E. F., Gladding, E. K., and Rogan, J. B., *J. Appl. Polymer Sci.*, **5**(13), 80 (1961).
- (144) Coles, R. F., and Levine, H. A. (to General Aniline and Film Corp.), U. S. Patent 2,942,025; *Chem. Abstr.*, **54**, 24,464a (1960).
- (145) Columbia-Southern Chemical Corp., Australian Patent 51,087 (1959).
- (146) Columbia-Southern Chemical Corp., Belgian Patent 573,065 (1959).
- (147) Columbia-Southern Chemical Corp., Belgian Patent 581,249 (1959).
- (148) Columbia-Southern Chemical Corp., Belgian Patent 584,933 (1959).
- (149) Columbia-Southern Chemical Corp., British Patent 878,115; *Chem. Abstr.*, **56**, 7216g (1962).
- (150) Compere, E. L., *Proc. Louisiana Acad. Sci.*, **6**, 93 (1942); *Chem. Abstr.*, **36**, 2779<sup>b</sup> (1942).
- (151) Conant, J. B., and Kirner, W. R., *J. Am. Chem. Soc.*, **46**, 232 (1924).
- (152) Conant, J. B., Kirner, W. R., and Hussey, R. E., *J. Am. Chem. Soc.*, **47**, 488 (1925).
- (153) Conrad, M., and Guthzeit, M., *Ann.*, **214**, 31 (1882).
- (154) Contardi, A., and Ercoli, A., *Gazz. chim. ital.*, **64**, 522 (1934); *Chem. Abstr.*, **29**, 1392<sup>a</sup> (1935).
- (155) Contardi, A., and Ercoli, A., *Congr. intern. quím. Pura apt.*, **9**<sup>o</sup>, Madrid, **5**, 163 (1934); *Chem. Abstr.*, **31**, 1764<sup>b</sup> (1937).
- (156) Copisarow, M., *J. Chem. Soc.*, 251 (1929).
- (157) Coppock, W. H., *J. Org. Chem.*, **22**, 325 (1957).
- (158) Cotter, R. J., Sauers, C. K., and Whelan, J. M., *J. Org. Chem.*, **26**, 10 (1961).
- (159) Councler, C., *Ber.*, **13**, 1697 (1880).
- (160) Criscuoli, A., *Chim. ind. agr. biol.*, **14**, 271 (1938); *Chem. Abstr.*, **34**, 7859<sup>a</sup> (1940).
- (161) Crunden, E. W., and Hudson, R. F., *J. Chem. Soc.*, 3748 (1961).
- (162) Curtius, Th., and Heidenreich, K., *Ber.*, **27**, 773 (1894).
- (163) Curtius, Th., and Heidenreich, K., *J. prakt. Chem.*, [2] **52**, 454 (1895).
- (164) Dadien, A., and Kohlrausch, K. W. F., *Monatsh.*, **57**, 225 (1931).
- (165) Dains, F. B., and Wertheim, E., *J. Am. Chem. Soc.*, **42**, 2307 (1920).
- (166) Danmiller, G. (to Badische Anilin- und Soda-Fabrik A.-G.), U. S. Patent 2,748,102; *Chem. Abstr.*, **50**, 16,184i (1956).
- (167) Davies, N. H., *J. Chem. Soc.*, 1357 (1951).
- (168) Davis, O. C. M., *J. Chem. Soc.*, **97**, 949 (1910).
- (169) Degering, E. F., Jenkins, G. L., and Sanders, B. E., *J. Am. Pharm. Assoc.*, **39**, 624 (1950); *Chem. Abstr.*, **45**, 1950a (1951).
- (170) De Groote, M., and Keiser, B. (to Petrolite Corp., Ltd.), U. S. Patent 2,472,573; *Chem. Abstr.*, **43**, 6818e (1949).
- (171) De Groote, M., and Keiser, B. (to Petrolite Corp., Ltd.), U. S. Patent 2,574,537; *Chem. Abstr.*, **46**, 2279b (1952).
- (172) Delaby, R., Damiens, R., and Capman, M. L., *Compt. rend.*, **246**, 3353 (1958).
- (173) Delepine, M., *Bull. soc. chim. France*, [4] **27**, 39 (1920).
- (174) Descude, P., *Compt. rend.*, **132**, 1567 (1901).
- (175) De Witt, H. D. (to Chemstrand Corp.), U. S. Patent 2,941,-984; *Chem. Abstr.*, **54**, 23,426f (1960).
- (176) Diels, O., *Ber.*, **36**, 736 (1903).
- (177) Diels, O., and Nawiaski, P., *Ber.*, **37**, 3672 (1904).
- (178) Diels, O., and Jacoby, E., *Ber.*, **41**, 2392 (1908).
- (179) Diels, O., and Gollman, R., *Ber.*, **44**, 3158 (1911).
- (180) Dirscherl, W., *Z. physiol. Chem.*, **239**, 49 (1936); *Chem. Abstr.*, **30**, 2982<sup>a</sup> (1936).
- (181) Dixon, A. E., *J. Chem. Soc.*, **89**, 909 (1906).
- (182) Dobrosserdov, D., *Zapiski Kazansk Univ.*, **326** (1909); *Chem. Zentr.*, I, 955 (1911).
- (183) Douglas, J. B., and Johnston, T. B., *J. Am. Chem. Soc.*, **60**, 1486 (1938).
- (184) Dow Chemical Co., British Patent 803,203; *Chem. Abstr.*, **53**, 14,972i (1959).
- (185) Dox, A. W., and Yoder, L., *J. Am. Chem. Soc.*, **45**, 723 (1923).
- (186) Dumas, J., *Ann. Chim.*, [2] **54**, 225 (1833).
- (187) Dumas, J., *Ann. Chim. Phys.*, [2] **56**, 113 (1834).
- (188) Dumas, J., *Ann.*, **10**, 277 (1834).
- (189) Dumas, J., and Peligot, E., *Ann.*, **15**, 1 (1935).
- (190) Dupont, G., *Compt. rend.*, **148**, 1522 (1909).
- (191) Dupont, L. A., French Patent 771,653; *Chem. Abstr.*, **29**, 816<sup>a</sup> (1935).
- (192) Duynstee, E. F. J., and Veerkamp, Th., *Rec. trav. chim.*, **81**, 241 (1962).
- (193) Dyson, G. M., *Chem. Rev.*, **4**, 149–152 (1927).
- (194) E. I. du Pont de Nemours and Co., Inc., British Patent 846,024 (1960).
- (195) Einhorn, A., *Ber.*, **42**, 2237 (1909).
- (196) Einhorn, A., *Ber.*, **42**, 2772 (1909).
- (197) Einhorn, A., German Patent 224,108; *Friedländer*, **10**, 1087 (1910).
- (198) Einhorn, A., and Rothlauf, L., *Ann.*, **382**, 237 (1911).
- (199) Engel, K., and Morel, C. J. (to J. R. Geigy and Co.), U. S. Patent 2,573,420; *Chem. Abstr.*, **46**, 3567g (1952).
- (200) Ettlinger, M. G., and Hodgkins, J. E., *J. Am. Chem. Soc.*, **77**, 1833 (1955).
- (201) Ettlinger, M. G., and Hodgkins, J. E., *J. Org. Chem.*, **21**, 204 (1956).
- (202) Ettlinger, M. G., and Lundeen, A. J., *J. Am. Chem. Soc.*, **78**, 1952 (1956).
- (203) Falck, E., *Ber.*, **18**, 2467 (1885).
- (204) Falck, E., *Ber.*, **19**, 1481 (1886).
- (205) Farbenfabriken Bayer A.-G., British Patent 706,717; *Chem. Abstr.*, **48**, 13,276e (1954).
- (206) Farbenfabriken Bayer A.-G., British Patent 772,627; *Chem. Abstr.*, **51**, 10,953i (1957).

- (207) Farbenfabriken Bayer A.-G., British Patent 800,815 (1958).
- (208) Farbenfabriken Bayer A.-G., Belgian Patent 577,943 (1959).
- (209) Farbenfabriken Bayer A.-G., British Patent 808,485; *Chem. Abstr.*, **53**, 15,641b (1959).
- (210) Farbenfabriken Bayer A.-G., British Patent 808,486; *Chem. Abstr.*, **53**, 15,641d (1959).
- (211) Farbenfabriken Bayer A.-G., British Patent 811,628; *Chem. Abstr.*, **53**, 20,908h (1959).
- (212) Farbenfabriken Bayer A.-G., Belgian Patent 589,144 (1960).
- (213) Farbenfabriken Bayer A.-G., British Patent 824,423; *Chem. Abstr.*, **54**, 10,965c (1960).
- (214) Farbenfabriken Bayer A.-G., South African Patent 60/1563.
- (215) Farbenfabriken Bayer A.-G., German Patent 1,117,598; *Chem. Abstr.*, **57**, 11,106d (1962).
- (216) Farbenfabriken Bayer A.-G., German Patent 1,133,727 (1962).
- (217) Farbwerte Hoechst A.-G., German Patent 232,879; *Friedländer*, **10**, 1241 (1911).
- (218) Farbwerte Hoechst A.-G., German Patent 860,205 (1952).
- (219) Farbwerte Hoechst A.-G., German Patent 875,648 (1953).
- (220) Farbwerte Hoechst A.-G., German Patent 895,899 (1953).
- (221) Farbwerte Hoechst A.-G., German Patent 915,868 (1954).
- (222) Farbwerte Hoechst A.-G., German Patent Application 10,-2001Vb/39c.
- (223) Farthing, A. G., *J. Chem. Soc.*, 3213 (1950).
- (224) Fatianow, Z. *Chem.*, **77** (1864).
- (225) Fatianow, *Jahresber. Fortschr. Chem.* **477** (1864).
- (226) Field, L., Lawson, J. E., and McFarland, J. W., *J. Am. Chem. Soc.*, **78**, 4389 (1956).
- (227) Fierz-David, H. E., and Muller, W., *J. Chem. Soc.*, **125**, 26 (1924).
- (228) Fischer, A. H. (to Minerec Corp.), U. S. Patent 2,608,572; *Chem. Abstr.*, **47**, 5433g (1953).
- (229) Fischer, E., *Ber.*, **41**, 2860 (1908).
- (230) Fischer, E., *Ber.*, **41**, 2875 (1908).
- (231) Flickinger, E., and Treischmann, H. G. (to I. G. Farbenindustrie "in Auflösung"), German Patent 821,207; *Chem. Abstr.*, **47**, 3329e (1953).
- (232) Flynn, E. H., Sigal, M. V., Jr., Wiley, P. F., and Gerzon, K., *J. Am. Chem. Soc.*, **76**, 3121 (1954).
- (233) Flynn, E. H., Murphy, A. W., and McMahon, R. E., *J. Am. Chem. Soc.*, **77**, 3104 (1955).
- (234) Forster, M. O., and Newman, S. H., *J. Chem. Soc.*, **97**, 2570 (1910).
- (235) Frazer, A. H., and Goldberg, E. J. (to E. I. du Pont de Nemours and Co., Inc.), U. S. Patent 2,888,440; *Chem. Abstr.*, **53**, 17,582f (1959).
- (236) Frazer, A. H. (to E. I. du Pont de Nemours and Co., Inc.), U. S. Patent 3,022,344 (1962).
- (237) Freudenberg, K., Eich, H., Knoevenagel, C., and Westphahl, W., *Ber.*, **B73**, 441 (1940).
- (238) Freudenberg, K., and Jacob, W., *Ber.*, **B74**, 1001 (1941).
- (239) Friedel, C., and Crafts, J. M., *Compt. rend.*, **84**, 1450 (1877).
- (240) Friedel, C., and Crafts, J. M., *Ann. Chim.*, [6] **1**, 449 (1884).
- (241) Friedel, C., and Crafts, J. M., *Ann. Chim.*, [6] **1**, 527 (1884).
- (242) Gadamer, J., and Knoch, F., *Arch. Pharm.*, **259**, 135 (1921); *Chem. Abstr.*, **15**, 3273 (1921).
- (243) Gaertner, R., *J. Am. Chem. Soc.*, **74**, 2185 (1952).
- (244) Garmaise, D. L., Schwartz, R., and McKay, A. F., *J. Am. Chem. Soc.*, **80**, 3332 (1958).
- (245) Gates, J. W., and Elius, H. S. (to Eastman Kodak Co.), U. S. Patent 2,713,575; *Chem. Abstr.*, **49**, 13,808c (1955).
- (246) Geiszler, A. O., and Spielman, M. A. (to Abbott Laboratories), U. S. Patent 2,806,053; *Chem. Abstr.*, **52**, 6399f (1958).
- (247) General Electric Co., British Patent 870,096; *Chem. Abstr.*, **56**, 14,173h (1962).
- (248) Gerrard, W., *J. Chem. Soc.*, **218** (1940).
- (249) Gerrard, W., and Schild, F., *Chem. Ind. (London)*, 1232 (1954).
- (250) Geuther, A., *J. prakt. Chem.*, [2] **6**, 160 (1873).
- (251) Geuther, A., *Ann.*, **205**, 223 (1880).
- (252) Gevaert Photo-Producten, N. V., Belgian Patent 552,176; *Chem. Abstr.*, **54**, 25,979c (1960).
- (253) Gevaert Photo-Producten, N.V., Belgian Patent 610,590 (1962).
- (254) Gevaert Photo-Producten, N.V., Belgian Patent 610,954 (1962).
- (255) Gevaert Photo-Producten, N.V., French Patent 1,283,-226 (1962).
- (256) Giacomello, G., and Leutini, M., *Gazz. chim. ital.*, **66**, 350 (1936).
- (257) Gialdi, F., and Barruffini, A., *Farmaco (Pavia)*, Ed. Sci., **12**, 318 (1957); *Chem. Abstr.*, **54**, 10,847h (1960).
- (258) Gleim, C. E., *J. Am. Chem. Soc.*, **76**, 107 (1954).
- (259) Glantz, M. L., University Microfilms, Inc., Ann Arbor, Mich., L.C. Card No. Mic. 60-5359; *Dissertation Abstr.*, **21**, 2476 (1961); *Chem. Abstr.*, **55**, 16,556h (1961).
- (260) Goldberg, E. J. (to E. I. du Pont de Nemours and Co.), U. S. Patent 2,927,098; *Chem. Abstr.*, **54**, 14,747f (1960).
- (261) Golz, H., and Mayer, F. (to Badische Anilin- und Soda-Fabrik A.-G.), British Patent 843,314; *Chem. Abstr.*, **55**, 10,962g (1961).
- (262) Golz, H., and Mayer, F. (to Badische Anilin- und Soda-Fabrik A.-G.), German Patent 1,044,406; *Chem. Abstr.*, **55**, 21,95e (1961).
- (263) Goodyear Tire and Rubber Co., British Patent 851,964; *Chem. Abstr.*, **55**, 11,919i (1961).
- (264) Goodyear Tire and Rubber Co., French Patent 1,278,258 (1962).
- (265) Green, M., and Hudson, R. F., *Proc. Chem. Soc.*, 149 (1959).
- (266) Green, M., *Chem. Ind. (London)*, 435 (1961).
- (267) Grignard, V., Rivat, G., and Urbain, E., *Compt. rend.*, **169**, 1074 (1919).
- (268) Grignard, V., Rivat, G., and Urbain, E., *Compt. rend.*, **169**, 1143 (1919).
- (269) Grignard, V., Rivat, G., and Urbain, E., *Ann. Chim.*, [9] **13**, 229 (1920).
- (270) Grosby, D. G., and Niemann, C., *J. Am. Chem. Soc.*, **76**, 4458 (1954).
- (271) Gross, F., *Ber.*, **18**, 2477 (1885).
- (272) Gross, H., Rieche, A., and Höft, E., *Chem. Ber.*, **94**, 544 (1961).
- (273) Gros, R., *Ann. Pharm. Franç.*, **5**, 459 (1947); *Chem. Abstr.*, **42**, 3702b (1948).
- (274) Gutmann, P., Doctoral Thesis, University of Heidelberg, Germany, 1903.
- (275) Haga, T. (to Mitsubishi Chemical Industries, Ltd.), Japanese Patent 4264 (1959); *Chem. Abstr.*, **54**, 299f (1960).
- (276) Haga, T., Sato, M., and Yoshida, M. (to Mitsubishi Chemical Industries, Ltd.), Japanese Patent 6323 (1958); *Chem. Abstr.*, **54**, 2174b (1960).
- (277) Hales, J. L., Jones, J. I., and Kynaston, W., *J. Chem. Soc.*, 618 (1957).
- (278) Hall, H. K., Jr., *J. Am. Chem. Soc.*, **77**, 5993 (1955).

- (279) Hall, H. K., Jr., *J. Org. Chem.*, **21**, 248 (1956).
- (280) Hall, H. K., Jr., and Morgan, P. W., *J. Org. Chem.*, **21**, 249 (1956).
- (281) Hall, H. K., Jr., *J. Am. Chem. Soc.*, **78**, 2717 (1956).
- (282) Hall, H. K., Jr., *J. Am. Chem. Soc.*, **79**, 5439 (1957).
- (283) Haller, A., *Compt. rend.*, **105**, 169 (1887).
- (284) Haller, A., *Ann. Chim.*, [6] **16**, 403 (1889).
- (285) Haller, A., and Bauer, E., *Compt. rend.*, **152**, 557 (1911).
- (286) Haller, A., and Bauer, E., *Ann. Chim.*, [10] **1**, 275 (1924).
- (287) Hamilton, C. S., and Sly, C., *J. Am. Chem. Soc.*, **47**, 435 (1925).
- (288) Hamilton, C. S., and Eztelmiller, R. E., *J. Am. Chem. Soc.*, **50**, 3360 (1928).
- (289) Hammond, J. A. S., U. S. Patent 1,618,824; *Chem. Abstr.*, **21**, 1128 (1927).
- (290) Hammond, J. A. S., U. S. Patent 2,496,091; *Chem. Abstr.*, **44**, 4033d (1950).
- (291) Hammond, R. (to Imperial Chemical Industries, Ltd.), British Patent 629,019; *Chem. Abstr.*, **44**, 3739i (1950).
- (292) Hantsch, A., *Ber.*, **27**, 1254 (1894).
- (293) Hantsch, A., and Kröber, W., *J. prakt. Chem.*, **223**, 126 (1927).
- (294) Haring, M., *Helv. Chim. Acta*, **42**, 1916 (1959).
- (295) Haring, M., *Helv. Chim. Acta*, **43**, 104 (1960).
- (296) Harford, M. B., Kenyon, J., and Phillips, H., *J. Chem. Soc.*, 179 (1933).
- (297) Harman, D. (to Shell Development Co.), U. S. Patent 2,629,731; *Chem. Abstr.*, **48**, 1418e (1954).
- (298) Haworth, R. D., Lamberton, A. H., and Woodecock, D., *J. Chem. Soc.*, 176 (1947).
- (299) Hechelkammer, W. (to Farbenfabriken Bayer A.-G.), German Patent 940,588; *Chem. Abstr.*, **51**, 12,967i (1957).
- (300) Hechenbleikner, R. (to American Cyanamid Co.), U. S. Patent 2,307,679; *Chem. Abstr.*, **37**, 3454<sup>g</sup> (1943).
- (301) Heiserman, C. E., and Edwards, W. R. (to Celanese Corporation of America), U. S. Patent 2,764,607; *Chem. Abstr.*, **51**, 6106b (1957).
- (302) Henry, L., *Compt. rend.*, **104**, 1618 (1887).
- (303) Hentrich, W., and Engelbrecht, H. J. (to Deutsche Hydrierwerke A.-G.), U. S. Patent 2,269,105; *Chem. Abstr.*, **36**, 2965<sup>3</sup> (1942).
- (304) Hentschel, W., *Ber.*, **17**, 1284 (1884).
- (305) Hentschel, W., *Ber.*, **18**, 1177 (1885).
- (306) Hentschel, W., *J. prakt. Chem.*, [2] **36**, 99 (1887).
- (307) Hentschel, W., *J. prakt. Chem.*, [2] **36**, 209 (1887).
- (308) Hentschel, W., *J. prakt. Chem.*, [2] **36**, 305 (1887).
- (309) Hentschel, W., *J. prakt. Chem.*, [2] **36**, 468 (1887).
- (310) Herzog, J., *Ber.*, **42**, 2557 (1909).
- (311) Heyden, Dr. F. v., and Co., German Patent 58,129; *Friedländer*, **3**, 850 (1891).
- (312) Heyden, Dr. F. v., and Co., German Patent 60,716; *Friedländer*, **3**, 852 (1891).
- (313) Heyden, Dr. F. v., and Co., German Patent 61,848; *Friedländer*, **3**, 853 (1892).
- (314) Heyden, Dr. F. v., and Co., German Patent 72,806; *Friedländer*, **3**, 854 (1893).
- (315) Hillmann-Elies, A., Hillman, G., and Jatzkewitz, H., *Z. Naturforsch.*, **8b**, 445 (1953); *Chem. Abstr.*, **49**, 230e (1955).
- (316) Himmelbauer, R., *J. prakt. Chem.*, [2] **54**, 177 (1896).
- (317) Hochstetter, A., German Patent 282,134; *Friedländer*, **12**, 65 (1915).
- (318) Hodel, E., and Gysin, H. (to Geigy, J. R. A.-G.), U. S. Patent 2,875,126; *Chem. Abstr.*, **53**, 15,101b (1959).
- (319) Hodel, E., and Gysin, H. (to Geigy J. R. A.-G.), British Patent 812,177; *Chem. Abstr.*, **54**, 9963i (1960).
- (320) Hodgkins, J. E., and Ettlinger, M. G., *J. Org. Chem.*, **21**, 404 (1956).
- (321) Hodgkins, J. E., Reeves, W. P., and Liu, Y.-T. G., *J. Am. Chem. Soc.*, **83**, 2532 (1961).
- (322) Hodogaya Chemical Co., Ltd., Japanese Patent 10,505 (1960); *Chem. Abstr.*, **55**, 10,390b (1961).
- (323) Hoeflake, J. M. A., *Rev. trav. chim.*, **40**, 488 (1921).
- (324) Hofman, F., *Z. angew. chem.*, **21**, 1986 (1908).
- (325) Holden, C. B. (to Columbia-Southern Chemical Corp.), U. S. Patent 2,732,914; *Chem. Abstr.*, **50**, 13,081e (1956).
- (326) Holmberg, B., *J. prakt. Chem.*, [2] **71**, 264 (1905).
- (327) Hood, H. P., and Murdock, H. R., *J. Phys. Chem.*, **23**, 498 (1919).
- (328) Hopff, H., and Uffer, H. (to I. G. Farbenindustrie A.-G.), German Patent 907,599 (1954).
- (329) Hopkins, T., *J. Chem. Soc.*, **117**, 278 (1920).
- (330) Hopkins, T. R., Neighbors, R. P., Strickler, P. D., and Phillips, L. V., *J. Org. Chem.*, **24**, 2040 (1959).
- (331) Hopkins, T. R., and Pullen, J. W. (to Spencer Chemical Co.), U. S. Patent 2,906,614; *Chem. Abstr.*, **54**, 2649g (1960).
- (332) Hoshimo, T., and Ichikiazaki, I., *Chem. High Polymers* (Tokyo), **2**, 328 (1945); *Chem. Abstr.*, **44**, 5150a (1950).
- (333) Houben, J., *Ber.*, **36**, 3087 (1903).
- (334) Houben-Weyl, "Methoden der Organischen Chemie," Vol. 8, Georg Thieme Verlag, Stuttgart, 1952, pp. 101-105.
- (335) Houssa, A. J. H., and Phillips, H., *J. Chem. Soc.*, 2510 (1929).
- (336) Houssa, A. J. H., and Phillips, H., *J. Chem. Soc.*, 2513 (1929).
- (337) Houssa, A. J. H., and Phillips, H., *J. Chem. Soc.*, 108 (1932).
- (338) Houssa, A. J. H., and Phillips, H., *J. Chem. Soc.*, 1232 (1932).
- (339) Hunter, H., *J. Chem. Soc.*, **125**, 1389 (1924).
- (340) Iarovenko, N. N., and Vasilieva, A. S., *J. Gen. Chem. USSR*, **28**, 2539 (1958); *Chem. Abstr.*, **53**, 3116h (1959).
- (341) Imperial Chemical Industries, Ltd., British Patent 588,660; *Chem. Abstr.*, **41**, 7162h (1947).
- (342) I. G. Farbenindustrie A.-G., Belgian Patent 450,026; *Chem. Abstr.*, **42**, 214d (1948).
- (343) I. G. Farbenindustrie A.-G., French Patent 895,395.
- (344) I. G. Farbenindustrie A.-G., French Patent 961,754; *Chem. Abstr.*, **46**, 6148i (1952).
- (345) I. G. Farbenindustrie A.-G., German Patent 857,806; *Chem. Abstr.*, **50**, 1901e (1956).
- (346) Imperial Chemical Industries, Ltd., French Patent 1,272,-377 (1962).
- (347) Issleib, K., and Anhoeck, H., *Z. Naturforsch.*, **16b**, 837 (1961); *Chem. Abstr.*, **57**, 859h (1962).
- (348) Ito, Y., and Hayashida, K. (to Hodogaya Chemical Co., Ltd.), Japanese Patent 10,505 (1960); *Chem. Abstr.*, **55**, 10,390b (1961).
- (349) Iwakura, Y., Ishizuka, Y., and Saito, T., *Nippon Kagaku Zasshi*, **76**, 1108 (1955); *Chem. Abstr.*, **51**, 17,750f (1957).
- (350) Iwakura, Y., et al., *Nippon Kagaku Zasshi*, **80**, 78 (1959); *Chem. Abstr.*, **55**, 4527g (1961).
- (351) Iwakura, Y., and Nabeya, A., *J. Org. Chem.*, **25**, 1118 (1960).
- (352) Iwakura, Y., and Tezuka, Y., *Sen-i Gakkaishi*, **17**, 794 (1961); *Chem. Abstr.*, **55**, 24,710f (1961).
- (353) Iwakura, Y., and Tezuka, Y., *Sen-i Gakkaishi*, **17**, 798 (1961); *Chem. Abstr.*, **55**, 24,710h (1961).
- (354) Iwakura, Y., Sakamoto, M., and Yasuda, H., *Nippon Kagaku Zasshi*, **82**, 606 (1961); *Chem. Abstr.*, **56**, 8534a (1962).
- (355) Jaeger, F. M., *Z. anorg. allgem. Chem.*, **101**, 67 (1917).

- (356) Jatkar, S. K. K., and Phansalkar, V. K., *J. Univ. Poona, Sci. Technol.*, **4**, 45 (1953); *Chem. Abstr.*, **48**, 11,859d (1954).
- (357) Job, A., and Dussollier, G., *Compt. rend.*, **184**, 1454 (1927).
- (358) Joglekar, M. S., *Phil. Mag.*, **24**, 405 (1937).
- (359) Johnson, D. A., *J. Am. Chem. Soc.*, **75**, 3636 (1953).
- (360) Jones, J. I., *J. Chem. Soc.*, 2735 (1957).
- (361) Jones, W. D. (to Celanese Corporation of America), U. S. Patent 2,630,302; *Chem. Abstr.*, **47**, 5183d (1953).
- (362) Jones, W. D., and McFarlane, S. B. (to Celanese Corporation of America); U. S. Patent 2,660,574; *Chem. Abstr.*, **48**, 9751i (1954).
- (363) Jones, W. D., and McFarlane, S. B. (to Celanese Corporation of America), U. S. Patent 2,660,575; *Chem. Abstr.*, **48**, 4255c (1954).
- (364) Joshi, B. S., Srinivasan, R., Talavdekar, R. V., and Venkataraman, K., *Tetrahedron*, **11**, 133 (1960).
- (365) Jung, S. L. (to E. I. du Pont de Nemours and Co., Inc.), U. S. Patent 2,962,470; *Chem. Abstr.*, **55**, 6903h (1961).
- (366) Kagarise, R. E., *J. Am. Chem. Soc.*, **77**, 1377 (1955).
- (367) Kahn, D. E., Yorston, F. H., and Cochinaris, N. E., *J. Chem. Soc.*, 1072 (1935).
- (368) Kaiser, D. W., and Thurston, J. T., *J. Org. Chem.*, **17**, 185 (1952).
- (369) Kaji, A., *Bull. Chem. Soc. Japan*, **34**, 254 (1961).
- (370) Kaluza, L., *Monatsh.*, **33**, 363 (1912).
- (371) Karvonen, *Ann. Acad. Sci. Fenniae*, **A10**, No. 4, 19; *Chem. Zentr.*, III, 808 (1919).
- (372) Kashima, M., *Bull. Chem. Soc. Japan*, **25**, 79 (1952); *Chem. Abstr.*, **47**, 1999h (1953).
- (373) Katz, M., and Wittbecker, E. L. (to E. I. du Pont de Nemours and Co., Inc.), U. S. Patent 2,787,630; *Chem. Abstr.*, **51**, 12,977a (1957).
- (374) Katz, M., *J. Polymer Sci.*, **40**, 337 (1959).
- (375) Katz, M. (to E. I. du Pont de Nemours and Co., Inc.), U. S. Patent 2,929,802; *Chem. Abstr.*, **54**, 17,937b (1960).
- (376) Katz, M., and Wittbecker, E. L. (to E. I. du Pont de Nemours and Co., Inc.), U. S. Patent 2,973,333; *Chem. Abstr.*, **55**, 17,082d (1961).
- (377) Kempf, T., *J. prakt. Chem.*, [2] **1**, 402 (1870).
- (378) Kern, W., Rauterkraus, K. J., and Weber, W., *Makromol. Chem.*, **43**, 98 (1961).
- (379) Kharasch, M. S., Lin, Y. C., and Nudenberg, W., *J. Org. Chem.*, **19**, 1150 (1954).
- (380) Kharasch, M. S., Weinhouse, S., and Jensen, E. V., *J. Am. Chem. Soc.*, **77**, 3145 (1955).
- (381) Kinugawa, J., Ochiai, M., and Yamamoto, H., *Yakugaku Zasshi*, **79**, 933 (1959); *Chem. Abstr.*, **54**, 497h (1960).
- (382) Kitano, H., and Fukui, K., *J. Chem. Soc. Japan*, **58**, 603 (1955); *Chem. Abstr.*, **50**, 11,262i (1956).
- (383) Kjaer, A., and Jensen, R. B., *Acta Chem. Scand.*, **10**, 141 (1956).
- (384) Kjaer, A., Gemlin, R., and Jensen, R. B., *Acta Chem. Scand.*, **10**, 432 (1956).
- (385) Kjaer, A., and Gemlin, R., *Acta Chem. Scand.*, **10**, 1193 (1956).
- (386) Kjaer, A., and Gemlin, R., *Acta Chem. Scand.*, **11**, 906 (1957).
- (387) Kleiner, H., and Lehman, W. (to Farbenfabriken Bayer A.-G.), U. S. Patent 2,779,686; *Chem. Abstr.*, **51**, 8448d (1957).
- (388) Klepl, A., *J. prakt. Chem.*, [2] **26**, 447 (1882).
- (389) Kling, A., Florentin, D., Lassieur, A., and Schmutz, R., *Compt. rend.*, **169**, 1046 (1919).
- (390) Kling, A., Florentin, D., Lassieur, A., and Schmutz, R., *Compt. rend.*, **169**, 1166 (1919).
- (391) Kling, A., Florentin, D., Lassieur, A., and Schmutz, R., *Ann. Chim.*, [9] **13**, 44 (1920).
- (392) Kling, A., Florentin, D., and Jacob, E., *Ann. Chim.*, [9] **14**, 189 (1920).
- (393) Kling, A., and Rouilly, M., *Compt. rend.*, **201**, 782 (1935).
- (394) Knabe, J., *Arch. Pharm.*, **289**, 479 (1956); *Chem. Abstr.*, **51**, 5777a (1957).
- (395) Knoll and Co., German Patent 117,267; *Friedländer*, **6**, 146 (1900).
- (396) Knunyants, I. L., Kil'disheva, O. V., and Bykhovskaya, E., *Zh. Obshch. Khim.*, **19**, 101 (1949); *Chem. Abstr.*, **43**, 6164c (1949).
- (397) Kobbe, H. (to Badische Anilin- und Soda-Fabrik A.-G.), German Patent 805,396; *Chem. Abstr.*, **45**, 9269c (1951).
- (398) Kodak, Soc. Anon., Belgian Patent 593,044; *Chem. Abstr.*, **55**, 18,683g (1961).
- (399) Kohlrausch, K. W. F., and Pongratz, A., *Z. physik. Chem.*, **B22**, 373 (1933).
- (400) Kohlrausch, K. W. F., and Sabathy, R., *Monatsh.*, **72**, 303 (1939); *Chem. Abstr.*, **33**, 9138<sup>a</sup> (1939).
- (401) Koller, C. R. (to E. I. du Pont de Nemours and Co., Inc.), U. S. Patent 2,929,801; *Chem. Abstr.*, **54**, 17,936i (1960).
- (402) Koontz, F. H., and Schaefgen, J. R. (to E. I. du Pont de Nemours and Co., Inc.), U. S. Patent 2,915,505; *Chem. Abstr.*, **54**, 4056h (1960).
- (403) Korshak, V. V., and Kolesnikov, G. S., *J. Gen. Chem. USSR*, **14**, 435 (1944); *Chem. Abstr.*, **39**, 4595<sup>b</sup> (1945).
- (404) Kost, A. N., *Vestn. Mosk. Univ.*, No. 2, 141 (1947); *Chem. Abstr.*, **42**, 3722g (1948).
- (405) Kost, A. N., *Uch. Zap. Mosk. Gos. Univ.*, **131**, 39 (1950); *Chem. Abstr.*, **47**, 9905i (1953).
- (406) Kovalenko, V. I., *J. Gen. Chem. USSR*, **22**, 1546 (1952); *Chem. Abstr.*, **47**, 8019 (1953).
- (407) Kraft, F., *Ber.*, **23**, 2785 (1890).
- (408) Kraft, M. Ya., and Lyutina, F. V., *J. Gen. Chem. USSR*, **1**, 190 (1931); *Chem. Abstr.*, **26**, 2167 (1932).
- (409) Kraft, M. Ya., and Aleksov, B. A., *J. Gen. Chem. USSR*, **2**, 726 (1932); *Chem. Abstr.*, **27**, 2426<sup>c</sup> (1933).
- (410) Krutman, A. I., *J. Gen. Chem. USSR*, **22**, 1342 (1952); *Chem. Abstr.*, **47**, 6337d (1953).
- (411) Krzikalla, H., and Merkel, K. (to Badische Anilin- und Soda-Fabrik A.-G.), German Patent 800,662; *Chem. Abstr.*, **45**, 1620g (1951).
- (412) Krzikalla, H., and Bauer, E. (to I. G. Farbenindustrie A.-G.), German Patent 801,989; *Chem. Abstr.*, **45**, 3622b (1951).
- (413) Kucherov, V. F., and Kocheshkov, K. A., *J. Gen. Chem. USSR*, **16**, 1137 (1946); *Chem. Abstr.*, **41**, 2703b (1947).
- (414) Kunckell, F., and Ulex, G., *J. prakt. Chem.*, [2] **86**, 518 (1912).
- (415) Kunckell, F., and Ulex, G., *J. prakt. Chem.*, [2] **87**, 227 (1913).
- (416) Kung, F. E. (to Pittsburgh Plate Glass Co.), U. S. Patent 2,370,589; *Chem. Abstr.*, **39**, 4891<sup>d</sup> (1945).
- (417) Kung, F. E., U. S. Patent 2,377,085; *Chem. Abstr.*, **39**, 3792<sup>e</sup> (1945).
- (418) Kurkjy, R. P., and Matzner, M., private communication.
- (419) LaLande, W. A., and Green, H. (to Pennsalt Manufacturing Co.), U. S. Patent 2,493,075; *Chem. Abstr.*, **44**, 3298i (1950).
- (420) Lambrech, J. A. (to Union Carbide Corp.), U. S. Patent 2,903,478; *Chem. Abstr.*, **54**, 2293c (1960).
- (421) Lambrech, J. A. (to Union Carbide Corp.), British Patent 841,141; *Chem. Abstr.*, **55**, 2591b (1961).
- (422) Langer, P., Prost, M., and Charlier, R., *Helv. Chim. Acta*, **42**, 2379 (1959).

- (423) Langer, P., Prost, M., and Charlier, R., *Helv. Chim. Acta*, **42**, 2394 (1959).
- (424) Lastovskii, R. P., USSR Patent 118,503; *Chem. Abstr.*, **53**, 22,016d (1959).
- (425) Leffler, M. T., and Matson, E. J., *J. Am. Chem. Soc.*, **70**, 3439 (1948).
- (426) Leimu, R., *Ber.*, **B70**, 1040 (1937).
- (427) Leister, N. A., and Tarbell, D. S., *J. Org. Chem.*, **23**, 1152 (1958).
- (428) Lessig, E. T., *J. Phys. Chem.*, **36**, 2325 (1932).
- (429) Lewis, E. S., and Herndon, W. C., Abstracts of Papers presented at the 133rd National Meeting of the American Chemical Society, San Francisco, Calif., April 1958, p. 44N.
- (430) Lewis, E. S., and Herndon, W. C., *J. Am. Chem. Soc.*, **83**, 1955 (1961).
- (431) Lewis, E. S., Herndon, W. C., and Duffey, D. C., *J. Am. Chem. Soc.*, **83**, 1959 (1961).
- (432) Lewis, E. S., and Herndon, W. C., *J. Am. Chem. Soc.*, **83**, 1961 (1961).
- (433) Lieb, H., Verdino, A., and Schadendorff, E., *Ann.*, **512**, 89 (1934); *Chem. Abstr.*, **29**, 172<sup>3</sup> (1935).
- (434) Lipovich, I. M., *Zh. Obsch. Khim.*, **23**, 811 (1953); *Chem. Abstr.*, **48**, 3940f (1954).
- (435) Lipovich, I. M., *J. Gen. Chem. USSR*, **23**, 849 (1953); *Chem. Abstr.*, **48**, 12,715h (1954).
- (436) Little, Arthur D., Inc., Belgian Patent 602,620 (1961).
- (437) Longosz, E. J., and Tarbell, D. S., *J. Org. Chem.*, **26**, 2161 (1961).
- (438) Lowenberg, *Chem. Ztg.*, 390 (1886).
- (439) Ludwig, B. J., and Piech, E. C., *J. Am. Chem. Soc.*, **73**, 5779 (1951).
- (440) Ludwig, B. J., and Piech, E. C., *J. Am. Chem. Soc.*, **73**, 5894 (1951).
- (441) Lyman, D. J., and Jung, S. L., *J. Polymer Sci.*, **40**, 407 (1959).
- (442) Lynn, J. W., *J. Org. Chem.*, **26**, 2119 (1961).
- (443) Mackareth, F. J. H. (to Imperial Chemical Industries, Ltd.), British Patent 588,660; *Chem. Abstr.*, **41**, 7162h, (1947).
- (444) Magat, E. E., and Strachan, D. R. (to E. I. du Pont de Nemours and Co., Inc.), U. S. Patent 2,708,617; *Chem. Abstr.*, **51**, 4759a (1957).
- (445) Mahala, M. M., and Deshpandale, S. S., *J. Indian Chem. Soc.*, **26**, 429 (1949); *Chem. Abstr.*, **44**, 3897g (1950).
- (446) Malatesta, P., *Ric. Sci.*, **22**, 1760 (1952); *Chem. Abstr.*, **48**, 1952h (1954).
- (447) Malinovsky, M. S., and Nadyantseva, N. M., *J. Gen. Chem. USSR*, **23**, 221 (1953); *Chem. Abstr.*, **48**, 2580c (1954).
- (448) Martin, H., Zubiani, R., and Koebel, K. (to Cilag-Chemie, Ltd.), Swiss Patent 343,889; *Chem. Abstr.*, **55**, 1530a (1961).
- (449) Martin, R. H., and Robinson, R., *J. Chem. Soc.*, 497 (1943).
- (450) Masaguer, J. R., *Anales Real Soc. Espan. Fis. Quim. (Madrid)*, **B51**, 705 (1955); *Chem. Abstr.*, **50**, 6237e (1956).
- (451) Matuszak, M. P., *J. Am. Chem. Soc.*, **56**, 2007 (1934).
- (452) Mazurewitsch, I., *J. Russ. Phys.-Chem. Soc.*, **42**, 1582 (1910); *Chem. Zentr.*, **1**, 1500 (1911).
- (453) McKay, A. F., and Brown, R. O., *J. Org. Chem.*, **16**, 1829 (1951).
- (454) McKay, A. F., and Vavasour, G. R., *Can. J. Chem.*, **31**, 688 (1953); *Chem. Abstr.*, **48**, 8238h (1954).
- (455) McLamore, W. M., P'An, S. Y., and Bavley, A., *J. Org. Chem.*, **20**, 1379 (1955).
- (456) Medlock, H., *Ann.*, **69**, 214 (1849).
- (457) Medlock, H., *Ann.*, **69**, 217 (1849) (see footnote on p. 219 by A. W. Hoffmann).
- (458) Medved, T. Ya., and Kabachnik, M. J., *Izv. Akad. Nauk SSSR, Otd. Khim. Nauk*, 684 (1956); *Chem. Abstr.*, **51**, 1818a (1957).
- (459) Meister, Lucius, and Brunning, *J. Chem. Soc.*, **7**, 205 (1888).
- (460) Meister, Lucius, and Brunning, German Patent 232,879; *Friedländer*, **10**, 1241 (1911).
- (461) Meister, Lucius, and Brunning, German Patent 256,843; *Friedländer*, **11**, 1034 (1913).
- (462) Melamed, S. (to Rohm and Haas Co.), U. S. Patent 2,920,-075; *Chem. Abstr.*, **54**, 11,058i (1960).
- (463) Merck, E., and Co., German Patent 247,453; *Chem. Zentr.*, **II**, 162 (1912).
- (464) Merck, E., and Co., German Patent 251,805; *Friedländer*, **11**, 82 (1912).
- (465) Merck, E., and Co., German Patent 254,471; *Friedländer*, **11**, 949 (1912).
- (466) Merck, E., and Co., German Patent 254,472; *Friedländer*, **11**, 948 (1912).
- (467) Merkel, K. (to Badische Anilin- und Soda-Fabrik A.-G.), German Patent 1,082,405; *Chem. Abstr.*, **55**, 13,923c (1961).
- (468) Merrill, S. H., *J. Polymer Sci.*, **55**, 343 (1961).
- (469) Meyer, V., *Ber.*, **2**, 298 (1869).
- (470) Michael, A., *J. prakt. Chem.*, [2] **37**, 473 (1888).
- (471) Michael, A., *J. prakt. Chem.*, [2] **45**, 580 (1892).
- (472) Michael, A., and Carlson, G. H., *J. Am. Chem. Soc.*, **58**, 353 (1936).
- (473) Michael, A., and Welner, N., *J. Org. Chem.*, **3**, 372 (1938).
- (474) Michaelis, L., *Ber.*, **24**, 3439 (1891).
- (475) Michejda, C. J., Tarbell, D. S., and Saunders, W. H., Jr., *J. Am. Chem. Soc.*, **84**, 4113 (1962).
- (476) Migushima, S., and Kubo, M., *Bull. Chem. Soc. Japan*, **13**, 174 (1938); *Chem. Abstr.*, **32**, 4399<sup>2</sup> (1938).
- (477) Miller, J. A., *Ber.*, **22**, 2790 (1889).
- (478) Mingoia, Q., *Gazz. chim. ital.*, **55**, 713 (1925).
- (479) Mingoia, Q., *Gazz. chim. ital.*, **58**, 667 (1928).
- (480) Minnesota Mining and Manufacturing Co., French Patent 1,273,547 (1962).
- (481) Mohler, H., *Helv. Chim. Acta*, **21**, 787 (1938); *Chem. Abstr.*, **32**, 6764<sup>b</sup> (1938).
- (482) Momose, T., Shoji, T., and Iwaskaki, M., *Yakugaku Zasshi*, **81**, 1045 (1961); *Chem. Abstr.*, **55**, 27,177h (1961).
- (483) Morel, A., *Bull. soc. chim. France*, [3] **21**, 815 (1899).
- (484) Morel, A., *Bull. soc. chim. France*, [3] **21**, 823 (1899).
- (485) Moureau, C., and Delange, R., *Bull. soc. chim. France*, [3] **29**, 648 (1903).
- (486) Moureau, C., and Delange, R., *Compt. rend.*, **136**, 552 (1903).
- (487) Muller, A. C., and Shokal, E. C. (to Shell Development Co.), U. S. Patent 2,795,572; *Chem. Abstr.*, **51**, 13,464c (1957).
- (488) Muller, E., and Bayer, O. (to Farbenfabriken Bayer A.-G.), U. S. Patent 2,999,844; *Chem. Abstr.*, **56**, 3639a (1962).
- (489) Muller, H., *Ann.*, **258**, 50 (1890).
- (490) Mumford, S. A., and Phillips, J. W. C., *J. Chem. Soc.*, 2112 (1929).
- (491) Muskat, I. E., and Strain, F. (to Pittsburgh Plate Glass Co.), U. S. Patent 2,370,565; *Chem. Abstr.*, **39**, 4526<sup>g</sup> (1945).
- (492) Muskat, I. E., and Strain, F. (to Pittsburgh Plate Glass Co.), U. S. Patent 2,370,566; *Chem. Abstr.*, **39**, 4527<sup>4</sup> (1945).

- (493) Muskat, I. E., and Strain, F. (to Pittsburgh Plate Glass Co.), U. S. Patent 2,370,567; *Chem. Abstr.*, **39**, 4087<sup>s</sup> (1945).
- (494) Muskat, I. E., and Strain, F. (to Pittsburgh Plate Glass Co.), U. S. Patent 2,370,568; *Chem. Abstr.*, **39**, 4087<sup>s</sup> (1945).
- (495) Muskat, I. E. (to Pittsburgh Plate Glass Co.), U. S. Patent 2,370,570; *Chem. Abstr.*, **39**, 4087<sup>s</sup> (1945).
- (496) Muskat, I. E., and Strain, F. (to Pittsburgh Plate Glass Co.), U. S. Patent 2,370,572; *Chem. Abstr.*, **39**, 4625<sup>t</sup> (1945).
- (497) Muskat, I. E., and Strain, F. (to Pittsburgh Plate Glass Co.), U. S. Patent 2,370,573; *Chem. Abstr.*, **39**, 3974<sup>s</sup> (1945).
- (498) Muskat, I. E., and Strain, F. (to Pittsburgh Plate Glass Co.), U. S. Patent 2,379,250; *Chem. Abstr.*, **39**, 4623<sup>t</sup> (1945).
- (499) Muskat, I. E., and Strain, F. (to Pittsburgh Plate Glass Co.), U. S. Patent 2,379,252; *Chem. Abstr.*, **39**, 4623<sup>t</sup> (1945).
- (500) Muskat, I. E., and Strain, F. (to Pittsburgh Plate Glass Co.), U. S. Patent 2,384,115; *Chem. Abstr.*, **40**, 2457<sup>t</sup> (1946).
- (501) Muskat, I. E., and Strain, F. (to Pittsburgh Plate Glass Co.), U. S. Patents 2,384,116–2,384,126; *Chem. Abstr.*, **40**, 2457<sup>t</sup>–2461<sup>t</sup> (1946).
- (502) Muskat, I. E., and Strain, F. (to Pittsburgh Plate Glass Co.), U. S. Patents 2,385,931–2,385,934; *Chem. Abstr.*, **40**, 2461<sup>t</sup>–2462<sup>t</sup> (1946).
- (503) Muskat, I. E., and Strain, F. (to Pittsburgh Plate Glass Co.), U. S. Patents 2,387,931–2,387,934; *Chem. Abstr.*, **40**, 2462<sup>t</sup>–2464<sup>t</sup> (1946).
- (504) Muskat, I. E., and Strain, F. (to Pittsburgh Plate Glass Co.), U. S. Patent 2,403,113; *Chem. Abstr.*, **40**, 5771<sup>t</sup> (1946).
- (505) Muskat, I. E., and Strain, F. (to Pittsburgh Plate Glass Co.), U. S. Patent 2,592,058; *Chem. Abstr.*, **46**, 8417<sup>g</sup> (1952).
- (506) Nagel, K., and Korn, F. (to Farbenfabriken Bayer A.-G.), German Patent 1,080,546; *Chem. Abstr.*, **55**, 17,510c (1961).
- (507) Naiki, K., *Yuki Gosei Kagaku Kyokai Shi*, **14**, 84 (1956); *Chem. Abstr.*, **51**, 7018d (1957).
- (508) Najar, H., Chabrier, P., and Giudicelli, R., *Bull. soc. chim. France*, 1189 (1955); *Chem. Abstr.*, **51**, 10,424g (1957).
- (509) Najar, H., Giudicelli, R., Chabrier, P., Joanic-Voisinet, E., and Sette, J., *Ann. Pharm. Franç.*, **17**, 200 (1959); *Chem. Abstr.*, **54**, 3189e (1960).
- (510) Najar, H., Chabrier, P., and Giudicelli, R., *Bull. soc. chim. France*, 611 (1959).
- (511) Najar, H., Chabrier, P., and Giudicelli, R., *Bull. soc. chim. France*, 1611 (1959).
- (512) Nakanishi, S., Myers, T. C., and Jensen, E. V., *J. Am. Chem. Soc.*, **77**, 3099 (1955).
- (513) Nakanishi, S., Myers, T. C., and Jensen, E. V., *J. Am. Chem. Soc.*, **77**, 5033 (1955).
- (514) Nazarov, I. N., Sokolov, D. V., and Litvinenko, G. S., *Izv. Akad. Nauk SSSR, Otd. Khim. Nauk*, 95 (1954); *Chem. Abstr.*, **49**, 6520c (1955).
- (515) Nazarov, I. N., Sokolov, D. V., and Litvinenko, G. S., *Bull. Acad. Sci. USSR, Div. Chem. Sci.*, **77** (1954); *Chem. Abstr.*, **49**, 8966b (1955).
- (516) Nazrov, I. N., and Shvekhgeimer, G. A., *J. Gen. Chem. USSR*, **29**, 464 (1959); *Chem. Abstr.*, **53**, 21,661c (1959).
- (517) Nef, J. U., *Ann.*, **266**, 52 (1891).
- (518) Nef, J. U., *Ann.*, **276**, 200 (1893).
- (519) Nekrassow, W., and Komissarow, J. F., *J. prakt. Chem.*, [2] **123**, 180 (1929); *Chem. Abstr.*, **24**, 62 (1930).
- (520) Nelson, J. A., Miller, T. G., and Smeltz, K. C. (to E. I. du Pont de Nemours and Co., Inc.), U. S. Patent 2,959,-611; *Chem. Abstr.*, **55**, 22,139a (1961).
- (521) Nemirovsky, J., *J. prakt. Chem.*, [2] **31**, 173 (1885).
- (522) Nencki, M., *Ber.*, **7**, 1588 (1874).
- (523) Nesmeyanov, A. N., and Reutov, O. A., *Izv. Akad. Nauk SSSR, Otd. Khim. Nauk*, 611 (1949); *Chem. Abstr.*, **44**, 7791h (1950).
- (524) Nesmeyanov, A. N., et al., *Izv. Akad. Nauk SSSR, Otd. Khim. Nauk*, 631 (1949).
- (525) Newmann, S. R., Heisler, R. Y., and Alpert, N. (to Texaco, Inc.), U. S. Patent 2,921,955; *Chem. Abstr.*, **54**, 22,360h (1960).
- (526) Nishide, M., Onodera, N., and Sera, M., *Kogyo Kagaku Zasshi*, **63**, 1464 (1960); *Chem. Abstr.*, **56**, 13,079i (1962).
- (527) Norton, F. H. (to Dow Chemical Co.), U. S. Patent 2,923,-732; *Chem. Abstr.*, **54**, 8674d (1960).
- (528) Nowak, K., Siemon, I. Z., and Siemeniewski, H., *Roczniki Chem.*, **36**, 557 (1962); *Chem. Abstr.*, **57**, 12,613h (1962).
- (529) Oesper, R. E., Broker, W., and Cook, W. A., *J. Am. Chem. Soc.*, **47**, 2609 (1925).
- (530) O'Gorman, J. M., Shand, W., Jr., and Schomaker, V., *J. Am. Chem. Soc.*, **72**, 4222 (1950).
- (531) Olah, G., and Pavlath, A., *Acta Chim. Acad. Sci. Hung.*, **3**, 191 (1953); *Chem. Abstr.*, **48**, 7533b (1954).
- (532) Olah, G., and Pavlath, A., *Acta Chim. Acad. Sci. Hung.*, **4**, 89 (1954); *Chem. Abstr.*, **49**, 6095e (1955).
- (533) Olivier, K. L., and Young, W. G., *J. Am. Chem. Soc.*, **81**, 5811 (1959).
- (534) Olomucki, M., *Bull. soc. chim. France*, 186 (1959).
- (535) Oppenheimer, E., *Ber.*, **22**, 2442 (1889).
- (536) Otto, P., *J. prakt. Chem.*, [2] **44**, 15 (1891).
- (537) Otto, R., and Rossing, A., *Ber.*, **18**, 2493 (1885).
- (538) Otto, R., and Otto, W., *Ber.*, **21**, 1516 (1888).
- (539) Owen, T.-Y., and Kao, Y.-S., *Hua Hsueh Hsueh Pao*, **25**, 312 (1959); *Chem. Abstr.*, **54**, 16,418c (1960).
- (540) Palomaa, N. H., and Leimu, R., *Ber.*, **B66**, 813 (1933).
- (541) Pajaczkowski, A., and Turner, J. M. (to Canadian Industries, Ltd., and Imperial Chemical Industries, Ltd.), British Patent 870,584; *Chem. Abstr.*, **56**, 4952c (1962).
- (542) Panneman, H. J., Marx, A. F., and Arens, J. F., *Rev. trav. chim.*, **78**, 487 (1959).
- (543) Parfent'ev, L. N., and Shamshurin, A. A., *Tr. Uzbeksk. Gos. Univ. Sbornik Rabot Khim.*, **15**, 67 (1939); *Chem. Abstr.*, **35**, 4351e (1941).
- (544) Pasedack, H. (to Badische Anilin- und Soda-Fabrik A.-G.), German Patent 837,699; *Chem. Abstr.*, **51**, 14,819h (1957).
- (545) Pawlewski, B., *Ber.*, **17**, 1205 (1884).
- (546) Pawlewski, B., *Ber.*, **25**, 1449 (1892).
- (547) Payal and Montagne, *Bull. soc. chim. France*, [4] **27**, 857 (1920).
- (548) Pechukas, A. (to Pittsburgh Plate Glass Co.), U. S. Patent 2,464,056; *Chem. Abstr.*, **43**, 5230b (1949).
- (549) Pennsalt Manufacturing Co., French Patent 970,947; *Chem. Abstr.*, **46**, 6428g (1952).
- (550) Perret, A., and Biechler, J., *Compt. rend.*, **203**, 84 (1936).
- (551) Perret, A., and Biechler, J., *Bull. soc. ind. Mulhouse*, **103**, 168 (1938); *Chem. Abstr.*, **32**, 2087<sup>s</sup> (1938).
- (552) Perron, Y. G., Pindell, M. H., Cast, L. B., and Cheney, L. C., *J. Med. Pharm. Chem.*, **4**, 41 (1961).
- (553) Pfeiffer, H., and Junkmann, K. (to Schering A.-G.), German Patent 1,007,764; *Chem. Abstr.*, **53**, 19,892i (1959).

- (554) Pfeiffer, H., and Junkmann, K. (to Schering A.-G.), German Patent 1,011,412; *Chem. Abstr.*, **54**, 2215d (1960).
- (555) Pianfetti, J. A., Williams, O. F., and Allen, J. F. (to Food Machinery and Chemical Corp.), U. S. Patent 2,786,009; *Chem. Abstr.*, **51**, 9998b (1957).
- (556) Pickard, R. H., and Littlebury, W. O., *J. Chem. Soc.*, **91**, 300 (1907).
- (557) Pierce, J. S., and Adams, R., *J. Am. Chem. Soc.*, **45**, 790 (1923).
- (558) Pierce, J. S., *J. Am. Chem. Soc.*, **50**, 241 (1928).
- (559) Pinchas, S., and Ben-Ishai, D., *J. Am. Chem. Soc.*, **79**, 4099 (1957).
- (560) Pinck, L. A., and Blair, J. S., *J. Am. Chem. Soc.*, **49**, 509 (1927).
- (561) Pittsburgh Plate Glass Co., British Patent 587,933; *Chem. Abstr.*, **42**, 206i (1948).
- (562) Pogosyan, G. M., and Matsoyan, S. G., *Izv. Akad. Nauk Arm. SSR, Khim. Nauk.*, **14**, 343 (1961); *Chem. Abstr.*, **57**, 7456c (1962).
- (563) Ponziò, G., and Bernardi, V., *Gazz. chim. ital.*, **53**, 813 (1923); *Chem. Abstr.*, **18**, 1490 (1924).
- (564) Putman, R. C. (to United Shoe Machinery, Inc.), U. S. Patent 2,523,324; *Chem. Abstr.*, **45**, 378d (1951).
- (565) Rabjohn, N., *J. Am. Chem. Soc.*, **70**, 1181 (1948).
- (566) Rabjohn, N., Hopkins, T. R., and Nagler, R. C., *J. Am. Chem. Soc.*, **74**, 3215 (1952).
- (567) Raiford, L. C., and Inman, G. O., *J. Am. Chem. Soc.*, **56**, 1586 (1934).
- (568) Raiford, L. C., and Dudley, J. R., *Proc. Iowa Acad. Sci.*, **51**, 313 (1944); *Chem. Abstr.*, **40**, 7178<sup>g</sup> (1946).
- (569) Ramsperger, H. C., and Waddington, G., *J. Am. Chem. Soc.*, **55**, 214 (1933).
- (570) Ravin, H. A., Tsou, K. C., and Seligman, A. M., *J. Biol. Chem.*, **191**, 843 (1951).
- (571) Razuvayev, G. A., and Ol'dekop, Yu. A., *J. Gen. Chem. USSR*, **23**, 1173 (1953); *Chem. Abstr.*, **47**, 12,215d (1953).
- (572) Recherches et Techniques Appliquées and Société Anonyme de Documentation d'Etudes et de Gestion, French Patent 1,176,912; *Chem. Abstr.*, **55**, 19,861e (1961).
- (573) Recherches et Techniques Appliquées and Société Anonyme de Documentation d'Etudes et de Gestion, French Patent Additions 70,903 and 70,904 (to French Patent 1,176,912); *Chem. Abstr.*, **55**, 19,861e (1961); *Chem. Abstr.*, **56**, 7216i-7217e (1962).
- (574) Reformatsky, A., *J. Russ. Phys. Chem. Soc.*, **19**, 298.
- (575) Reformatsky, A., *Ber.*, **41**, 4083 (1908).
- (576) Rehberg, C. E., Dixon, M. B., and Fisher, C. H., *J. Org. Chem.*, **13**, 254 (1948).
- (577) Rennie, E. H., *J. Chem. Soc.*, **41**, 33 (1882).
- (578) Reppe, W., *Ann.*, **596**, 80 (1955).
- (579) "Research" N. V., Belgian Patent 566,669 (1958).
- (580) "Research" N. V., British Patent 843,881; *Chem. Abstr.*, **55**, 2187b (1961).
- (581) Ressler, C., and du Vigneaud, V., *J. Am. Chem. Soc.*, **76**, 3107 (1954).
- (582) Reynolds, D. D., and Laakso, T. M. (to Eastman Kodak Co.), U. S. Patent 2,784,163; *Chem. Abstr.*, **51**, 9219e (1957).
- (583) Reynolds, D. D., and Van den Berghe, J. (to Eastman Kodak Co.), U. S. Patent 2,789,509; *Chem. Abstr.*, **51**, 14,809c (1957).
- (584) Reynolds, D. D., and Van den Berghe, J. (to Eastman Kodak Co.), U. S. Patent 2,789,964; *Chem. Abstr.*, **51**, 14,808c (1957).
- (585) Reynolds, D. D., and Van den Berghe, J. (to Eastman Kodak Co.), U. S. Patent 2,789,970; *Chem. Abstr.*, **51**, 14,808g (1957).
- (586) Reynolds, D. D., Fields, D. L., and Johnson, D. L., *J. Org. Chem.*, **26**, 5119 (1961).
- (587) Reynolds, D. D., Fields, D. L., and Johnson, D. L., *J. Org. Chem.*, **26**, 5130 (1961).
- (588) Richter, E., *Ber.*, **22**, 2449 (1889).
- (589) Riedmair, J., and Kindler, H. (to Casella Farbwerke Mainkur A.-G.), German Patent 960,896; *Chem. Abstr.*, **53**, 16,149h (1959).
- (590) Rivier, H., and Richard, P., *Helv. Chim. Acta*, **8**, 490 (1925).
- (591) Riviere, C., *Ann. Chim.*, **1**, 157 (1946).
- (592) Roberts, L. E., Lashbrook, R., Treat, M. J., and Yates, W., *J. Am. Chem. Soc.*, **74**, 5787 (1952).
- (593) Roemer, H., *Ber.*, **6**, 1101 (1873).
- (594) Roese, B., *Ann.*, **205**, 227 (1880).
- (595) Rogovin, Z. A., and Kozlova, Yu. S., *Zh. Obshch. Khim.*, **29**, 1667 (1959); *Chem. Abstr.*, **54**, 8650f (1960).
- (596) Rohm and Haas Co., British Patent 834,575; *Chem. Abstr.*, **55**, 6438c (1961).
- (597) Rona, M., and Ben-Ishai, D., *J. Org. Chem.*, **26**, 1446 (1961).
- (598) Sachs, C. C., and Bond, J. (to A. H. Kerr Co.), U. S. Patent 2,548,685; *Chem. Abstr.*, **45**, 6430a (1951).
- (599) Salomon, F., *J. prakt. Chem.*, [2] **6**, 433 (1873).
- (600) Sauers, C. K., and Cotter, R. J., *J. Org. Chem.*, **26**, 6 (1961).
- (601) Sauers, C. K., and Cotter, R. J. (to Union Carbide Corp.), U. S. Patent 2,980,701; *Chem. Abstr.*, **55**, 19,869a (1961).
- (602) Saunders, J. H., Slocombe, R. J., and Hardy, E. E., *J. Am. Chem. Soc.*, **73**, 3796 (1951).
- (603) Saville, B., *Analyst*, **82**, 269 (1957).
- (604) Schadendorff, E., and Verdino, A., *Monatsh.*, **65**, 338 (1935); *Chem. Abstr.*, **29**, 3684<sup>g</sup> (1935).
- (605) Schaefgen, J. R., Koontz, F. H., and Tietz, R. F., *J. Polymer Sci.*, **40**, 377 (1959).
- (606) Scheibler, H., and Voss, J., *Ber.*, **53**, 388 (1920).
- (607) Scheibler, H., and Stein, H., *J. prakt. Chem.*, **139**, 105 (1934).
- (608) Scheibler, H., and Scheibler, U., *Chem. Ber.*, **87**, 379 (1954).
- (609) Schiemenz, G. P., and Engelhard, H., *Chem. Ber.*, **92**, 857 (1959).
- (610) Schiff, H., *Ann.*, **291**, 367 (1896).
- (611) Schipper, E., and Nichols, J., *J. Am. Chem. Soc.*, **80**, 5714 (1958).
- (612) Schlack, P. (vested in the Alien Property Custodian), U. S. Patent 2,325,552; *Chem. Abstr.*, **38**, 432<sup>1</sup> (1944).
- (613) Schmerling, L., and Ipatieff, V. N. (to Universal Oil Products Co.), U. S. Patent 2,404,537; *Chem. Abstr.*, **40**, 6252<sup>3</sup> (1946).
- (614) Schnell, H., and Bottenbruch, L. (to Farbenfabriken Bayer A.-G.), German Patent Application F 13040 IVb/39c (Oct. 16, 1953).
- (615) Schnell, H., Bottenbruch, L., and Fritz, G. (to Farbenfabriken Bayer A.-G.), German Patent Application, F 17166 IVb/39c (March 26, 1955).
- (616) Schnell, H., *et al.* (to Farbenfabriken Bayer A.-G.), German Patent Application F 17168 IVb/39c.
- (617) Schnell, H., *et al.* (to Farbenfabriken Bayer A.-G.), German Patent Application F 17528 IVb/39c.
- (618) Schnell, H., Bottenbruch, L., and Krimm, H. (to Farbenfabriken Bayer A.-G.), U. S. Patent 3,028,365; *Chem. Abstr.*, **51**, 10,953i (1957).
- (619) Schnell, H., Kimpel, W., Bottenbruch, L., Krimm, H., and

- Fritz, G. (to Farbenfabriken Bayer A.-G.), German Patent 1,011,148; *Chem. Abstr.*, **54**, 8148b (1960).
- (620) Schnell, H., and Bottenbruch, L. (to Farbenfabriken Bayer A.-G.), German Patent 1,045,657; *Chem. Abstr.*, **55**, 1075f (1961).
- (621) Schnell, H., and Bottenbruch, L., *Makromol. Chem.*, **57**, 1 (1962).
- (622) Scholl, R., and Egner, W., *Ann.*, **397**, 301 (1913).
- (623) Scholtissek, C., *Chem. Ber.*, **89**, 2562 (1956).
- (624) Schopff, M., *Ber.*, **18**, 1063 (1885).
- (625) Schopff, M., *Ber.*, **18**, 2472 (1885).
- (626) Schotte, H., Preiwe, H., and Roescheisen, H., *Z. physiol. Chem. Hoppe-Seyler's*, **174**, 119 (1928); *Chem. Abstr.*, **22**, 1759 (1928).
- (627) Schotten, C., *Ber.*, **15**, 421 (1882).
- (628) Schreiner, L., *J. prakt. Chem.*, [2] **22**, 353 (1880).
- (629) Schubart, L. H., *Ber.*, **22**, 2433 (1889).
- (630) Schwing, P., and Sabetay, S., *Bull. soc. chim. France*, [4] **43**, 857 (1928).
- (631) Scott, G. (to Imperial Chemical Industries, Ltd.), U. S. Patent 2,951,834; *Chem. Abstr.*, **55**, 6036f (1961).
- (632) Shapiro, S. L., Rose, I. M., and Freedman, L., *J. Am. Chem. Soc.*, **81**, 6322 (1959).
- (633) Shemyakin, M. M., Arbuzov, Yu. A., Kosolov, M. N., Shatenstein, G. A., Onoprienko, V. V., and Konnova, Yu. V., *Zh. Obshch. Khim.*, **30**, 542 (1960); *Chem. Abstr.*, **54**, 24,575i (1960).
- (634) Sheppard, W. A., *J. Org. Chem.*, **27**, 3756 (1962).
- (635) Sims, J. G., and Moyle, C. L. (to Dow Chemical Co.), U. S. Patent 2,917,534; *Chem. Abstr.*, **54**, 7652c (1960).
- (636) Slimowicz, C. E., and Degering, E. F., *J. Am. Chem. Soc.*, **71**, 1043 (1949).
- (637) Smith, E. S. (to Goodyear Tire and Rubber Co.), U. S. Patent 3,022,281; *Chem. Abstr.*, **55**, 11,919i (1961).
- (638) Smith, L. H., "Synthetic Fiber Developments in Germany," Textile Research Institute, New York, N. Y., 1946.
- (639) Smyth, C. P., *J. Am. Chem. Soc.*, **47**, 1894 (1925).
- (640) Societe pour l'Ind. Chim. a Bâle, Swiss Patent 219,857; *Chem. Abstr.*, **42**, 7072b (1948).
- (641) Societe pour l'Ind. Chim. a Bâle, Swiss Patent 221,823; *Chem. Abstr.*, **43**, 886d (1949).
- (642) Societe pour l'Ind. Chim. a Bâle, Swiss Patent 221,824; *Chem. Abstr.*, **43**, 441e (1949).
- (643) Societe pour l'Ind. Chim. a Bâle, Swiss Patent 221,825; *Chem. Abstr.*, **43**, 441f (1949).
- (644) Societe pour l'Ind. Chim. a Bâle, Swiss Patent 221,826; *Chem. Abstr.*, **43**, 441f (1949).
- (645) Societe pour l'Ind. Chim. a Bâle, Swiss Patent 223,217; *Chem. Abstr.*, **43**, 1990i (1949).
- (646) Societe anon. d'innovations chimiques, dite: Sinnova ou Sadic, French Patent 1,017,280; *Chem. Abstr.*, **51**, 17-, 206d (1957).
- (647) Spica, M., and DeVarda, G., *Gazz. chim. ital.*, **17**, 165 (1887).
- (648) Spiegler, L. (to E. I. du Pont de Nemours and Co., Inc.), U. S. Patent 2,873,291; *Chem. Abstr.*, **53**, 11,230h (1959).
- (649) Staab, H. A., *Ann.*, **609**, 83 (1957).
- (650) Stefanova, A. V., *Annuaire Univ. Sofia, Fac. Phys.-Math.*, **40**, 147 (1943-1944).
- (651) Stefanye, D., Howard, W. L., and Beidler, W., *J. Am. Chem. Soc.*, **77**, 3663 (1955).
- (652) Stefanye, D., Howard, W. L., Brown, J. W., and Weintraub, R. L. (to U. S. Dept. of the Army), U. S. Patent 2,916,491; *Chem. Abstr.*, **54**, 5716i (1960).
- (653) Steuber, W. (to E. I. du Pont de Nemours and Co., Inc.), U. S. Patent 3,044,990; *Chem. Abstr.*, **57**, 12,758e (1962).
- (654) Stilmar, F. B. (to E. I. du Pont de Nemours and Co., Inc.), U. S. Patent 2,757,191; *Chem. Abstr.*, **51**, 4056f (1957).
- (655) Stolz, F., *J. prakt. Chem.*, [2] **55**, 145 (1897).
- (656) Strain, F. (to Pittsburgh Plate Glass Co.), U. S. Patent 2,370,588; *Chem. Abstr.*, **39**, 4891<sup>b</sup> (1945).
- (657) Strain, F. (to Pittsburgh Plate Glass Co.), U. S. Patent 2,397,630; *Chem. Abstr.*, **40**, 3769<sup>b</sup> (1946).
- (658) Strain, F. (to Pittsburgh Plate Glass Co.), U. S. Patent 2,446,145; *Chem. Abstr.*, **43**, 6652h (1949).
- (659) Strain F., and Newton, W. F. (to Pittsburgh Plate Glass Co.), U. S. Patent 2,476,637; *Chem. Abstr.*, **44**, 655d (1950).
- (660) Strain, F., Bissinger, W. E., Dial, W. R., Rudolf, H., DeWitt, B. J., Stevens, H. C., and Langston, J. H., *J. Am. Chem. Soc.*, **72**, 1254 (1950).
- (661) Strain, F. (to Columbia-Southern Chemical Corp.), U. S. Patent 2,777,871; *Chem. Abstr.*, **51**, 12,969a (1957).
- (662) Stuehmer, W., and Funke, S. (to Kali-Chemie A.-G.), U. S. Patent 2,878,158; *Chem. Abstr.*, **53**, 13,110d (1959).
- (663) Stuehmer, W., and Funke, S. (to Kali-Chemie A.-G.), British Patent 812,256; *Chem. Abstr.*, **54**, 2261e (1960).
- (664) Stuehmer, W., and Funke, S. (to Kali-Chemie A.-G.), German Patent 1,020,017; *Chem. Abstr.*, **54**, 4500b (1960).
- (665) Sweeny, W., *J. Appl. Polymer Sci.*, **5**, No. 16, S15 (1961).
- (666) Swerdluff, J., and McFarlane, S. B., Abstracts of Papers presented at the American Chemical Society Meeting, New York, N. Y., Sept., 1954, p. 145.
- (667) Synievski, V., *Ber.*, **28**, 1874 (1895).
- (668) Tarbell, D. S., and Price, J. A., *J. Org. Chem.*, **21**, 144 (1956).
- (669) Tarbell, D. S., and Price, J. A., *J. Org. Chem.*, **22**, 245 (1957).
- (670) Tarbell, D. S., and Leister, N. A., *J. Org. Chem.*, **23**, 1149 (1958).
- (671) Tarbell, D. S., and Longosz, E. J., *J. Org. Chem.*, **24**, 774 (1959).
- (672) Theis, T. W., and Richardson, K. W., Jr. (to Columbia-Southern Chemical Corp.), U. S. Patent 2,778,846; *Chem. Abstr.*, **51**, 10,563g (1957).
- (673) Theobald, R. S., *J. Chem. Soc.*, 5359 (1961).
- (674) Theobald, R. S., *J. Chem. Soc.*, 5365 (1961).
- (675) Theobald, R. S., *J. Chem. Soc.*, 5370 (1961).
- (676) Thiele, J., and Dent, F., *Ann.*, **302**, 245 (1898).
- (677) Thoma, W., and Rinke, H., *Ann.*, **624**, 30 (1959).
- (678) Tiemann, F., *Ber.*, **18**, 2456 (1885).
- (679) Tiemann, F., *Ber.*, **19**, 1475 (1886).
- (680) Timmermans, J., *Bull. soc. chim. Belges*, **31**, 392 (1923).
- (681) Traube, W., and Gockel, H., *Ber.*, **56**, 384 (1923).
- (682) Traverso, G., *Gazz. chim. ital.*, **87**, 67 (1957).
- (683) Trosken, O. (to Cassella Farbwerke Mainkur A.-G.), German Patent 1,053,492; *Chem. Abstr.*, **55**, 10,412e (1961).
- (684) Tsou, K. C., *J. Am. Chem. Soc.*, **76**, 6108 (1954).
- (685) Tursich, J. A. (to Columbia-Southern Chemical Co.), U. S. Patent 2,788,268; *Chem. Abstr.*, **51**, 11,646h (1957).
- (686) Twiss, D., *J. Am. Chem. Soc.*, **49**, 491 (1927).
- (687) Ugi, T., and Beck, F., *Chem. Ber.*, **94**, 1839 (1961).
- (688) Ulrich, H. (to General Aniline and Film Corp.), U. S. Patent 2,382,185; *Chem. Abstr.*, **40**, 220<sup>b</sup> (1946).
- (689) Ulsch, K., *Ann.*, **226**, 281 (1884).
- (690) Underwood, H. W., Jr., and Baril, O. L., *J. Am. Chem. Soc.*, **53**, 2200 (1931).
- (691) Union Chimique Belge, S. A., Belgian Patent 532,057; *Chem. Abstr.*, **54**, 5572i (1960).

- (692) Van Dorp, D. A., and Arens, J. F., *Rec. trav. chim.*, **65**, 722 (1946).
- (693) Van Dorp, D. A., and Arens, J. F., *Rec. trav. chim.*, **66**, 189 (1947).
- (694) Vaughan, J. R., Jr., *J. Am. Chem. Soc.*, **73**, 3547 (1951).
- (695) Vaughan, J. R., Jr., and Osato, R. L., *J. Am. Chem. Soc.*, **74**, 676 (1952).
- (696) Vaughan, J. R., Jr., *J. Am. Chem. Soc.*, **74**, 6137 (1952).
- (697) Vaughan, J. R., Jr., and Eichler, J. A., *J. Am. Chem. Soc.*, **75**, 5556 (1953).
- (698) Vaughan, J. R., Jr., and Eichler, J. A., *J. Am. Chem. Soc.*, **76**, 2474 (1954).
- (699) Vaughan, W. R., and Spencer, J. L., *J. Org. Chem.*, **25**, 1077 (1960).
- (700) Vavon, G., and Riviere, C., *Compt. rend.*, **220**, 286 (1945).
- (701) Verdino, A., and Schadendorff, E., *Monatsh.*, **66**, 169 (1935).
- (702) du Vigneaud, V., Ressler, C., Swan, J. M., Roberts, C. W., Katsoyannis, P. G., and Gordon, S., *J. Am. Chem. Soc.*, **75**, 4879 (1953).
- (703) Walker, H. G., Jr., Levine, R., Kibler, R. F., and Hauser, C. R., *J. Am. Chem. Soc.*, **68**, 672 (1946).
- (704) Weatherly, T. L., and Williams, Q., *J. Chem. Phys.*, **22**, 958 (1954); *Chem. Abstr.*, **48**, 11,922f (1954).
- (705) Wegler, R., *Ann.*, **498**, 62 (1932).
- (706) Weise, J., *Ber.*, **22**, 2418 (1889).
- (707) Welch, V. A., and Kent, P. W., *J. Chem. Soc.*, 2266 (1962).
- (708) Whaley, W. M., and Meadow, M., *J. Org. Chem.*, **19**, 666 (1954).
- (709) Wheeler, H. L., and Dustin, G. K., *Am. Chem. J.*, **24**, 424 (1900).
- (710) Wiberg, K. B., and Schryne, T. M., *J. Am. Chem. Soc.*, **77**, 2774 (1955).
- (711) Wieland, H., Honold, E., and Pascual-Vila, J., *Z. physiol. Chem.*, **130**, 326 (1923); *Chem. Abstr.*, **18**, 1830 (1924).
- (712) Wieland, H., van Hove, H., and Borner, K., *Ann.*, **446**, 31 (1926).
- (713) Wieland, T., and Bernhard, H., *Ann.*, **572**, 190 (1951).
- (714) Wieland, T., Schaftner, W., and Bokelmann, E., *Ann.*, **573**, 99 (1951).
- (715) Williams, J. L. R., Reynolds, D. D., Dunham, K. R., and Tinker, J. F., *J. Org. Chem.*, **24**, 64 (1959).
- (716) Willersinn, H. (to Badische Anilin- und Soda-Fabrik A.-G.), German Patent 1,100,952; *Chem. Abstr.*, **55**, 27,970g (1961).
- (717) Willstätter, R., Ulbrich, E., Pogany, L., and Maimeri, C., *Ann.*, **477**, 161 (1929).
- (718) Wilm, T., and Wischin, G., *Ann.*, **147**, 150 (1868).
- (719) Wilm, T., and Wischin, G., *Ann.*, **147**, 157 (1868).
- (720) Wilm, T., *Ann.*, **192**, 243 (1878).
- (721) Windholz, T. B., *J. Org. Chem.*, **23**, 2044 (1958).
- (722) Windholz, T. B., Abstracts of Papers presented at the 136th National Meeting of the American Chemical Society, Atlantic City, N. J., Sept., 1959, p. 63P.
- (723) Windholz, T. B., *J. Org. Chem.*, **25**, 1703 (1960).
- (724) Wingfoot Corp., British Patent 613,280; *Chem. Abstr.*, **43**, 4293b (1949).
- (725) Wislicenus, J., and Ehrlich, L., *Ber.*, **7**, 892 (1874).
- (726) Wittbecker, E. L. (to E. I. du Pont de Nemours and Co., Inc.), U. S. Patent 2,731,445; *Chem. Abstr.*, **50**, 11,054d (1956).
- (727) Wittbecker, E. L. (to E. I. du Pont de Nemours and Co., Inc.), U. S. Patent 2,731,446; *Chem. Abstr.*, **50**, 11,054e (1956).
- (728) Wittbecker, E. L., and Katz, M. (to E. I. du Pont de Nemours and Co., Inc.), U. S. Patent 2,787,630; *Chem. Abstr.*, **51**, 12,977a (1957).
- (729) Wittbecker, E. L., and Morgan, P. W., *J. Polymer Sci.*, **40**, 289 (1959).
- (730) Wittbecker, E. L., and Katz, M., *J. Polymer Sci.*, **40**, 367 (1959).
- (731) Wittbecker, E. L. (to E. I. du Pont de Nemours and Co., Inc.), U. S. Patent 2,913,433; *Chem. Abstr.*, **54**, 6200h (1960).
- (732) Wittbecker, E. L. (to E. I. du Pont de Nemours and Co., Inc.), U. S. Patent 2,970,987; *Chem. Abstr.*, **55**, 5978h (1961).
- (733) Wittbecker, E. L. (to E. I. du Pont de Nemours and Co., Inc.), U. S. Patent 3,036,979; *Chem. Abstr.*, **57**, 7484h (1962).
- (734) Wolf, G., and Seligman, A. M., *J. Am. Chem. Soc.*, **73**, 2080 (1951).
- (735) Wolff, H., *Ber.*, **22**, 2395 (1889).
- (736) Woolley, J. M. (to Imperial Chemical Industries, Ltd.), British Patent 729,505; *Chem. Abstr.*, **50**, 712e (1956).
- (737) Wright, W. B., Jr. (to American Cyanamid Co.), U. S. Patent 2,714,105; *Chem. Abstr.*, **50**, 12,120b (1956).
- (738) Wright, W. B., Jr., and Brabander, H. J., *J. Org. Chem.*, **26**, 4051 (1961).
- (739) Wright, W. B., Jr., and Brabander, H. J., *J. Org. Chem.*, **26**, 4057 (1961).
- (740) Wurtz, A., *Compt. rend.*, **68**, 1298 (1869).
- (741) Wurtz, A., et al., *Bull. soc. chim. France*, [2] **44**, 26 (1885).
- (742) Yamamoto, M., Sakuma, M., and Yamaguchi, T., *J. Pharm. Soc. Japan*, **64**, 214 (1944); *Chem. Abstr.*, **45**, 2897a (1951).
- (743) Yoder, L., *J. Am. Chem. Soc.*, **45**, 475 (1923).
- (744) Yura, S., and Ono, T., *J. Soc. Chem. Ind. Japan*, **48**, 31 (1945); *Chem. Abstr.*, **42**, 6332a (1948).
- (745) Yura, S., *J. Chem. Soc. Japan, Ind. Chem. Sect.*, **51**, 157 (1948); *Chem. Abstr.*, **45**, 547b (1951).
- (746) Zahn, H., and Dominik, M., *Makromol. Chem.*, **44/46**, 290 (1961).
- (747) Ziegenbein, W., and Franke, W., *Angew. Chem.*, **71**, 573 (1959).
- (748) Zinkeisen, W., *Ber.*, **22**, 2946 (1889).
- (749) Zinner, G., *Angew. Chem.*, **69**, 480 (1957).
- (750) Zinner, G., *Arch. Pharm.*, **292**, 329 (1959); *Chem. Abstr.*, **54**, 3196i (1960).