

ADDITIVITY RULES FOR THE ESTIMATION OF THERMOCHEMICAL PROPERTIES^{1a}

S. W. BENSON, F. R. CRUICKSHANK,^{1b} D. M. GOLDEN, G. R. HAUGEN, H. E. O'NEAL,^{1c}
A. S. RODGERS,^{1d} R. SHAW, AND R. WALSH^{1e}

Department of Thermochemistry and Chemical Kinetics, Stanford Research Institute, Menlo Park, California 94025

Received April 24, 1968

Contents

I. Introduction	279
II. General Discussion	279
III. Hydrocarbons	280
A. Alkanes	280
B. Alkenes	281
C. Alkynes	283
D. Aromatics	283
E. Polyaromatics	283
F. Cyclic Alkanes and Alkenes	285
IV. Oxygen-Containing Compounds	293
V. Nitrogen-Containing Compounds	294
A. Amines	294
B. Imines	295
C. Cyanides	295
D. Nitro Compounds	295
E. Nitrites and Nitrates	296
F. Hydrazines, Azo Compounds, and Tetrazines	296
G. Amides and Imides	296
H. Nitrogen-Containing Ring Compounds	296
I. Some Unique Group Compounds	298
J. Miscellaneous Compounds	299
VI. Halogen-Containing Compounds	301
VII. Sulfur, Boron, Phosphorus Compounds and Organometallic Compounds	304
A. Organosulfur Compounds	305
B. Organophosphorus Compounds	309
C. Organoboron Compounds	309
D. Organometallic Compounds	310
E. Miscellaneous Compounds	311
VIII. Guide to the Use of the Group Tables	313

I. Introduction

The purpose of this review is to extend, as a result of the large increase in data since 1958, the group additivity scheme for estimation of thermochemical properties as originally proposed by Benson and Buss² (BB). Because the principal interest of the authors is the application of thermochemistry to chemical kinetics, C_p° contributions have been tabulated as a func-


tion of temperature and all values predicted from the group tables refer to the gas phase. The original concept has been expanded to include cyclic compounds and, whenever possible, the experimental data are critically reviewed up to December of 1967. Sufficient data are available for hydrocarbons to render suspect any data which deviate from the value predicted by the scheme, by more than 1 cal/(mole deg) in C_p° or S° , or 1 kcal/mole in ΔH_f° . Somewhat larger errors in experimental values may therefore be detected by comparison with predicted values. From the tables of experimental data, it is clear that there are many common compounds for which the values of heat capacity and entropy are unmeasured.

In the interests of accuracy, most weight has been given to data produced after 1930, but the accuracy of thermochemical data is not uniform over all classes of compounds, and the reader is referred to the text on the appropriate compound type for a critical appraisal of the associated errors. Sources of inaccuracy in heats of formation are the inherent error in the use of heats of combustion, which are typically about ± 1 kcal/mole, the neglect by many authors of the change in latent heat of vaporization with temperature, and the necessity of estimating unmeasured latent heats. This last is smaller than the first two sources, the formula used in this review having been found to give satisfactory results. It is $\Delta H_{\text{vap}}(25^\circ) = S_T[(1.76 \times 10^{-3})t_B + 0.253]$, where $\Delta H_{\text{vap}}(25^\circ)$ is in kilocalories/mole, S_T is the Trouton constant, usually 22 cal/(mole deg); and t_B , the boiling point, is in degrees centigrade. Entropies and heat capacities are determined either calorimetrically or from vibrational assignment and statistical formulas. In the former case, errors are likely to be quite small, and in the latter case errors depend on the accuracy of the assignment, especially of the lower frequency modes, and often on the suitability of estimated barriers to internal rotation.

II. General Discussion

The reader is referred to BB² for an extensive discussion of additivity rules and the hierarchy of approximations which exist. It is sufficient for the purposes of the present review to remind the reader of the basic concept and the assumptions involved.

For the disproportionation reaction



the group additivity law assumes that $\Delta\Phi = \Delta\Phi_s$, where Φ is any molecular property and $\Delta\Phi_s$ is the contribution to that property due to symmetry changes and optical isomerism.

(1) (a) This work has been supported in part by Grant No. AP-00353-03 from the Department of Air Pollution, Public Health Service. (b) University of Strathclyde, Glasgow Ci, U. K. (c) San Diego State College, Department of Chemistry, San Diego, Calif. (d) Texas A & M University, Department of Chemistry, College Station, Texas. (e) The University, Department of Chemistry, Whiteknights Park, Reading, Berkshire, England.

(2) S. W. Benson and J. H. Buss, *J. Chem. Phys.*, **29**, 546 (1958).


Table I^a
Alkanes
(C-(C)(H)₃; C-(C)₂(H)₂; C-(C)₃(H); C-(C)₄; Alkane *gauche* Correction)

	ΔH_f°		Δ^b	σ	S°		Δ	$C_p^{300^\circ}$		Δ	$C_p^{800^\circ}$		Δ
	Obsd	Estd			Obsd	Estd		Obsd	Estd		Obsd	Estd	
Ethane	-20.24	-20.16	-0.08	2 × 3 ^a	54.85	55.09	-0.2	12.65	12.38	0.3	25.83	26.04	-0.2
Propane	-24.82	-25.11	0.29	2 × 3 ^a	64.51	64.51	0.0	17.66	17.88	-0.2	37.08	37.11	0.0
<i>n</i> -Butane	-30.15	-30.06	-0.09	2 × 3 ^a	74.12	73.93	0.2	23.40*	23.38	0.0	48.23	48.19	0.0
<i>n</i> -Hexane	-39.96	-39.96	0.00	2 × 3 ^a	92.83	92.77	0.1	34.37*	34.37	0.0	70.36	70.34	0.0
<i>n</i> -Octane	-49.82	-49.86	0.04	2 × 3 ^a	111.55	111.61	-0.1	43.35*	45.36	0.0	92.50	92.48	0.0
<i>n</i> -Decane	-59.67	-59.76	0.09	2 × 3 ^a	(130.17)	130.45	-0.3	(56.34)	56.35	0.0	(114.63)	114.63	0.0
<i>n</i> -Dodecane	-69.52	-69.66	0.14	2 × 3 ^a	(148.78)	149.29	-0.5	(67.33)	67.34	0.0	(136.76)	136.78	0.0
2-Methylpropane	-32.15	-32.14	-0.01	3 ^a	70.42	70.44	0.0	23.25*	23.11	0.1	48.49	48.37	0.1
2-Methylbutane	-36.92	-36.29	-0.63	3 ^a	82.12	82.04	0.1	28.54*	28.61	-0.1	59.71	59.45	0.3
2,2-Dimethylpropane	-39.67	-39.82	0.15	3 ^b × 2 ^a	72.23	72.88	-0.7	29.21*	29.14	0.1	60.78	60.85	-0.1
2,3-Dimethylbutane	-42.49	-42.52	0.03	3 ^a	87.42	87.39	0.0	33.76	33.84	0.1	70.7	70.70	0.0
2,2,3-Trimethylbutane	-48.96	-48.60	-0.36	3 ^a	91.61	91.70	-0.1	39.54*	39.86	-0.3	82.73	85.18	-0.5
2,2,3,3-Tetramethylbutane	-53.99	-54.68	0.69	3 ^b × 2	93.06	93.45	-0.4	46.29*	45.89	0.4	96.18	95.66	-0.5

^a Data from API compilation, except when noted. ^b Alkane group values for ΔH_f° and S° are from ref 2. ^c Data marked with an asterisk used in a regression coefficient analysis to obtain the group values.

For the molecular properties of interest here, $\Delta H_T \rightarrow 0$, $\Delta C_p^\circ \rightarrow 0$, and $\Delta S_T \rightarrow \Delta S_\sigma = R \ln K_\sigma$, where $K_\sigma = \sigma(\text{RNN}'\text{R})_\sigma(\text{SNN}'\text{S})/\sigma(\text{RNN}'\text{S})_\sigma(\text{SNN}'\text{R})$, $\sigma(x)$ being the symmetry number including both internal and external symmetry. An additional term for entropy of mixing, due to the existence of optical isomers, will be discussed below.

If the molecular framework NN' is two atoms or greater, these relationships imply the additivity of group properties, which include all nearest neighbor interactions, since a group is defined as an atom together with its ligands (e.g., in the group C-(H)₃(C) the central C atom is bonded to three H atoms and one C atom). Thus the equation



implies the additivity of the properties of the groups C-(H)₃(C), C-(H)₂(O), O-(C)(H), C-(H)₂(C)(O), and O-(C)₂, if the appropriate $\Delta\Phi = \Delta\Phi_\sigma$.

Several groups can be uniquely determined from simple molecules with well-known properties, but there are some groups which exist only in conjugate pairs, either as sums or differences. In this latter case, we have arbitrarily assigned values to one of the pairs. This was not done previously but greatly facilitates calculations without affecting the value of any property of any molecule.³

The basic approximation of group additivity makes no allowance for next-nearest neighbor interactions, but there are some molecules in which such interactions have considerable influence on the molecular properties. Such refinements are tabulated in the present scheme under the label of conformational corrections. The group values in the tables are based on the most stable conformation.

Since all carbon-carbon double bonds are between two sp² carbon atoms, the nomenclature C_d-(H)₂ is used to replace the former, C_d-(C_d)(H)₂, etc. Similar treatment is given to triple bonds, C_t-(X), aromatic ring carbon atoms, C_B-(X), and C_a, the allenic group >C=C=C<. Other group abbreviations are noted at the end of the appropriate table.

(3) This represents a change of policy; see ref 42 in ref 2.

Arbitrarily assigned group values are indicated by placing the group name in square brackets. Individual values which have been estimated are in parentheses. Values of Δ , which is defined as the difference between observed and estimated values, are quoted as [0] when the compound in question is the sole basis for a particular group. Entropies and heat capacities calculated from spectroscopically determined vibrational frequencies are here considered as measured, even if internal rotation barriers were transferred from another molecule. All original references are given where possible with the exception of values obtained from the American Petroleum Institute compilation, "Selected Values of Physical and Thermochemical Properties of Hydrocarbons and Related Compounds," by F. D. Rossini, *et al.*, Carnegie Press, Pittsburgh, Pa., 1953, which is the source of all unreferenced entries. If any entry comes from more than one source, one of which is the above, it is referenced as A.

Recognizing that their search for data, while it was exhausting, may not have been exhaustive, the authors would be grateful to have any values of thermodynamic properties of the type listed here brought to their notice. When sufficient such numbers are available, we will issue a set of revised tables.

III. Hydrocarbons

A. ALKANES (C-(C)(H)₃; C-(C)₂(H)₂; C-(C)₃(H); C-(C)₄, ALKANE *gauche*)

The values of the group contribution to heat of formation and entropy were taken directly from BB,² as was the value for the *gauche* correction. Values for the heat capacity, obtained from a multiple linear regression coefficient analysis (MLRCA), are listed in Table I.

Corrections to the heat of formation for *gauche* configurations are made one for each such conformation in a molecule. Thus, observing along a C(sp³)-C(sp³) bond, as shown in Figure 1, the *gauche* corrections are as listed for the substituted butanes.

Analysis of the simply branched alkanes will show deviations predominantly small and of the same sign, i.e., $\Delta H_{\text{obsd}} - \Delta H_{\text{estd}} \sim 0.4$ kcal/mole. This is done deliberately to avoid




Figure 1.

Table II
The C₉ Hydrocarbons^a

Compound	ΔH_f°		Δ
	Obsd	Estd	
3,3-Diethylpentane	-55.44	-53.22	-2.22
2,2,3,3-Tetramethylpentane	-56.70	-58.03	1.33
2,2,3,4-Tetramethylpentane	-56.64	-58.98	2.34
2,2,4,4-Tetramethylpentane	-57.83	-61.23	3.40
2,3,3,4-Tetramethylpentane	-56.46	-59.00	2.54

^a Values from A. Labbauf, J. B. Greenshields, and F. D. Rossini, *J. Chem. Eng. Data*, 6, 261 (1961).

excessively large errors in the heavily substituted alkanes. The largest error of +3.4 kcal/mole among the latter occurs with 2,2,4,4-tetramethylpentane. This compound has two *t*-butyl groups on the same atom and six *gauche* interactions. Data from related compounds, such as di-*t*-butyl ether, will show that *gauche* interactions do not account sufficiently for the strain which is between H atoms and C atoms separated by three C atoms, a next-next-next-nearest neighbor interaction. No simple scheme can ever hope to account for such a distant interaction, and these may be expected to occur frequently in heavily branched compounds. Thus the discrepancies shown in Table II are probably real.

B. ALKENES

1. Monoolefins (C_d-(H)₂; C_d-(C)(H);
C_d-(C)₂; C-(C_d)(C)(H)₂;
C-(C_d)(C)₂(H); C-(C_d)(C)₂;
alkene *gauche*;
cis correction)

As can be seen from Table III,⁴⁻¹³ there is a comparatively large number of values of the heats of formation of this class of compounds. Group contributions to the heat of formation were arrived at by subtracting all alkane group and *gauche* contributions and obtaining a MLRCA for the molecules indicated in the table. (The *cis* interaction was fixed at 1.00 kcal/mole, except when one of the interacting groups is a *t*-butyl, in which case the *cis* correction is 4.00 kcal/mole, or when a molecule has two *cis* interactions, such as 2,3-dimethylbut-2-ene, where the total *cis* correction is 3.00 kcal/mole.)

- (4) H. F. Bartolo and F. D. Rossini, *J. Phys. Chem.*, 64, 1686 (1960).
- (5) K. W. Egger, *J. Am. Chem. Soc.*, 89, 504 (1967).
- (6) D. M. Golden, *et al.*, *ibid.*, 86, 5416 (1964).
- (7) K. W. Egger and S. W. Benson, *ibid.*, 88, 236 (1966).
- (8) D. W. Scott, *et al.*, *ibid.*, 71, 2769 (1949).
- (9) J. P. McCullough and D. W. Scott, *ibid.*, 81, 1331 (1959).
- (10) J. D. Rockenfeller and F. D. Rossini, *J. Phys. Chem.*, 65, 267 (1961).
- (11) F. M. Fraser and E. J. Prosen, *J. Res. Natl. Bur. Std.*, 54, 143 (1955).
- (12) K. W. Egger and S. W. Benson, *J. Am. Chem. Soc.*, 87, 3311 (1965).
- (13) N. P. Bekkedahl and L. A. Wood, *J. Res. Natl. Bur. Std.*, 19, 55 (1937).




Figure 2.

The available data for entropy and heat capacity are considerably less than for the heats, but sufficient measured values exist to determine the group contributions from the MLRCA fit of the data so marked in the table.

The question of *gauche* interactions along a C(sp³)-C(sp²) bond has been taken into account, as indicated in Figure 2. In this figure (1a, 2a, and 3a) is depicted the supposed most stable conformations when the sp³ carbon has a hydrogen substituent (indicated by dashed line), as the sp³ carbon increases in substitution from one to three heavy groups (indicated by full lines). It seems clear that in cases 1a and 2a no significant interaction need arise. In case 3a, while significant interaction may be expected, it will always be taken into account by the group C-C_d(C)_s, which always implies the presence of the second sp² carbon, which is interacting with the heavy substituents on the sp³ carbon.

Figure 2 (1b, 2b, and 3b) depicts what are supposed to be the most stable conformations with heavy substituents on the central sp² carbon (1a, 2a, and 3a). It is seen that in 1b, we might expect to have one alkene *gauche* correction, while molecules of the class of 2b and 3b have two alkene *gauche* corrections.

Comparing the C-C distances of the interacting groups in the alkene and in the alkane cases, about 3.2 Å in the former and 2.8 Å in the latter, we expect a smaller effect. In fact, the effect is small enough that we have chosen to ignore it in cases of the type 1b (*i.e.*, one *gauche*) and to make a correction only in cases 2b and 3b. The correction in Table XXXIII is this total correction.

Entropy and heat capacity have been determined from a MLRCA fit of the indicated data. The *cis* correction to the entropy in entropy units is +1.2 for but-2-ene, 0 for all other 2-ene's, and -0.6 for 3-ene's.

There are no measured data to lead to entropy and heat capacity values for the groups C-(C_d)(C)₂(H) and C-(C_d)(C)₂. The group values have been estimated as shown in the footnotes of Table III. These estimates lead to significantly different values than do the incremental estimates of the API original reference. These are first encountered in 3-methylbut-1-ene and 3,3-dimethylbut-1-ene, respectively. (Entropy and heat capacity values in API, other than those tabulated here, are all estimated.)

2. Dienes (C_a; C_d-(C_d)(H);
C_d-(C_d)(C); C-(C_d)₂(H)₂)

These molecules with allenic double bonds can be fitted to group values by the introduction of the C_a group, which accounts for the properties of >C=C=C<. The end carbon atoms are treated as normal C_d's. As can be seen from Table III, this accounts quite well for both the measured data and the incremental estimates in API with a mild exception in the heat capacity values of 1,2-pentadiene.

For the nonallenic dienes, there are enough data to get a value for all group contributions to the heat of formation, but the contributions of the group C-(C_d)₂(H)₂ to the entropy

Structure	Allenic Dienes: C _a			Nonallenic Dienes: C _a -(C _b)(H); C _t -(C _d)(C); C _d -(C _a)(H) ₂			Allenic Dienes: C _a			
	Heat of Formation	Entropy	Heat Capacity	Heat of Formation	Entropy	Heat Capacity	Heat of Formation	Entropy	Heat Capacity	
Oct-1-ene	-19.82*	-20.14	0.32	-0.4	14.16*	14.1	0.1	25.42*	25.4	0.0
2,2-Dimethyl-cis-hex-3-ene	-21.77*	-22.23	0.46	0.0	19.23*	19.4	-0.2	36.01*	36.0	0.0
2,2-Dimethyl-trans-hex-3-ene	-26.16*	-26.23	0.07	-0.1	(25.3)	24.5	0.8	(47.7)	47.4	0.3
2-Methyl-3-ethylpent-1-ene	-24.40*	-24.55	0.15	0.4	(24.3)	24.6	-0.3	(46.6)	46.6	0.0
2,4,4-Trimethylpent-1-ene	-26.68*	-26.38	-0.30	0.2	(25.3)	25.5	-0.2	(47.2)	47.2	0.0
2,4,4-Trimethylpent-2-ene	-25.50*	-25.80	0.30							
Allene	45.92*	46.72	-0.80							
1,2-Butadiene	38.77*	38.97	-0.20							
1,2-Pentadiene	33.61*	34.21	-0.60							
2,3-Pentadiene	31.79*	31.12	0.57							
3-Methyl-1,2-butadiene	(31.00)	30.64	0.36							
1,3-Butadiene	26.33*	26.48	-0.15		19.11	19.1	[0]	36.84	36.8	[0]
cis-1,3-Pentadiene	19.77*	19.73	0.04	0.0	(22.7)	23.0	-0.3	(47.0)	47.2	-0.2
trans-1,3-Pentadiene	18.77*	18.73	0.04	-0.1	(24.9)	24.4	0.5	(47.7)	47.6	0.1
1,4-Pentadiene	25.41	25.41	[0]	-0.3	(25.2) ^d	23.2	2.0	(47.6) ^d	46.7	0.9
2-Methyl-1,3-butadiene (isoprene)	18.09	18.09	[0]		(25.2) ^e	24.8	0.4	(48.0) ^e	47.2	0.8

* Starred values used in regression coefficient analysis to obtain group values. ^b C_a-(C_b)(C_c)(H) = C-(C_a)(H) + [C-(C_b)(C)(H)₂-C-(C_a)(H)] + [C-(C_b)(C)(H)₂-C-(C_a)(H)] - C-(C_a)(H). ^c C_a-(C_b)(C)(H) = C-(C_a)(H) + [C-(C_b)(C)(H)₂-C-(C_a)(H)] - C-(C_a)(H). ^d The value in API has been raised by R ln 2 to take into account the effect of optical isomers which was neglected in the original reference: J. Kilpatrick, *et al.*, *J. Res. Natl. Bur. Std.*, **42**, 225 (1949). ^e Group values preferred, as quoted values differ from other allenic molecules by relatively large amounts.

and heat capacity, and that of the group C_d-(C_a)(C) to heat capacity, had to be estimated as indicated in the footnotes of Table III. These estimates lead to heat capacity values which are somewhat at variance with the estimates in the API tables.

C. ALKYNES (C_t-(H); C_t-(C); C-(C_t)(C)(H)₂; C-(C_t)(C)₂(H); C_t-(C_d); C_d-(C_t)(H))

The contributions of the first four groups mentioned above to heat of formation can be obtained from the data in the API tables in the usual manner. The entropy and heat capacity contributions of all but C-(C_t)(C)₂(H) can also be ascertained from measured values, the contributions of this group being determined as noted in Table IV.¹⁴

From some observed values of heats of formation of some mixed "yne-ene" compounds, based on solution heats of hydrogenation, one can ascertain the heat of formation contribution of the sum of groups C_t-(C_d) + C_d-(C_t)(H). We have arbitrarily assigned to the C_d-(C_t)(H) group the same values as the C_d-(C_a)(H) group, and thus obtained values for the C_t-(C_d) group.

D. AROMATICS (TABLE V)

1. Alkylbenzenes (C_B-(H); C_B-(C); C-(C_B)(C)(H)₂; C-(C_B)(C)₂(H); C-(C_B)(C)₃; *ortho* correction)

Values were obtained in a straightforward manner from those values so marked. Where group values were estimated, the nature of the estimation is noted.

2. Unsaturated Benzenes (C_B-(C_d); C_d-(C_B)(H); C_d-(C_B)(C); C_B-(C_t); C_t-(C_B))

There seems to be very little in the way of measured values for this class of compounds. The contributions of the sum of the C_B-(C_d) and C_d-(C_B)(H) groups can be obtained from styrene itself, and this value allows the estimation of the properties of the other vinylbenzenes listed, with the exception of α -methylstyrene. For convenience, the C_d-(C_B)(H) group is arbitrarily assigned the values of the C_d-(C_a)(H) group. The measured value of the heat of formation of α -methylstyrene yields values for the C_d-(C_B)(C) group contribution. The entropy and heat capacity contributions of this group are estimated as equal to those of the C_d-(C_a)(C) group.

The values of entropy and heat capacity contributions of the sum [C_B-(C_t) + C_t-(C_B)] are estimated from phenylacetylene, and the C_B-(C_t) group is assigned the values of the C_B-(C_d) group. The heat of formation is estimated as shown in Table V.¹⁵⁻¹⁹

E. POLYAROMATICS (TABLE VI²⁰⁻²²)

The measured values for biphenyl yield the contributions of the group C_B-(C_B) in a straightforward manner. This group seems to reproduce the heats of naphthalenic compounds

- (14) H. A. Skinner and A. Snelson, *Trans. Faraday Soc.*, **55**, 404 (1959).
- (15) E. J. Prosen, *et al.*, *J. Res. Natl. Bur. Std.*, **36**, 455 (1946). This paper is the source of all the API values, as well.
- (16) S. H. Hastings and D. E. Nicholson, *J. Phys. Chem.*, **61**, 730 (1957).
- (17) D. W. Scott, *et al.*, *ibid.*, **66**, 911 (1962).
- (18) D. J. Coleman and G. Pilcher, *Trans. Faraday Soc.*, **62**, 821 (1966).
- (19) J. C. Evans and R. A. Nyquist, *Spectrochim. Acta*, **16**, 918 (1960).
- (20) D. M. Spero and F. D. Rossini, *J. Phys. Chem.*, **64**, 1723 (1960).
- (21) J. A. Katon and E. R. Lippincott, *Spectrochim. Acta*, **15**, 627 (1959).
- (22) C. J. Eagon, *J. Chem. Eng. Data*, **8**, 532 (1963).

Table IV^a
Alkynes
(C₁-(H); C₁-(C); C-(C)₂(H)₂; C-(C)₂(C)₂(H); C-(C)₂(C)₂(H); C₁-(C₂); C₂-(C₁)(H))

	ΔH_f°		Δ	Ref	σ	S°		Δ	Ref	C_p^{300}		Δ	Ref
	Obsd	Estd				Obsd	Estd			Obsd	Estd		
Ethyne	54.19*	53.86	0.33		2	48.00*	48.0	0.0		10.53	10.54	0.0	
Propyne	44.32*	44.40	-0.08		3	59.30*	59.3	0.0		14.55	14.6	0.0	
But-1-yne	39.48*	39.67	-0.19		3	69.51*	69.6	-0.1		19.54	19.5	0.0	
But-2-yne	34.97*	34.94	0.03		3 ² X 2	67.71*	67.8	-0.1		18.70	18.6	0.0	
Pent-1-yne	34.50*	34.72	-0.22		3	(79.10)	79.0	0.1		(25.65)	25.0	0.7	
Pent-2-yne	30.80*	30.21	0.59		3 ²	(79.30)	79.5	-0.2		(23.69)	23.6	0.1	
3-Methylbut-1-yne	32.60*	32.60	[0]		3 ²	(76.23) ^b	76.2	0.0		(25.1) ^b	24.8	0.3	
Hexa-1,5-diyne (dipropargyl)	99.44*	99.50	-0.06	14 ^c									
cis-Pent-3-en-1-yne	60.60*	61.64	-1.04	14 ^c									
trans-Pent-3-en-1-yne	60.92*	60.64	0.28	14 ^c									
cis-Dec-3-en-1-yne	36.25	36.81	-0.56	14 ^c									
trans-Dec-3-en-1-yne	36.95*	35.91	1.04	14 ^c									

^a Starred values used in regression coefficient analysis to obtain group values. ^b C-(C₁)(C)₂(H) = C-(C)₂(H) + [C-(C)₂(C)(H)₂-C-(C)₂(H)]₂. ^c These are calculated from heats of hydrogenation studies in solution which are often low by 1 or 2 kcal/mole.

Table V

	ΔH_f°		Δ	Ref	σ	S°		Δ	Ref	C_p^{300}		Δ	Ref
	Obsd	Estd				Obsd	Estd			Obsd	Estd		
Benzene	19.82	19.79	0.03		3 X 2 ²	64.34*	64.26	0.1		19.65*	19.42	0.2	
Toluene	11.95*	11.92	0.03		3 X 2	76.64*	76.81	-0.2	17	24.94*	25.04	-0.1	
Ethylbenzene	7.12*	7.06	0.06		3 X 2	86.15*	86.15	[0]		30.88*	30.88	[0]	
Isobutylbenzene	-5.15	-4.92	-0.23	15	3 ² X 2	...	102.33	41.61	...	
1,2-Dimethylbenzene	4.54*	4.63	-0.09		3 ² X 2	84.34*	84.21	0.1	16	32.10*	31.79	0.3	
1,3-Dimethylbenzene	4.12*	4.06	0.06		3 ² X 2	85.49*	85.82	-0.3		30.66*	30.66	0.0	
1,4-Dimethylbenzene	4.29*	4.06	0.23		3 ² X 2	84.23*	84.45	-0.2		30.49*	30.66	-0.2	
n-Propylbenzene	1.87*	2.11	-0.24		3 X 2	(95.74)	95.59	0.2		(36.99)	36.48	0.5	
n-Butylbenzene	-3.30*	-2.84	-0.46		3 X 2	(105.04)	105.01	0.0		(42.09)	41.98	0.1	
1-Propylbenzene	0.94*	0.86	0.08		3 ² X 2	(92.87) ^a	92.91	0.0		(36.47) ^a	36.11	0.4	
sec-Butylbenzene	-4.17*	-4.09	-0.08	15	3 ² X 2	...	102.33	41.61	...	
t-Butylbenzene	-5.42*	-5.42	[0]	15	3 ¹ X 2	...	95.93	42.13	...	
1-Methyl-2-ethylbenzene	0.29*	-0.23	0.52		3 ²	(95.42)	94.93	0.5		(37.94)	37.63	0.3	

Alkylbenzenes: (C_B-(H); C_B-(C); C-(C_B)(C)(H)₂; C-(C_B)(C)₂(H); C-(C_B)(C)₂(H); C-(C_B)(C)₂; ortho Correction)

1-Methyl-3-ethylbenzene	-0.46	-0.80	0.34	3 ²	(96.60)	96.54	0.1	(36.59)	36.51	0.1	(77.8)	77.59	0.2
1-Methyl-4-ethylbenzene	-0.78	-0.80	0.02	3 ² × 2	(95.34)	95.18	0.2	(36.42)	36.51	-0.1	(77.6)	77.59	0.0
1,2,3-Trimethylbenzene	-2.29*	-2.66	0.37	3 ³ × 2	91.79*	91.60	0.2	16	37.82*	-0.7	77.56*	78.30	-0.7
1,2,4-Trimethylbenzene	-3.33*	-3.24	-0.09	3 ³	94.54*	94.58	0.0	16	36.99*	-0.4	76.93*	77.41	-0.5
1,3,5-Trimethylbenzene	-3.84*	-3.81	-0.03	3 ⁴ × 2	92.15*	92.65	-0.5		36.10*	-0.2	76.84*	76.51	0.3
1,2,3,4-Tetramethylbenzene	-10.02*	-9.96	-0.06	3 ⁴ × 2	99.55*	99.00	0.6	16	45.50*	0.2	89.42*	89.62	-0.2
1,2,3,5-Tetramethylbenzene	-10.71*	-10.54	-0.17	3 ⁴ × 2	100.99*	100.62	0.4	16	44.57*	0.4	88.79*	88.74	0.1
1,2,4,5-Tetramethylbenzene	-10.82*	-10.53	-0.29	3 ⁴ × 2 ²	100.03*	99.26	0.8	16	44.77*	0.6	88.41*	88.74	-0.3
Pentamethylbenzene	(-17.79)	-17.27	-0.52	3 ⁵ × 2	106.09*	106.40	-0.3	16	51.99*	0.0	101.29*	100.94	0.4
Hexamethylbenzene	(-25.26)	-24.00	-1.26	3 ⁷ × 2 ²	108.12*	108.65	-0.5	16	59.73*	-0.2	113.51*	113.14	0.4
Bibenzyl	32.41	34.30	-1.9	2 ²	...	129.90	44.08	85.98	...
Styrene	35.22	35.22	[0]	2	82.48	82.48	[0]		29.35	[0]	61.40	61.40	[0]
S-Methylstyrene	27.00	27.00	[0]	3 × 2	(91.7) ^c	89.73	2.0		35.48	-0.6	(71.8) ^c	72.57	-0.8
cis-β-Methylstyrene	(29.00)	28.77	0.23	3 × 2	(91.7)	91.07	0.6		33.26	0.6	(71.8)	71.61	0.2
trans-β-Methylstyrene	(28.00)	27.77	0.23	3 × 2	(90.9)	91.07	0.8		34.60	0.5	(72.2)	72.00	0.2
o-Methylstyrene	(28.30)	27.92	0.38	3	(91.7)	91.24	0.5		36.09	-1.2	(71.8)	71.82	0.0
m-Methylstyrene	(27.60)	27.35	0.25	3	(93.1)	92.85	0.3		34.97	-0.1	(71.8)	70.94	0.9
p-Methylstyrene	(27.40)	27.35	0.05	3 × 2	(91.7)	91.49	0.2		34.97	-0.1	(71.8)	70.94	0.9
Ethynylbenzene	...	(78.31)	...	2	76.88	76.88	[0]	19	27.63	[0]	55.79	55.79	[0]

^a C-(C_B)(C_B)(H) = C-(C_B)(H) + [C-(C_B)(C)₂] - C-(C_B)(C)(H)₂ - C-(C_B)(C)(H)₂ + [C-(C_B)(C)(H)]₂ - C-(C_B)(C)(H)₂ - C-(C_B)(C)(H)₂ = C₁-(C_B)(C) = C₁-(C_B)(B). ^d C₁-(C_B)(B) = C₁-(C_B)(C) = C₁-(C_B)(C_B) = C₁-(C_B)(C_B).

quite well, but begins to fall off with larger compounds of this class. While the value for anthracene is not bad, it is clear from the value of phenanthrene that this type of pseudo group additivity does not take into account some important interactions. Note that biphenyl is not planar in the gas phase so that it may be lacking in some conjugation energy. The value shown for $\Delta H_f^\circ(\text{C}_B-\text{C}_B)$ was an average over biphenyl and naphthalenes.

Entropy and heat capacity can be treated with a ring correction, as for the cycloalkanes and alkenes treated in the next section.

F. CYCLIC ALKANES AND ALKENES

Our approach to cyclic compounds has been to use groups from the open-chain compounds and a ring correction. In the case of heats of formation, this correction is what is often defined as the strain energy. Since the available data are sparse, all ring strains were evaluated from the unsubstituted rings where data for these were available. (For the cyclohexanes, enough data are available to allow us to relax this rule.) The transferability of this strain to substituted ring compounds is generally attested to by the data of Table VII.²³⁻⁵¹ It will be noticed that the deviations are somewhat larger than previously encountered. This is due to the absence of gas-phase data, the use of the unsubstituted ring as standard, and perhaps small differences in the strain energies of various substituted small rings.

The entropy and heat capacity correction terms are also seen to be transferable. Again, an examination of the deviations will show that choice of the unsubstituted ring as the standard often throws the values of the substituted compounds off by a small amount.

The magnitude of the entropy corrections is not surprising when one realizes that a disproportionate amount of the entropy is tied up in the C-(H)₃(C) and C_d-(H)₂ groups of open-chain molecules.

- (23) J. W. Knowlton and F. D. Rossini, *J. Res. Natl. Bur. Std.*, **43**, 113 (1949).
- (24) K. B. Wiberg, *et al.*, *J. Am. Chem. Soc.*, **84**, 3980 (1962).
- (25) S. Kaarsemaker and J. Coops, *Rec. Trav. Chim.*, **71**, 261 (1952).
- (26) R. B. Turner, *et al.*, *Tetrahedron Letters*, **15**, 997 (1965).
- (27) R. B. Turner and R. H. Garner, *J. Am. Chem. Soc.*, **80**, 1424 (1958).
- (28) R. B. Turner, Kekule Symposium, IUPAC, Butterworth & Co., Ltd., London, 1959.
- (29) M. A. Dolliver, *et al.*, *J. Am. Chem. Soc.*, **59**, 831 (1937).
- (30) J. E. Bloor and S. Gartside, *Nature*, **184**, 1313 (1959).
- (31) G. B. Kistiakowsky, *et al.*, *J. Am. Chem. Soc.*, **58**, 137 (1936).
- (32) G. B. Kistiakowsky, *et al.*, *ibid.*, **58**, 146 (1936).
- (33) R. B. Turner, private communication (ΔH_f° , at 25°, acetic acid, and is 0.3 kcal more negative than for 1,3-cyclohexadiene).
- (34) K. A. Kobe and R. E. Pennington, *Petrol. Refiner.*, **29**, 93 (1950).
- (35) K. B. Wiberg and W. J. Batley, *J. Am. Chem. Soc.*, **82**, 6375 (1960).
- (36) G. W. Rathyens, Jr., *et al.*, *ibid.*, **75**, 5634 (1953).
- (37) A. Danti, *J. Chem. Phys.*, **27**, 1227 (1957).
- (38) D. W. Scott, *et al.*, *J. Phys. Chem.*, **64**, 906 (1960).
- (39) J. B. Conn, *et al.*, *J. Am. Chem. Soc.*, **61**, 1868 (1939).
- (40) J. B. Conn, *et al.*, *ibid.*, **60**, 440 (1938).
- (41) R. B. Turner and W. R. Meeder, *ibid.*, **79**, 4133 (1957).
- (42) R. B. Turner, *et al.*, *ibid.*, **79**, 4127 (1957).
- (43) D. M. Speros and F. D. Rossini, *J. Phys. Chem.*, **64**, 1723 (1960).
- (44) H. Leroux, *Compt. Rend.*, **151**, 384 (1910).
- (45) C. C. Browne and F. D. Rossini, *J. Phys. Chem.*, **64**, 927 (1960).
- (46) J. M. Fraser and F. J. Prosen, *J. Res. Natl. Bur. Std.*, **54**, 143 (1955).
- (47) H. L. Finke, *et al.*, *J. Am. Chem. Soc.*, **78**, 5489 (1956).
- (48) D. W. Scott, *et al.*, *ibid.*, **72**, 4664 (1950).
- (49) R. Srinivasan, *et al.*, *J. Phys. Chem.*, **69**, 1775 (1965).
- (50) R. H. Boyd, *et al.*, Abstracts, 152nd National Meeting of the American Chemical Society, New York, N. Y., Sept 1966, No. V164.
- (51) R. B. Williams, *J. Am. Chem. Soc.*, **64**, 1395 (1942).

Table VI

	ΔH_f°		Δ	Ref	σ	S°		Δ	Ref	C_p^{300}		Δ	C_p^{800}		Δ	Ref
	Obsd	Estd				Obsd	Estd			Obsd	Estd		Obsd	Estd		
Biphenyl	43.52	42.92	0.60	18	2 ^s	93.91	93.91	[0]	21	39.05	39.05	[0]	86.92	86.92	[0]	21
Naphthalene	36.12	36.32	-0.20	18, 20	2 ^s	80.5 ^a	80.5	[0]	22							
1-Methylnaphthalene	27.93	28.43	-0.50	20												
2-Methylnaphthalene	27.75	28.43	-0.6	20												
Anthracene	55.17	52.84	2.67	18												
Phenanthrene	49.46	52.84	-3.38	18												

^a Naphthalenic ring correction = 8.1 gibbs/mole.

Table VII

Nonaromatic Rings: Ring Corrections

	ΔH_f°		Δ	Ref	n	σ	S°		Δ	Ref	C_p^{300}		Δ	C_p^{800}		Δ	Ref
	Obsd	Estd					Obsd	Estd			Obsd	Estd		Obsd	Estd		
Cyclopropane	12.74	12.7	[0]	23	3	2	56.8	56.8	[0]	34	13.44	13.44	[0]	31.44	31.44	[0]	34
Methylenecyclopropane	48.0	48.0	[0]	^e													
Cyclopentane	66.6	66.6	[0]	24	2		58.38	58.38	[0]	35							
1-Methylcyclopentane	58.2	58.3	-0.1	^e													
Cyclobutane	16.4	6.4	[0]	25	2 ^s		63.43	63.43	[0]	36	17.37	17.37	[0]	42.42	42.42	[0]	36
Cyclobutene	37.5	37.5	0.0	^e	2		62.98	62.98	[0]	37	16.03	16.03	[0]	32.26	32.26	[0]	37
Methylcyclobutane	...	-0.6													
1,2-Dimethylcyclobutene	20.8	21.7	-0.9	^f													
1-Methylcyclobutene	29.1	29.1	[0]	27 ^{a,b}													
1-Methyl-1,3-methylenecyclobutene	48.3	48.6	-0.3	28 ^{a,b}													
Methylenecyclobutane	30.0	28.0	2.0	27 ^{a-c}													
1,3-Dimethylenecyclobutane	53.3	51.4	1.9	28 ^{a-c}													
Cyclopentane	-18.46	-18.46	[0]		10		70.0	70.0	[0]		19.98	19.98	[0]	52.44	52.44	[0]	
Cyclopentene	8.56	8.56	[0]	29	2		69.23	69.23	[0]		18.08	18.08	[0]	45.78	45.78	[0]	
Cyclopentadiene	32.44	32.44	[0]	30													
Methylcyclopentane	-25.50	-25.49	-0.0		3		81.22	81.34	-0.1	38	26.43	25.21	1.2	63.8	63.7	-0.1	38
1-Methylcyclopentene	-1.5	-0.29	-1.2	27 ^a	3		(78.2)	78.1	0.1		(24.3)	24.2	-0.1	(57.0)	57.1	-0.1	
1-Ethylcyclopentene	-5.8	-5.05	-0.8	27 ^a													
Methylenecyclopentane	2.4	3.32	-0.9	27 ^{a-c}													
Ethylidenecyclopentane	-4.5	-4.13	-0.4	27 ^{a-c}													
1,1-Dimethylcyclopentane	-33.05	-33.17	0.1		3 ^s × 2		(85.87)	85.18	0.69		(32.16)	31.23	0.93	(76.18)	76.20	0.0	
cis-1,2-Dimethylcyclopentane	-30.96	-31.51	0.6		3 ^s		(87.51)	88.14	-0.63		(32.34)	30.44	1.9	74.98	74.94	1.0	
trans-1,2-Dimethylcyclopentane	-32.67	-32.52	-0.2		2		(87.67)	88.14	-0.47		(32.44)	30.44	2.0	(75.84)	74.94	0.9	
cis-1,3-Dimethylcyclopentane	-32.47	-32.52	0.6		3 ^s		87.54	88.14	-0.6	38	32.72	30.44	2.3	75.68	74.94	0.8	38

<i>trans</i> -1,3-Dimethylcyclopentane	-31.93	-32.52	0.1	2	3 ² × 2	(87.54)	88.14	-0.6	(32.72)	30.44	2.3	(75.68)	74.94	0.8
Cyclohexane	-29.43	-29.73	0.3	3	3 × 2	71.28	71.58	-0.3	25.58	26.58	-1.0	66.76	67.26	-0.5
Cyclohexene	-0.84	-0.84	[0]	31	2	74.27	74.27	[0]	25.28	25.28	[0]	59.49	59.49	[0]
1,3-Cyclohexadiene	26.0	26.0	[0]	32	3				32.51	31.81	0.7	78.74	78.5	0.2
1,4-Cyclohexadiene	26.3	26.3	[0]	33	3				38.23	37.31	0.9	90.1	89.6	0.5
Methylcyclohexane	-36.99	-36.76	-0.2		3 ²	87.24	87.12	0.1	(37.2)	37.0	0.2	(90.7)	91.0	-0.3
Ethylcyclohexane	-41.05	-40.91	-0.1		3 ²	89.51	89.96	-0.5	(37.7)	37.0	0.7	(90.1)	89.8	0.3
1,1-Dimethylcyclohexane	-43.26	-42.81	-0.5		3 ² × 2	88.65	88.60	0.1	(38.3)	37.0	1.3	(90.5)	89.8	0.7
<i>cis</i> -1,2-Dimethylcyclohexane	-41.15	-41.39	0.2	2	3 ²	88.54	88.60	-0.1	(37.9)	37.0	1.9	(90.5)	89.8	0.7
<i>trans</i> -1,2-Dimethylcyclohexane	-43.02	-42.99	0.0	2	3 ²	89.92	89.96	0.0	(37.9)	37.0	0.9	(89.8)	89.8	0.0
<i>cis</i> -1,3-Dimethylcyclohexane	-44.16	-43.79	-0.4		3 ² × 2	88.54	88.60	-0.1	(37.9)	37.0	0.9	(89.8)	89.8	0.0
<i>trans</i> -1,3-Dimethylcyclohexane	-42.20	-42.19	0.0	2	3 ²	88.54	88.60	-0.1	(37.9)	37.0	0.9	(89.8)	89.8	0.0
<i>cis</i> -1,4-Dimethylcyclohexane	-42.22	-42.19	0.0		3 ² × 2	87.19	87.24	-0.1	(38.0)	37.0	1.0	(90.6)	89.8	0.8
<i>trans</i> -1,4-Dimethylcyclohexane	-44.12	-43.79	-0.3											
1-Methylcyclohexene	-10.0	-9.7	-0.3	27 ^a										
1-Ethylcyclohexene	-15.0	-14.5	-0.5	27 ^a										
Methylenecyclohexane	-7.2	-8.1	0.9	27 ^{a,e}										
Ethylenecyclohexane	-13.7	-15.5	1.8	27 ^{a,e}										
Cycloheptane	-28.3	-28.3	[0]	25	1	81.8	81.8	[0]	47					
Cycloheptene	-1.8	-1.8	[0]	39										
1,3-Cycloheptadiene	22.9	22.9	[0]	39										
1,3,5-Cycloheptatriene	44.5	44.5	[0]	40	1	75.4	75.4	[0]	47					
1-Methylcycloheptene	-10.5	-10.7	0.2	27 ^{a,b}										
Methylenecycloheptane	-8.2	-6.7	-1.5	27 ^{a-c}										
Cyclooctane	-29.7	-29.7	[0]	25	2 ³	87.7	87.7	[0]	47					
<i>cis</i> -Cyclooctene	-6.2	-6.2	[0]	39										
<i>trans</i> -Cyclooctene	3.1	3.1	[0]	41										
1,3,5-Cyclooctatriene	43.7	43.7	[0]	42 ^a										
Cyclooctatetraene	71.3	71.3	[0]	42 ^a										
Cyclononane	-31.8	-31.8	[0]	25										
<i>cis</i> -Cyclononene	-7.2	-7.2	[0]	41 ^a										
<i>trans</i> -Cyclononene	-4.3	-4.3	[0]	41 ^a										
<i>trans</i> -Decalin	-43.6	-42.8	-0.8	43										
<i>cis</i> -Decalin	-40.4	-41.2	0.8	43										
Tetralin	2.6	6.6	-4.0	44 ^a										
Hydrindene	14.3	14.8	-0.5	29										
<i>trans</i> -Hydrindane	-31.5	-31.4	-0.1	45										
Spiropentane	44.23	44.23	[0]	46	2 ²	67.46	67.46	[0]	48	21.19	0	47.91	47.91	[0]
Bicyclo[1.1.0]butane	51.9	53.3	-1.4	<i>e</i>	2	62.49	52.49	[0]	69					
1,3-Dimethylcyclo[1.1.0]butane	39.3	37.9	-1.4	26										
Bicyclo[2.1.0]pentane	37.6	37.6	[0]	28 ^s										
Bicyclo[3.1.0]hexane	9.07	9.07	[0]	50										
Bicyclo[4.1.0]hexane	0.32	0.32	[0]	49										
Bicyclo[5.1.0]octane	-3.87	-3.87	[0]	50										
Bicyclo[6.1.0]nonane	-7.42	-7.42	[0]	50										
1,4-Dihydronaphthalene	33.3	33.3	[0]	51										
1,2-Dihydronaphthalene	30.3	30.6	0.3	51										

^a 1.0 kcal/mole added to heats derived from measurements in solution. ^b Depends upon the group value of the corresponding cycloalkane. ^c Using the ring correction of the corresponding cycloalkane. ^d Work done in 1910. ^e Added in proof: K. B. Wiberg and R. A. Fenoglio, *J. Am. Chem. Soc.*, **90**, 3395 (1968). ^f Added in proof: R. B. Turner, *et al.*, *ibid.*, **90**, 4315 (1968).





Methyl <i>n</i> -propyl ether, CH ₃ OC ₃ H ₇	-56.8	-56.7	-0.1	73
Methyl propyl ether, CH ₃ OCH(CH ₂) ₂	-60.2	-60.0 ^b	-0.2	73
Di- <i>n</i> -propyl ether, C ₃ H ₇ OC ₃ H ₇	-70.0	-70.1	+0.1	76
Diisopropyl ether, (CH ₃) ₂ CHOCH(CH ₂) ₂	-76.1	-76.8 ^c	+0.7	76
Di- <i>n</i> -butyl ether, C ₄ H ₉ OC ₄ H ₉	-79.8	-80.0	+0.2	76
Di- <i>sec</i> -butyl ether, (CH ₃ CH ₂ CH(CH ₃)) ₂ O	-86.2	-85.7 ^d	-0.5	76
Methyl <i>t</i> -butyl ether, CH ₃ OC(CH ₃) ₃	-70.0	-69.3	-0.7	77
Isopropyl <i>t</i> -butyl ether, (CH ₃) ₂ CHOCH(CH ₂) ₂	-85.6	-86.2 ^e	+0.6	77
Di- <i>t</i> -butyl ether, * (CH ₃) ₂ COC(CH ₃) ₂	-87.2	-87.2 ^f	[0.0]	77
Methyl phenyl ether, CH ₃ OC ₆ H ₅	-18.0 ^g	-18.0	0.0	78
Ethyl phenyl ether, C ₂ H ₅ OC ₆ H ₅	-25.4 ^h	-25.5	+0.1	78
Diphenyl ether, * C ₆ H ₅ OC ₆ H ₅	-10.1 ⁱ	-10.1	[0.0]	79
Divinyl ether, * CH ₂ CHOCHCH ₂	-3.0	-3.0	[0.0]	75
Ethyl vinyl ether, C ₂ H ₅ OCHCH ₂	-33.5	-33.3	-0.2	75
2-Ethoxyprop-1-ene, C ₂ H ₅ OC(CH ₃)CH ₂	-43.4 ^j	-43.4	0.0	61
2-Methoxy- <i>n</i> -but-2-ene, CH ₃ OC(CH ₃)CHCH ₃	-39.9 ^k	-40.0	0.1	61
2,2-Dimethoxypropane, (CH ₃) ₂ C(OCH ₂) ₂	-102.9	-103.3 ^l	+0.4	80
2,2-Diethoxypropane, (CH ₃) ₂ C(OC ₂ H ₅) ₂	-120.6	-102.3 ^l	-0.3	80

^a An asterisk denotes unique group. ^b One oxygen *gauche* correction of 0.3. ^c Two oxygen *gauche* corrections of 0.3. ^d Two oxygen plus two-alkyl *gauche* corrections. ^e Three oxygen *gauche* corrections. ^f Four oxygen *gauche* corrections plus strain. ^g Using calculated $\Delta H_{\text{vap}} = 11.6$. ^h Using calculated $\Delta H_{\text{vap}} = 12.3$. ⁱ From a heat of combustion of the solid plus heat of fusion of 4 and heat of vaporization of 15.2 (ref 79). ^j From a calculated heat of formation of -68.5 for ethyl isopropyl ether. ^k From a calculated heat of formation of -64.7 for methyl *sec*-butyl ether and includes one *cis* correction. ^l Includes four oxygen *gauche*.

Table X
Peroxides and Hydroperoxides

Compound	ΔH_f°		Δ	Ref	n, σ	S°		Δ	C_p^{300}		Δ	Ref				
	Obsd	Estd				Obsd	Estd		Obsd	Estd						
Dimethyl peroxide, CH ₃ OOCH ₃	-30.0	-29.2	-0.8	81												
Diethyl peroxide, C ₂ H ₅ OOCC ₂ H ₅	-46.1	-46.2	+0.1	81												
Di- <i>t</i> -butyl peroxide, (CH ₃) ₂ COOC(CH ₃) ₂	-81.5	-81.6 ^a	+0.1	81												
Hydrogen peroxide, HOOH	-32.5	-32.5	[0.0]	82	2, 2	55.7	55.7	[0.0]	82	10.3	10.2	0.1	14.3	14.4	-0.1	82
<i>t</i> -Butyl hydroperoxide, (CH ₃) ₂ COOH	-52.1	-57.1 ^b	+5.0	83												
Hexyl-1 hydroperoxide	-58.6	-58.4	-0.2	84, 85												
Hexyl-2 hydroperoxide	-61.1	-62.9	+1.8	84, 85												
Hexyl-3 hydroperoxide	-59.9	-62.9	+3.0	84, 85												
Heptyl-1 hydroperoxide	-66.0	-63.4	-2.6	84, 85												
Heptyl-2 hydroperoxide	-66.8	-67.9	+1.1	84, 85												
Heptyl-3 hydroperoxide	-66.9	-67.9	+1.0	84, 85												
Heptyl-4 hydroperoxide	-63.8	-67.9	+4.1	84, 85												
Cyclohexyl hydroperoxide	-52.3	-52.8	+0.5	84, 85												
Methyl cyclohexyl hydroperoxide	-63.0	-59.8	-3.2	84, 85												
Acetyl peroxide, (CH ₃ COO) ₂	-119.0	-129.0	+10.0	87												
Propionyl peroxide, (C ₂ H ₅ COO) ₂	-138.0	-139.0	+1.0	87												
Butyryl peroxide, (C ₄ H ₇ COO) ₂	-150.0	-149.0	-1.0	87												

^a Including four oxygen *gauche* corrections. ^b Including two oxygen *gauche* corrections. ^c Gives unique group.

Cyclohexanone*		-53.1	-53.1	[0.0]	96, 67
Succinic anhydride*		-122.6	-122.6	[0.0]	97
Glutaric anhydride*		-131.3	-131.3	[0.0]	97
Maleic anhydride*		-97.9	-97.9	[0.0]	98

* An asterisk denotes unique group. ^b Including one hydrocarbon *gauche* correction. ^c Including two oxygen *gauche* corrections. ^d Including one oxygen and one hydrocarbon *gauche* corrections.

Table XII
Aldehydes and Ketones

Compound ^a	Ref	σ	S°		Δ	ΔH_f°		Δ	Ref	$C_p^{\circ 300}$		Δ	$C_p^{\circ 800}$		Δ	Ref
			Obsd	Estd		Obsd	Estd			Obsd	Estd		Obsd	Estd		
Acetaldehyde, CH ₃ CHO	99	3	63.2	63.2	[0.0]	-39.7	-39.7	[0.0]	99	13.2	13.2	[0.0]	24.2	24.2	[0.0]	99, 100
Propionaldehyde, C ₂ H ₅ CHO	57, 101	3	73.4	72.7	-0.9	-44.7	-44.7	-0.9	57, 101	18.8	19.4	-0.6	35.2	35.3	-0.1	101
Butyraldehyde, C ₃ H ₇ CHO	57	3	82.6	82.1	+0.8	-48.9	-49.7	+0.8	57							
Heptaldehyde, C ₆ H ₁₃ CHO	57				+1.9	-62.8	-64.7	+1.9	57							
Acetone, * CH ₃ COCH ₃	102	2 × 3 ²	70.5	70.5	[0.0]	-51.7	-51.7	[0.0]	102	18.0	18.0	[0.0]	34.9	34.9	[0.0]	102
Methyl ethyl ketone, CH ₃ CH ₂ COCH ₃	103	3 ²	80.8	81.5	-0.3	-57.0	-56.7	-0.3	103	24.7	24.2	0.5	46.1	46.0	0.1	103
Diethyl ketone	105	2 × 3 ²	88.6	89.3	-0.7	-61.65	-61.56	-0.9	105	<i>e</i>						
Methyl <i>n</i> -propyl ketone	105	3 ²	90.5	90.8	-0.3	-61.92	-61.56	-0.36	105	<i>e</i>						
Ethyl <i>n</i> -propyl ketone	105				0	-66.51	-66.51	0	105							
Methyl <i>n</i> -butyl ketone	105				-0.45	-66.96	-66.51	-0.45	105							
Di- <i>n</i> -butyl ketone	105				-1.21	-82.57	-81.36	-1.21	105							
Methyl isopropyl ketone	105				+0.31	-62.76	-63.07	+0.31	105							
Ethyl isopropyl ketone	105				-0.36	-68.38	-68.02	-0.36	105							
Diisopropyl ketone	105				+0.04	-74.40	-74.44	+0.04	105							
Ethyl <i>t</i> -butyl ketone	105				+0.31	-75.00	-74.69	+0.31	105							
Isopropyl <i>t</i> -butyl ketone	105				+0.31	-80.84	-81.15	+0.31	105							
<i>t</i> -Butyl neopentyl ketone	105				-0.83	-94.15	-93.32	-0.83	105							
Di- <i>t</i> -butyl ketone	105				+5.18	-82.65	-87.83	+5.18	105							
Crotonaldehyde, * CH ₃ CH=CHCHO	61				[0.0]	-25.5	-25.5	[0.0]	61							
Furfuraldehyde*	104				0.0	-35.4	-34.4 ^b	0.0	104							
Benzaldehyde, * C ₆ H ₅ CHO	52				[0.0]	-6.0	-6.0	[0.0]	52							
Acetophenone, * C ₆ H ₅ COCH ₃	85				[0.0]	-22.0	-22.0	[0.0]	85							
Benzophenone, * C ₆ H ₅ COC ₆ H ₅	85				[0.0]	12.3	12.3	[0.0]	85							
Acetylacetone, * CH ₃ COCH ₂ COCH ₃	106				[0.0]	-90.4	-90.4 ^c	[0.0]	106							
Diacetyl, CH ₃ COCOCCH ₃	62				[0.0]	-78.6	-78.6 ^d	[0.0]	62							

^a An asterisk denotes unique group. ^b Heat of vaporization estimated to be 12 kcal/mole. ^c Heat of vaporization estimated to be 11 kcal/mole. ^d Heat of vaporization estimated to be 9 kcal/mole. ^e Added in proof: R. J. L. Anden, *et al.*, *J. Chem. Soc., A*, 1894 (1968).

Table XIII
Acids, Esters, and Anhydrides

Compound ^a	ΔH_f°		Δ	n, σ	Ref	σ	S°		Δ	Ref	$C_p^{300^\circ}$		Δ	$C_p^{800^\circ}$		Δ	Ref
	Obsd	Estd					Obsd	Estd			Obsd	Estd		Obsd	Estd		
Formic acid, HCOOH	-90.5	-89.9	-0.6		98	1	59.4	59.4	0.0	98	10.8	10.8	0.0	18.3	18.4	-0.1	98
Acetic acid, CH ₃ COOH	-103.8	-103.8	0.0		107, 108	3	67.5	68.3	-0.8	107	16.0	16.0	0.0	29.1	29.1	0.0	107
Methyl formate, HCOOCH ₃	-81.0	-81.0	0.0		52	3	84.6	72.3	12.3	52							
Ethyl acetate, CH ₃ COOC ₂ H ₅	-103.4	-103.4	0.0		52	3 ²	90.1	89.9	0.2	52							
Methyl benzoate, C ₆ H ₅ COOCH ₃	-71.7	-71.2	-0.5		85												
Ethyl benzoate, C ₆ H ₅ COOC ₂ H ₅	-79.1	-79.7	+0.6		85												
Vinyl acetate, * CH ₃ COOCH=CH ₂	-72.4	-72.4	[0.0]		52, 61												
Methyl methacrylate, * CH ₂ C(CH ₃)COOCH ₃	-79.3 ^b	-79.3	[0.0]		61												
Methyl <i>trans</i> -cinnamate, C ₆ H ₅ CH=CHCOOCH ₃	-49.3 ^c	-48.2	-1.1		109												
Diethyl fumarate, (CHCOOC ₂ H ₅) ₂	-168.9 ^d	-170.1	+1.2		109												
Acetic anhydride, * (CH ₃ CO) ₂ O	-137.9	-137.9	[0.0]		97												

^a An asterisk denotes unique group. ^b Using a calculated value of -107.9 for methyl isobutyrate. ^c Using a calculated value of -79.0 for methyl β -phenylpropionate. ^d Using a calculated value of -198.2 for diethyl succinate.

Table XIV
Aliphatic and Aromatic Amines^a

Compound	ΔH_f°		Δ	Ref	n, σ	Ref	S°		Δ	Ref	$C_p^{300^\circ}$		Δ	$C_p^{800^\circ}$		Δ	Ref
	Obsd	Estd					Obsd	Estd			Obsd	Estd		Obsd	Estd		
CH ₃ NH ₂	-5.5	-5.3	-0.2	110	3	57.93*	57.93	[0.0]	111	11.91*	11.91	[0.0]	22.43*	22.43	[0.0]	111	
(CH ₃) ₂ NH	-4.5	-4.8	0.3	110	3 ²	65.39*	65.39	[0.0]	112	16.58*	16.58	[0.0]	33.94*	33.94	[0.0]	112	
(CH ₃) ₃ N	-5.9	-5.8	-0.1	110	3 ⁴	69.15*	69.15	0.0	112	22.05*	22.05	[0.0]	45.62*	45.62	[0.0]	112	
C ₂ H ₅ NH ₂	-11.0	-11.9	0.9	110		...	67.7		...	17.56	17.56		...	33.52		...	
(C ₂ H ₅) ₂ NH	-17.2	-18.0	0.8	110		...	85.0		...	27.08	27.08		...	56.12		...	
(C ₂ H ₅) ₃ N	-23.9	-23.2	-0.7	110		...	98.4		...	37.80	37.80		...	78.89		...	
<i>n</i> -C ₃ H ₇ NH ₂	-16.77	-16.80	0.03	120		
<i>i</i> -C ₃ H ₇ NH ₂	-20.02	-20.6	0.58	120		
<i>n</i> -C ₄ H ₉ NH ₂	-21.8	-22.8	1.0	113 ^b		
<i>sec</i> -C ₄ H ₉ NH ₂	[-24.7]*	-24.7	[0.0]	113 ^b		
<i>t</i> -C ₄ H ₉ NH ₂	-28.6*	-28.6	[0.0]	113 ^b		
<i>n</i> -C ₄ H ₉ NH- <i>i</i> -C ₄ H ₉	-40.2	-39.1	-1.1	113 ^c		
C ₆ H ₅ NH ₂	20.8*	20.8	[0.0]	115	2	76.44*	76.44	[0.0]	115	26.07*	26.07	[0.0]	53.79*	53.79*	[0.0]	115	
C ₆ H ₅ NHCH ₃	21.6	20.8	0.8	116													
C ₆ H ₅ N(CH ₃) ₂	22.8	22.0	0.8	116													
C ₆ H ₅ NHC ₂ H ₅	13.4	14.2	-0.8	117													
C ₆ H ₅ N(C ₂ H ₅) ₂	9.6	10.4	-0.8	117													
(C ₆ H ₅) ₂ NH	49.9*	49.9	[0.0]	118													
				119													

^a Each piece of data marked with an asterisk was the sole source for determining a particular group value, and therefore these data necessarily fit the group predictions. ^b ΔH_f° revised (see text). ^c ΔH_f° revised (see text) to 11.4 kcal/mole.

Table XV

Imines

Compound	ΔH_f°		Δ	Ref	S°		$300^\circ K$		$800^\circ K$	
	Obsd	Estd			Obsd	Estd	Obsd	Estd	Obsd	Estd
$CH_3CH=N-n-C_3H_7$	-1.2	-1.6	0.8	121 ^a						
$C_2H_5CH=NC_2H_5$	-0.0	-1.7	1.7	121 ^a						
$C_2H_5CH=N-n-C_3H_7$	-8.1	-6.6	-1.5	121 ^a						
$C_2H_5CH=N-i-C_3H_7$	-9.6	-9.6	0.0	121 ^a						
$i-C_3H_7CH_2=NC_2H_5$	-7.5	-8.6	1.1	121 ^a						
$i-C_3H_7CH=N-n-C_4H_9$	-20.4	-18.5	-1.9	114 ^b						

^a ΔH_f° for all the others calculated from liquid-phase heats of hydrogenation quoted in ref 121 as follows. To obtain gas-phase heats of hydrogenation 0.5 kcal/mole is added to liquid-phase values to allow for differences in ΔH_v° . From these gas-phase heats of hydrogenation and group estimates of ΔH_f° for corresponding amines, ΔH_f° values for the imines were obtained. ^b ΔH_v° revised (see text) to 11.4 kcal/mole.




Figure 3.

Additional independent evidence for the size of the alkane *gauche* interaction is also obtained from the substituted cyclohexanes. For example, ΔH_f° for *cis*-1,4-dimethylcyclohexane is 1.9 kcal/mole greater than for the *trans* isomer. In Figure 3 we see that in the *trans* isomer the methyl groups can be all equatorial or all axial, whereas in the *cis* isomer there must always be one axial methyl group. From the Newman diagrams in Figure 3, it is seen that the axial methyl group has *gauche* interaction with each of the two next-nearest ring carbon atoms. This would lead to an expected decrease in stability by $2 \times 0.8 = 1.6$ kcal/mole for the *cis* isomer, and this is very close to the observed 1.9 kcal/mole. For 1,3-dimethylcyclohexane, the opposite is true, *i.e.*, the *trans* isomer must have one axial methyl group, and we would predict it should be less stable than the *cis* isomer by 1.6 kcal/mole. The observed difference is 1.96 kcal/mole. In *trans*-1,2-dimethylcyclohexane, the two Me groups are equatorial, generating one *gauche* interaction. In the *cis* isomer, one Me group is equatorial and the other axial, generating three *gauche* interactions, since they are on adjacent carbon atoms. The observed difference in ΔH_f° is 1.87 kcal/mole, again equal to two *gauche* interactions within experimental error. *Gauche* interactions between ring carbon atoms are included in the ring correction.

This treatment of the cyclic compounds does not strictly adhere to our definition of the law of group additivity, since we cannot write disproportionation reactions that open the ring. However, this is generally a minor deviation.

Use of the open-chain hydrocarbon groups can cause some difficulties. For instance, the use of the values derived from conjugated dienes for the entropy of ring systems, like methylcyclobutene in conjunction with the entropy ring correction for cyclobutene itself, overcorrects for the stiffness of the ring since the diene has some stiffness already. In cases of this type, it is better to use the groups from unconjugated olefins.

Having established the ring corrections, we may use them to get group values for cyclic compounds. For instance, the

heat of hydrogenation of 1,4-dihydronaphthalene to tetralin yields a value for the group $C-(C_B)(C_d)(H)_2$ if the ring correction for tetralin is taken as the sum of the strains in cyclohexene and that in 1,4-cyclohexadiene.

Such transferability of ring corrections must involve caution. For instance, the heat of formation of 1,2-dihydronaphthalene is predicted exactly, only if the ring correction is adjusted for the difference in conjugation energy of styrene and 1,2-dienes. Where ΔH_2 = heat of hydrogenation, the conjugation energies are defined as

$$\begin{aligned} \text{conjugation energy of styrene} &= \Delta H_{H_2}^\circ(\text{styrene}) - \\ &\quad \Delta H_{H_2}^\circ(\text{mono-1-olefin}) = 1.7 \text{ kcal/mole} \\ \Delta H_{H_2}^\circ(\text{styrene}) &= [C_B-(C)] - [C_B-(C_d)] + [C-(C)(C_B)(H)_2] - \\ &\quad [C_d-(C_B)(H)] - [C_d-(H)_2] + [C-(H)_3(C)] \\ \Delta H_{H_2}^\circ(\text{mono-1-olefin}) &= [C-(C)_2(H)_2] - [C-(C_d)(C)(H)_2] + \\ &\quad [C-(C)_2(H)_2] - [C_d-(C)(H)] - [C_d-(H)_2] + [C-(H)_3(C)] \end{aligned}$$

$$\begin{aligned} \text{conjugation energy of 1,3-dienes} &= \Delta H_{H_2}^\circ(\text{diene}) - \\ &\quad \Delta H_{H_2}^\circ(\text{mono-1-olefin}) = 4.07 \text{ kcal/mole} \\ \Delta H_{H_2}^\circ(1,3\text{-diene}) &= [C_d-(C)(H)] + [C-(C_d)(C)(H)_2] - \\ &\quad [2C_d-(C_d)(H)] + [C-(C)(H)_3] - [C_d-(H)_2] \end{aligned}$$

Thus, the ring correction for 1,2-dihydronaphthalene becomes

$$\text{ring correction (cyclohexadiene-1,3)} - (4.0 - 1.7) = 2.5 \text{ kcal/mole}$$

In conclusion, we may summarize the usefulness of the law of group additivity, as applied to hydrocarbons, as follows. In the prediction of entropy and heat capacity, the additivity of group properties generally gives agreement with experiment within ± 0.5 gibbs/mole. In the estimation of heats of formation, the agreement is generally in the range ± 0.5 kcal/mole. For hydrocarbons, enough data of this type exist to make suspect any data which deviates from group additivity by more than 1 gibbs/mole or 1 kcal/mole. Exceptions to this rule will appear in the case of very heavily branched compounds.

This section prepared by S. W. Benson, H. E. O'Neal, and D. M. Golden.

IV. Oxygen-Containing Compounds

The three main sources of original data used were the Benson and Buss paper,² a valuable review by Green,⁵² and the IUPAC Bulletins of Thermodynamics and Thermochemistry.⁵³

(52) J. H. Green, *Quart. Rev. (London)*, **15**, 125 (1961).

(53) Bulletins of Thermodynamics and Thermochemistry, IUPAC, University of Michigan, Ann Arbor, Mich.

The group values are in Table XXXIV. In Tables VIII–XIII,^{54–109} observed data are compared with those estimated

- (54) National Bureau of Standards Technical Note 270-1, U. S. Government Printing Office, Washington, D. C., 1965.
- (55) E. V. Ivash, J. C. M. Li, and K. S. Pitzer, *J. Chem. Phys.*, **23**, 1814 (1955).
- (56) J. H. S. Green, *Trans. Faraday Soc.*, **57**, 2132 (1961).
- (57) E. Buckley and J. D. Cox, *ibid.*, **63**, 895 (1967).
- (58) J. F. Mathews and J. J. McKetta, *J. Phys. Chem.*, **65**, 758 (1961).
- (59) J. H. S. Green, *Chem. Ind.* (London), 1215 (1960).
- (60) H. A. Skinner and A. Snelson, *Trans. Faraday Soc.*, **56**, 1776 (1960).
- (61) M. A. Dolliver, T. L. Gresham, G. B. Kistiakowsky, E. A. Smith, and W. E. Vaughan, *J. Am. Chem. Soc.*, **60**, 440 (1938).
- (62) G. S. Parks, K. E. Manchester, and L. M. Vaughan, *J. Chem. Phys.*, **22**, 2089 (1954).
- (63) G. S. Parks, T. J. West, B. F. Naylor, P. S. Fujii, and L. A. McLaine, *J. Am. Chem. Soc.*, **68**, 2524 (1946).
- (64) M. E. Dyatkina, *Zh. Fiz. Khim.*, **28**, 377 (1954).
- (65) J. H. S. Green, *Trans. Faraday Soc.*, **59**, 1559 (1963).
- (66) N. S. Berman and J. J. McKetta, *J. Phys. Chem.*, **66**, 1444 (1962).
- (67) G. S. Parks, J. R. Moseley, and P. V. Patterson, *J. Chem. Phys.*, **18**, 152 (1950).
- (68) E. T. Beynon and J. J. McKetta, *J. Phys. Chem.*, **67**, 2761 (1963).
- (69) R. P. Bell and J. C. Clunie, *Trans. Faraday Soc.*, **48**, 439 (1952).
- (70) R. J. L. Andon, D. P. Biddiscombe, J. D. Cox, R. Handley, D. Harrop, E. F. G. Herington, and J. F. Martin, *J. Chem. Soc.*, 5246 (1960).
- (71) J. C. Evans, *Spectrochim. Acta*, **16**, 1382 (1960).
- (72) J. H. S. Green, *Chem. Ind.* (London), 1575 (1962).
- (73) G. Pilcher, A. S. Pell, and D. J. Coleman, *Trans. Faraday Soc.*, **60**, 499 (1964).
- (74) Z. Seha, *Chem. Listy*, **49**, 1569 (1955).
- (75) G. Pilcher, H. A. Skinner, A. S. Pell, and A. E. Pope, *Trans. Faraday Soc.*, **59**, 316 (1963).
- (76) M. Colomina, A. S. Pell, H. A. Skinner, and D. J. Coleman, *ibid.*, **61**, 2641 (1965).
- (77) E. J. Smutny and A. Bondi, *J. Phys. Chem.*, **65**, 546 (1961).
- (78) M. Badoche, *Bull. Soc. Chim. France*, **8**, 212 (1941).
- (79) R. C. Cass, S. E. Fletcher, C. T. Mortimer, H. D. Springall, and T. R. White, *J. Chem. Soc.*, 1406 (1958).
- (80) J. H. Stern and F. H. Dover, *J. Phys. Chem.*, **66**, 97 (1962).
- (81) G. Baker, J. H. Littlefair, R. Shaw, and J. C. J. Thynne, *J. Chem. Soc.*, 6970 (1965).
- (82) P. A. Giguere and J. D. Liu, *J. Am. Chem. Soc.*, **77**, 6477 (1955).
- (83) W. E. Vaughan, *Discussions Faraday Soc.*, **10**, 330 (1951).
- (84) W. Pritzkow and K. A. Muller, *Chem. Ber.*, **89**, 2316 (1956).
- (85) P. Gray and A. Williams, *Chem. Rev.*, **59**, 239 (1959).
- (86) J. W. Breitenbach and J. Derkosk, *Monatsh.*, **82**, 117 (1951).
- (87) L. Jaffe, E. J. Prosen, and M. Szwarc, *J. Chem. Phys.*, **27**, 416 (1957).
- (88) A. S. Pell and G. Pilcher, *Trans. Faraday Soc.*, **61**, 71 (1965).
- (89) S. Sundaram, *Z. Physik. Chem.* (Frankfurt), **36**, 376 (1963).
- (90) A. A. Vvedenskii, *Russ. J. Phys. Chem.*, **40**, 1048 (1966).
- (91) F. L. Oetting, *J. Chem. Phys.*, **41**, 149 (1964).
- (92) R. F. Zurcher and H. H. Gunthard, *Helv. Chim. Acta*, **38**, 849 (1955).
- (93) A. Snelson and H. A. Skinner, *Trans. Faraday Soc.*, **57**, 2125 (1961).
- (94) K. Pihlaja and J. Heikkilä, *Acta Chem. Scand.*, **21**, 2390 (1967).
- (95) G. B. Guthrie, D. W. Scott, W. N. Hubbard, C. Katz, J. P. McCullough, M. E. Gross, K. D. Williamson, and G. Waddington, *J. Am. Chem. Soc.*, **74**, 4662 (1952).
- (96) J. B. Conn, G. B. Kistiakowsky, and E. A. Smith, *ibid.*, **61**, 1868 (1939).
- (97) C. P. Keszthelyi and H. E. O'Neal, submitted for publication.
- (98) J. H. S. Green, *J. Chem. Soc.*, 2241 (1961).
- (99) K. S. Pitzer and W. Weltner, *J. Am. Chem. Soc.*, **71**, 2842 (1949).
- (100) A. A. Antonov and P. G. Maslov, *Russ. J. Phys. Chem.*, **38**, 318 (1964).
- (101) I. A. Vasil'ev and A. A. Vvedenskii, *ibid.*, **40**, 453 (1966).
- (102) R. E. Pennington and K. E. Kobe, *J. Am. Chem. Soc.*, **79**, 300 (1957).
- (103) G. C. Sinke and F. L. Oetting, *J. Phys. Chem.*, **68**, 1354 (1964).
- (104) P. Landrieu, F. Bayloca, and J. R. Johnson, *Bull. Soc. Chim. France*, **45**, 36 (1929).
- (105) P. Sellers, IUPAC, "Thermodynamik Symposium," Heidelberg, Sept 1967; *Acta Chem. Scand.*, in press.
- (106) G. R. Nicholson, *J. Chem. Soc.*, 2431 (1957).
- (107) W. Weltner, *J. Am. Chem. Soc.*, **77**, 3941 (1955).
- (108) F. W. Evans and H. A. Skinner, *Trans. Faraday Soc.*, **55**, 260 (1959).
- (109) R. B. Williams, *J. Am. Chem. Soc.*, **64**, 1395 (1942).

from groups. There are more groups and less observed data for oxygen compounds than for hydrocarbons. Consequently, there are many more sole sources for groups. But in the cases where there is enough information to make a reasonable test, such as in the heats of formation of the alcohols and ethers, it is clear that the principle of group additivity is valid. Compared with the number of heats of formation, there are very few entropies and heat capacities. For example, it is surprising that dimethyl ether is the only aliphatic ether for which the entropy or heat capacity has been measured.

The data are insufficiently precise or extensive to fix a good value for the *gauche* interactions in branched compounds. The value selected (0.3 kcal/mole) fits the data quite well, but values smaller or larger by 0.2 kcal/mole would do about as well.

This section prepared by S. W. Benson and R. Shaw.

V. Nitrogen-Containing Compounds

As with the oxygen-containing compounds, there are much less data for the nitrogen compounds than for the hydrocarbons. Consequently, the accuracy of the nitrogen groups is more variable and not so high as for the hydrocarbons. In some cases interpolation methods and similar approximations have been necessary to obtain a group value. No values are available for nitrogen compounds containing carbonyl groups, halogen atoms, etc., so that any complex intramolecular interactions in such compounds cannot be included in any estimate by group additivity.

A. AMINES

1. Aliphatic

The data and group predictions are presented in Table XIV.^{110–120} In order to obtain the estimates for entropy and specific heats for all except the three methylamines, group values were used which have been obtained from compounds (RNH₂) by interpolating between the corresponding alcohols (ROH) and alkanes (RCH₃). This procedure was checked for CH₃NH₂, taking the arithmetic mean of values for CH₃OH and CH₃CH₃ (symmetry corrected for the entropy). The errors involved were less than 0.5 eu.

Some of the gas-phase heats of formation were suspected to be slightly in error, due to errors in estimated or measured heats of vaporization. These we revised, using Trouton's rule (and in the case of these amines, a Trouton constant, $S_T = 22$ eu, which may still be a trifle low) and corrected to 25°, using $\Delta C_p(\text{vap}) \approx 15$ eu.

- (110) I. Jaffe, Thesis, University of Maryland, 1958.
- (111) A. A. Vvedenskii and V. M. Petrov, *Russ. J. Phys. Chem.*, **39**, 812 (1965).
- (112) K. A. Kobe and R. H. Harrison, *Petrol. Refiner*, **33** (11), 161 (1954).
- (113) F. W. Evans, D. M. Fairbrother, and H. A. Skinner, *Trans. Faraday Soc.*, **55**, 399 (1959).
- (114) A. F. Bedford, P. B. Edmondson, and C. T. Mortimer, *J. Chem. Soc.*, 2927 (1962).
- (115) W. E. Hatton, D. L. Hildebrand, G. C. Sinke, and D. R. Stull, *J. Chem. Eng. Data*, **7**, 229 (1962).
- (116) J. A. Kerr, R. C. Sekhar, and A. F. Trotman-Dickenson, *J. Chem. Soc.*, 3217 (1963).
- (117) G. N. Vriens and A. G. Hill, *Ind. Eng. Chem.*, **44**, 2732 (1952).
- (118) L. G. Cole and E. C. Gilbert, *J. Am. Chem. Soc.*, **73**, 5426 (1951).
- (119) D. J. Coleman and H. A. Skinner, *Trans. Faraday Soc.*, **62**, 2057 (1966).
- (120) N. K. Smith and W. D. Good, *J. Chem. Eng. Data*, **12**, 572 (1967).

2. Aromatic

The data and group predictions are shown in Table XIV. It should be noted at this point that for single side chains attached to benzene rings the possibility of nonbonded steric interactions arises, even if the side chains are unbranched. For instance, in *N*-methylaniline, by analogy with ethylbenzene, the terminal methyl group, will be close ($\leq 90^\circ$) to one or other of the *ortho* positions of the benzene ring. This interaction cannot be classed as *gauche* in the strictest sense, and for simplicity's sake we have included it in the relevant group. For example, $C_6H_5N(C_2H_5)_2$ has only two "*gauche*" corrections worth 1.6 kcal/mole in all.

B. IMINES

As can be seen from Table XV,¹²¹ only ΔH_f° values are known for a few imines, but the internal consistency is fairly reasonable.

In Table XXXV, as stated, the N_I refers to the nitrogen in these compounds. In order to separate this group value from others in the molecule, the carbon atom end of the double bond was taken as a C_2 atom (as in olefins), and it was assumed that the groups, $C-(N_I)(C)(H)_2$, $C-(N_I)(C)_2(H)$, were the same as $C-(N)(C)(H)_2$, $C-(N)(C)_2(H)$. This assumption holds well for olefins where it can be seen that $C-(C_d)(C)(H)_2$ is almost the same as $C-(C)_2(H)_2$.

C. CYANIDES (OR NITRILES)

The data are compared with the group predictions in Table XVI.¹²²⁻¹³⁶ Recent ΔH_f° values for alkyl cyanides are particularly sparse. A check on the group predictions for ethyl cyanide would be particularly valuable, for instance. The deviations observed for *trans*-dicyanoethylene may be real and may indicate a conjugation of electrons sufficient to cause departure from group additivity.

D. NITRO COMPOUNDS

The data are compared with group predictions in Table XVII. Since the only entropy data for nitro compounds were those of CH_3NO_2 , it was necessary to use bond additivity to generate the entropy estimates listed here. It was from these estimates (symmetry corrected) that the group values listed in Table XXXV were calculated.

- (121) L. M. Jackson and D. I. Packam, *Proc. Chem. Soc.*, 349 (1957).
 (122) N. E. Duncan and G. J. Janz, *J. Chem. Phys.*, **23**, 434 (1955).
 (123) L. A. Weber and J. E. Kilpatrick, *ibid.*, **36**, 829 (1962).
 (124) F. W. Evans and H. A. Skinner, *Trans. Faraday Soc.*, **55**, 255 (1959).
 (125) J. H. S. Green, *Spectrochim. Acta*, **17**, 607 (1961).
 (126) L. H. Daly and S. E. Wiberly, *J. Mol. Spectry.*, **2**, 177 (1958).
 (127) C. A. Wulff and E. F. Westrum, *J. Phys. Chem.*, **67**, 2376 (1963).
 (128) H. L. Clever, C. A. Wulff, and E. F. Westrum, *ibid.*, **69**, 1983 (1965).
 (129) F. Halverson, R. F. Stamm, and J. J. Whalen, *J. Chem. Phys.*, **16**, 808 (1948).
 (130) H. S. Davis and O. F. Wiedeman, *Ind. Eng. Chem.*, **37**, 482 (1945).
 (131) H. R. Wyss and H. H. Günthard, *Helv. Chim. Acta*, **44**, 625 (1961).
 (132) R. H. Boyd, *J. Chem. Phys.*, **38**, 2529 (1963).
 (133) "JANAF Interim Thermochemical Tables," D. R. Stull, Ed., Dow Chemical Co., Midland, Mich., 1960-1966.
 (134) E. F. Westrum, Jr., and A. Ribner, *J. Phys. Chem.*, **71**, 1216 (1967).
 (135) A. Ribner and E. F. Westrum, Jr., *ibid.*, **71**, 1208 (1967).
 (136) R. H. Boyd, K. R. Guha, and R. Winthrick, *ibid.*, **71**, 2187 (1967).

Table XVI
Cyanides^a

Compound	ΔH_f°		Δ	Ref	n, σ	S°		Δ	C_p° (300°K)		Δ	Ref	
	Obsd	Estid				Obsd	Estid		Obsd	Estid			Obsd
C_2H_5CN	12.4	68.75	68.4	0.35	123	17.30*	17.30	[0.0]	122
<i>n</i> - C_3H_7CN	7.45*	7.45	[0.0]	124	77.8	43.8
<i>i</i> - C_3H_7CN	5.62*	5.62	[0.0]	124	76.2	43.4
C_4H_9CN	52.3*	52.3	[0.0]	124	...	76.89*	76.89	[0.0]	125	26.2*	26.2	[0.0]	125
<i>c</i> - C_3H_6CN	...	43.5	70.7*	70.7	[0.0]	126 ^b	18.95*	18.95	[0.0]	126
$NCCH_2CH_2CN$...	45.0	79.10	79.0	0.1	127	...	22.2
$NCCH_2CH_2CH_2CN$...	40.0	88.1	88.51	-0.41	128	...	27.7
$(CH_3)_3CN$	79.63*	79.63	[0.0]	134
$CH_2=CHCN$	43.7	43.6	0.1	130	...	65.56	64.20	1.36	129	15.30	14.90	0.4	129
<i>trans</i> - $CH_2CH=CHCN$	35.8	35.9	-0.1	131	...	71.43	72.8	-1.37	131	19.71	20.10	-0.39	131
<i>trans</i> - $NCCH=CHCN$	81.3	74.8	6.5	136
$C_2H_3(CN)_2$	123.9	122.5	1.4	132
$C_2(CN)_4$	168.5	171.3	-2.8	132
$C_2(CN)_2$	127.5*	127.5	[0.0]	133	...	69.44*	69.44	[0.0]	133	20.58*	20.60	[0.0]	133
$(CH_3)_2C(CN)_2$	83.42*	83.42	[0.0]	135

^a Each piece of data marked with an asterisk was the sole source for determining a particular group value, and therefore these data necessarily fit the group predictions. ^b Note the quoted value for S° in ref 126 is in error.

Table XVII
Nitro Compounds^a

Compound	ΔH_f°		Δ	Ref	n, σ	S°		$300^\circ K$		$800^\circ K$	
	Obsd	Estd				Obsd	Estd	Obsd	Estd	Obsd	Estd
$C_2H_5NO_2$	-23.8	-24.4	0.6	139 ^b	3 × 2						
$n-C_3H_7NO_2$	-29.7	-29.3	-0.4	139	3 × 2						
$i-C_3H_7NO_2$	-33.9	-34.4	0.5	139	3 ² × 2						
$n-C_4H_9NO_2$	-34.4	-34.3	-0.1	139	3 × 2						
$sec-C_4H_9NO_2$	-39.1	-38.5	-0.6	139	3 ² × 2						
$C_2H_5CH(NO_2)_2$	-25.9*	-25.9	[0.0]	139							

^a Each piece of data marked with an asterisk was the sole source for determining a particular group value, and therefore these data necessarily fit the group predictions. ^b ΔH_f° revised to 9.7 kcal/mole.

E. NITRITES AND NITRATES

The data are compared with group predictions in Table XVIII. Heats of vaporization were generally consistent with a Trouton constant of 22.5 eu, for both classes of compounds, with the exception of ΔH_v° ($n-C_3H_7ONO$). Consequently, the latter was revised in accordance with $S_T = 22.5$ eu.

F. HYDRAZINES, AZO COMPOUNDS, AND TETRAZINES

The data are compared with group predictions in Table XIX.¹⁴⁸⁻¹⁵⁰ Information on azo compounds is particularly lacking. It is interesting to note here a much quoted value for $\Delta H_f^\circ(CH_3N=NCH_3)$ of 43.2 kcal/mole.¹³⁷ This value was never measured, but merely estimated from $\Delta H_f^\circ(i-C_3H_7N=NiC_3H_7)$ ¹³⁸ by a cruder additivity rule, which is, in principle, less reliable than that used in this paper. The data of Table XXXV would predict $\Delta H_f^\circ(CH_3N=NCH_3) = 44.7$ kcal/mole. It should be pointed out, however, that in deriving the group values for azo compounds, it was assumed that the increments in replacing H by Me in CH_3NNCH_3 were the same as the corresponding increments in the isoelectronic olefins $CH_3CH=CHCH_3$.

G. AMIDES AND IMIDES

The data are compared with group predictions in Table XX.¹⁶¹⁻¹⁵⁴ Unfortunately, there are no checks whatsoever on the group values for heats of formation among the amides,

and therefore this table probably contains the least reliable information of any class of nitrogen-containing compounds listed thus far. However, the method of measuring heats of hydrolysis for many of these amides ought to give values for ΔH_f° at least as reliable as combustion measurements, and it is felt that these data are probably good to better than ± 2 kcal/mole.

H. NITROGEN-CONTAINING RING COMPOUNDS

The data for saturated nitrogen-containing ring compounds are included as corrections to group predictions and listed in Table XXXV. Comparison of these ring corrections with those of sections III and IV shows once again that, both for heats and entropies, they tend to depend primarily on ring size and not significantly on whether a $>CH_2$ has been replaced by an $>O$ or an $>NH$ group. Such a rule has been used to estimate several of the ring corrections, where data are lacking. Application of this rule must obviously be made with caution, since the ring correction for ΔH_f° for succinimide appears to be slightly high for a five-membered ring. In this particular case, however, we feel the data and imide group values are not sufficiently reliable to predict this ring correction to better than ± 2 kcal/mole and thus invalidate the rule.

The data for unsaturated, nitrogen-containing heterocyclic ring compounds (most ΔH_f° values) are listed in Table XXI.¹⁵⁵⁻¹⁶² These data do not readily extend themselves to the method of ring corrections by group predictions, since all of these compounds show resonance. The problem of estimation of the resonance energies involved depends on establishment of a proper standard of comparison. This is rather an involved and complex problem, and we feel that it would not be appropriate to discuss it here, particularly since it would not contribute to the task of empirical prediction.

(137) M. Page, H. O. Pritchard, and A. F. Trotman-Dickenson, *J. Chem. Soc.*, 3878 (1953).

(138) G. E. Coates and L. E. Sutton, *ibid.*, 1187 (1948).

(139) D. E. Holcomb and C. L. Dorsey, *Ind. Eng. Chem.*, **41**, 2788 (1949).

(140) G. Geiseler and M. Rätzsch, *Z. Physik. Chem. (Frankfurt)*, **26**, 131 (1960).

(141) J. D. Ray and A. A. Gershon, *J. Phys. Chem.*, **66**, 1750 (1962).

(142) J. D. Ray and R. A. Ogg, *ibid.*, **63**, 1522 (1959).

(143) J. C. D. Brand and T. M. Cawthorn, *J. Am. Chem. Soc.*, **77**, 319 (1955).

(144) P. Gray and P. L. Smith, *J. Chem. Soc.*, 769 (1954).

(145) J. G. Aston, H. L. Fink, G. J. Janz, and K. E. Russell, *J. Am. Chem. Soc.*, **73**, 1939 (1951).

(146) R. C. Harshman, *Jet Propulsion*, **27**, 398 (1957).

(147) J. G. Aston, G. J. Janz, and K. E. Russell, *J. Am. Chem. Soc.*, **73**, 1943 (1951).

(148) J. G. Aston, J. L. Wood, and T. P. Zolki, *ibid.*, **75**, 6202 (1953).

(149) T. M. Donovan, C. H. Shomate, and W. R. McBride, *J. Phys. Chem.*, **64**, 281 (1960).

(150) J. G. Aston, T. P. Zolki, and J. L. Wood, *J. Am. Chem. Soc.*, **77**, 281 (1955).

(151) Z. Seha, *Collection Czech. Chem. Commun.*, **26**, 2435 (1961).

(152) A. Bauder and H. H. Günthard, *Helv. Chim. Acta*, **41**, 670 (1958).

(153) A. A. Antonov, *Zh. Obsch. Khim.*, **34**, 2340 (1964).

(154) I. Wadsö, *Acta Chem. Scand.*, **19**, 1079 (1965).

(155) D. W. Scott, W. T. Berg, I. A. Hossenlopp, W. N. Hubbard, J. F. Messerley, S. S. Todd, D. R. Douslin, J. P. McCullough, and G. Waddington, *J. Phys. Chem.*, **71**, 2263 (1967).

(156) W. N. Hubbard, F. R. Frow, and G. Waddington, *J. Phys. Chem.*, **65**, 1326 (1961).

(157) J. P. McCullough, D. R. Douslin, J. F. Messerley, I. A. Hossenlopp, T. C. Kinchenloe, and G. Waddington, *J. Am. Chem. Soc.*, **79**, 4289 (1957).

(158) D. W. Scott, W. N. Hubbard, J. F. Messerley, S. S. Todd, I. A. Hossenlopp, W. D. Good, D. R. Douslin, and J. P. McCullough, *J. Phys. Chem.*, **67**, 680 (1963).

(159) D. W. Scott, W. D. Good, G. B. Guthrie, S. S. Todd, I. A. Hossenlopp, A. G. Osborn, and J. P. McCullough, *ibid.*, **67**, 685 (1963).

(160) R. J. L. Andon, J. D. Cox, E. F. G. Herington, and J. F. Martin, *Trans. Faraday Soc.*, **53**, 1074 (1957).

(161) J. D. Cox, *ibid.*, **56**, 959 (1960).

(162) J. Tjebbes, *Acta Chem. Scand.*, **16**, 916 (1962).

Table XVIII
Nitrites and Nitrates^a

Compound	ΔH_f°		Δ	Ref	n, σ	S°		Δ	Ref	C_p°		Δ	Ref
	Obsd	Estd				Obsd	Estd			Obsd	Estd		
CH ₃ ONO	-15.6	-16.0	0.4	141	3	70.14*	70.14	[0.0]	140	15.36*	15.36	[0.0]	140
C ₂ H ₅ ONO	-24.2	-24.5	0.3	85	3	...	80.6	20.35
<i>n</i> -C ₃ H ₇ ONO	-30.1	-29.4	-0.7	85/	3	...	90.10	25.85
CH ₃ ONO ₂	-28.6	-28.7	0.1	142, 141	3 × 2	75.97	75.30	0.67	143
C ₂ H ₅ ONO ₂	-36.8	-37.2	0.4	85	3 × 2	85.14	85.80	-0.66	144
<i>n</i> -C ₃ H ₇ ONO ₂	-41.8	-42.1	0.3	85	3 × 2	...	95.20
<i>i</i> -C ₃ H ₇ ONO ₂	-45.7	-45.0	-0.7	85	3 × 2	...	92.20

^a Each piece of data marked with an asterisk was the sole source for determining a particular group value, and therefore these data necessarily fit the group predictions. ^b ΔH_f° revised to 7.9 kcal/mole.

Table XIX
Hydrazines, Azo Compounds, and Tetrazines

Compound	ΔH_f°		Δ	Ref	n, σ	S°		Δ	Ref	C_p°		Δ	Ref
	Obsd	Estd				Obsd	Estd			Obsd	Estd		
N ₂ H ₄	22.8	22.8	0.0	133	2, 2	57.03	56.9	0.13	133	12.19*	12.19	[0.0]	133
CH ₃ NNH ₂	22.3	22.2	0.1	146	2, 1	66.72	67.0	-0.28	145	17.1*	17.1	[0.0]	145
CH ₃ NNHCH ₃	21.6	21.6	0.0	146	2, 2	74.39	74.3	0.09	147	...	22.0
(CH ₃) ₂ NNH ₂	20.4	20.4	0.0	148, 149	2, 1	72.82	71.80	1.02	148
(CH ₃) ₂ NNHCH ₃	...	20.7	2, 1	79.45	80.50	-1.05	150
C ₆ H ₅ NNH ₂	50.1	50.3	-0.2	118, 119
C ₆ H ₅ NNHCH ₃	77.9	77.8	0.1	118, 119
NH=NH	50.2*	50.2	[0.0]	133	...	52.22*	52.22	[0.0]	133	8.75*	8.75	[0.0]	133
<i>i</i> -C ₃ H ₇ N=N- <i>i</i> -C ₃ H ₇	19.4*	19.5	[0.0]	138
(CH ₃) ₂ NN=N(CH ₃) ₂	65.7*	65.7	[0.0]	149


















^a Each piece of data marked with an asterisk was the sole source for determining a particular group value, and therefore these data necessarily fit the group predictions.

Table XX
Amides and Imides^a

Compound	ΔH_f°		Δ	Ref	n, σ	S°		Δ	Ref	C_p°		Δ	Ref
	Obsd	Estd				Obsd	Estd			Obsd	Estd		
HCONH ₂	-44.5*	-44.5	[0.0]	152	...	59.62*	59.62	[0.0]	151	11.10*	11.1	[0.0]	151
CH ₃ CONH ₂	-57.8*	-57.8	[0.0]	154	3	69.10*	69.1	[0.0]	153	15.63	15.63	[0.0]	153
CH ₃ CONH- <i>n</i> -C ₄ H ₉	-73.1*	-73.1	[0.0]	154 ^b	3 ²	...	105.2
CH ₃ CONHC ₆ H ₅	-24.9*	-24.9	[0.0]	154 ^b
(CH ₃) ₂ CO ₂ NH	-102.7*	-102.7	[0.0]	154 ^b
(CH ₃) ₂ CO ₂ N- <i>n</i> -C ₄ H ₉	-114.2*	-114.2	[0.0]	154 ^b
(CH ₃) ₂ CO ₂ NC ₆ H ₅	-65.5*	-65.5	[0.0]	154 ^b

^a Each piece of data marked with an asterisk was the sole source for determining a particular group value, and therefore these data necessarily fit the group predictions. ^b Calculated from the data of ref 154; known values for ΔH_f° (CH₃COOH) and ΔH_f° (H₂O) given in part IV and known or group estimates for ΔH_f° of various amines.

Table XXI
Heterocyclic N-Atom-Containing Ring Compounds in the Ideal Gas State (1 Atm)

Compound Name	Formula	ΔH_f°		Δ	Ref	n, σ	S° (obsd)	Ref	$300^\circ K$	$800^\circ K$	Ref
		Obsd	Estd ^a						C_p°	C_p°	
									(obsd)	(estd)	
Pyrrole		25.88		...	155	2	64.66		17.16	38.38	
2,5-Dimethylpyrrole		9.1		...	119	...					
Pyrazole		43.3		...	114	...					
Imidazole		30.6		...	114	...					
Pyridine		33.5	33.2	0.3	156	2	67.7	159	18.80	42.49	157
Picolines, α		23.6	23.8	-0.2	158	3	77.8	160	24.05	53.21	158
β		25.3	25.3	0.0	159	3	77.8	161	23.94	53.12	159
γ		24.4	23.8	0.6	160	...					
Dimethylpyridines (lutidines) 2,3		16.3	16.5	-0.2	161	...					
2,4		15.3	14.4	0.9	161	...					
2,5		15.9	15.9	0.0	161	...					
2,6		14.0	14.4	-0.4	161	...					
3,4		16.7	16.5	0.2	161	...					
3,5		17.4	17.4	0.0	161	...					
Pyridazine		66.5		...	162	...					
Pyrimidine		47.0		...	162	...					
Pyrazine		46.9		...	162	...					

^a See text for explanation of method of estimation.


There appear to be small, systematic differences in positional isomers in the pyridine-substituted compounds, favoring *ortho* and *para* substitution. The simplest way to bring these compounds into the system of ring corrections is to use the $N_1-(C_B)$ group to define the pyridine ring, and then make a -1.5 kcal correction for the extra stability due to *ortho* and *para* substitution, and otherwise the same *ortho* corrections as in benzene. The agreement between estimates arrived at thus and observed ΔH_f° values is seen from Table XXI to be good. A comparison of these effects of methyl substitution with those for benzene itself [$\Delta H_f^\circ(C_6H_5CH_3) - \Delta H_f^\circ(C_6H_6) = 7.8$ kcal/mole] shows that methyl substitution on pyridine always produces a slightly greater stabilization than on benzene.

As far as the various isomeric heterocycles containing two ring nitrogen atoms are concerned, it is evident that isomers in which the nitrogen atoms are bonded to one another are significantly less stable than those in which they are not. This could have been qualitatively anticipated on the general grounds of electronegativity.

I. SOME UNIQUE GROUP COMPOUNDS

The above tables represent the extent of the comparison possible between existing data and group predictions. The group additivity method offers no rationale of the properties of particular low molecular weight compounds which are themselves unique groups. In addition, methods are available

Table XXII
Unique Groups

Compound	ΔH_f°	Ref	S°	Ref	C_p°						Ref	
					300°	400°	500°	600°	800°	1000°		1500°
NH ₃	-10.98	133	46.09	133	8.53	9.24	10.04	10.81	12.23	13.47	15.85	133
HCN	32.3	133	48.27	133	8.61	9.42	10.04	10.56	11.45	12.18	13.48	133
CH ₃ CN	(19.1) ^a		58.2	165	12.55	14.66	16.65	18.42	21.32	23.55		165
(CN) ₂	73.87	133	57.80	133	13.59	14.77	15.62	16.31	17.42	18.24	19.43	133
CH ₂ (CN) ₂	63.5	136	69.16	166	17.38	20.22	22.61	24.60	27.72	30.00		166
HNC			48.78	167	9.33	9.97	10.44	10.83	11.49	12.06	13.10	167
CH ₃ NC	35.9	168	59.06	177	12.83	14.80	16.72	18.45	21.34	23.56		177
HN ₃	71.6	169	57.14	169	10.46	11.74	12.81	13.71	15.10	16.12	17.68	169
CH ₂ N ₃	67	170, 163										
CH ₂ N ₂	71	172	58.11	171	12.58	14.23	15.62	16.87	18.73	20.19	22.52	171
	79	174	56.93	173	10.23	12.39	14.34	15.94	18.35	20.04		173
HCNO	-27.9	133	56.98	133	10.57	12.09	13.11	13.93	15.19	16.13	17.62	133
HNO	23.8	133	52.78	133	8.29	8.79	9.34	9.87	10.77	11.45	12.51	133
CH ₃ NO	(16) ^a											
HONO	-18.6	133	59.63	133	10.95	12.36	13.48	14.37	15.66	16.55	17.90	133
HONO ₂	-32.1	133	65.12	133	12.80	15.10	16.93	18.35	20.33	21.61	23.42	133
CH ₃ NO ₂	-17.9	175	65.81	175	13.76	16.80	19.56	21.92	25.56	28.17	32.05	175
N ₂	0	133	45.77	133	6.96	6.99	7.07	7.20	7.51	7.82	8.33	133
NO	21.58	133	50.39	133	7.13	7.16	7.29	7.47	7.83	8.12	8.55	133
NO ₂	7.91	133	57.40	133	8.85	9.60	10.33	10.96	11.88	12.47	13.20	133
NO ₂ sym	16.6	133	60.42	133	11.26	13.37	14.96	16.11	17.51	18.28	19.12	133
NO ₂ unsym	14	176	68	176								
N ₂ O	19.61	133	52.60	133	9.25	10.20	10.95	11.57	12.49	13.11	13.96	133
N ₂ O ₃	19.8	133	74.01	133	15.72	17.39	18.73	19.82	21.39	22.38	23.61	133
N ₂ O ₄	2.54	133	72.84	133	18.52	21.16	23.23	24.86	27.11	28.49	30.16	133
N ₂ O ₅	2.70	133	82.94	133	23.09	26.49	28.97	30.68	32.74	33.81	34.91	133
CH ₂ =CHNO ₂	...		70.6	178	17.43	21.5	24.9	27.7	31.7	34.5	38.5	178

^a Estimates. See text.

for estimation of heats of formation of unstable or nonexistent compounds, as well as those of some classes for which no gas-phase data exist. All this information is discussed in the following sections.

The thermodynamic data for these particular compounds are shown in Table XXII.¹⁶⁵⁻¹⁶⁸ The entropy and specific heat values are almost certainly reliable as are most of the heats of formation. A few of these latter, however, require comment. $\Delta H_f^\circ(\text{CH}_2\text{N}_2)$ is uncertain to ± 5 kcal/mole and $\Delta H_f^\circ(\text{C}-\text{CH}_2\text{N}=\text{N})$ may be even less certain. $\Delta H_f^\circ(\text{CH}_3\text{CN})$ was obtained by interpolation between $\Delta H_f^\circ(\text{HCN})$ and $\Delta H_f^\circ(\text{C}_2\text{H}_5\text{CN})$. The series HCN, CH₃CN, and C₂H₅CN is analogous to HCHO, CH₃CHO, C₂H₅CHO, in that both $-\text{CN}$ and $>\text{C}=\text{O}$ are very polar groups. Thus, we would expect the increments in ΔH_f° to be the same between analogous pairs in these homologous series. This is approximately the case for $\Delta H_f^\circ(\text{HCN}) - \Delta H_f^\circ(\text{HCHO}) = 60.0$ kcal/mole and $\Delta H_f^\circ(\text{C}_2\text{H}_5\text{CN}) - \Delta H_f^\circ(\text{C}_2\text{H}_5\text{CHO}) = 58.2$ kcal/mole. Thus, we have used $\Delta H_f^\circ(\text{CH}_3\text{CN}) - \Delta H_f^\circ(\text{CH}_3\text{CHO}) = 58.8$ kcal/mole to calculate $\Delta H_f^\circ(\text{CH}_3\text{CN})$.

$\Delta H_f^\circ(\text{CH}_3\text{NO})$ was obtained from $\Delta H_f^\circ(\text{CH}_3)$ ¹⁶³ and $\Delta H_f^\circ(\text{NO})$ ¹³³ by assuming $DH^\circ(\text{CH}_3\text{NO}) = 40$ kcal/mole. The

latter value is supported by kinetic evidence from NO-catalyzed isomerization of olefins,¹⁶⁴ although it is not in agreement with an earlier estimate of 57 kcal/mole proposed by Gowenlock, Trotman, and Batt.¹⁷⁹

J. MISCELLANEOUS COMPOUNDS

There are little or no data on gas-phase heats of formation of 1,1-diamines, azines, Schiff bases, oximes, hydroxylamines, nitroso compounds, nitramines, and azides. However, we have been able to estimate some of these, and our results are shown in terms of the group contributions contained in Table XXXVI. Heats of formation, calculated from this table, are unlikely to be more accurate than ± 2 kcal/mole, but are probably not worse. Part of the table is self-explanatory, but the origin of some of the quoted values requires detailed explanation.

Bond dissociation energies can sometimes be useful for estimates of heats of formation, and these have been used to


(169) D. A. Dows and G. C. Pimentel, *J. Chem. Phys.*, **23**, 1258 (1955).(170) B. L. Evans, A. D. Yoffe, and P. Gray, *Chem. Rev.*, **59**, 515 (1959).(171) C. B. Moore and G. C. Pimentel, *J. Chem. Phys.*, **40**, 342 (1964).(172) G. C. Hassler and D. W. Setser, *J. Am. Chem. Soc.*, **87**, 3793 (1965).(173) R. Ettinger, *J. Chem. Phys.*, **40**, 1693 (1964).(174) G. S. Paulett and R. Ettinger *ibid.*, **39**, 825 (1963).(175) J. P. McCullough, D. W. Scott, R. E. Pennington, I. A. Hosselopp, and G. Waddington, *J. Am. Chem. Soc.*, **76**, 4791 (1954).(176) W. A. Guillory and H. S. Johnston, *J. Chem. Phys.*, **42**, 2457 (1965).(177) M. G. Krishna-Pillai and F. F. Cleveland, *J. Mol. Spectry.*, **5**, 212 (1960).(178) K. R. Loos and H. H. Gunthard, *J. Chem. Phys.*, **46**, 1200 (1967).

(179) B. G. Gowenlock, J. Trotman, and L. Batt, Special Publication No. 10, The Chemical Society, London, 1957, p 75.

(163) D. M. Golden, R. Walsh, and S. W. Benson, *J. Am. Chem. Soc.*, **87**, 4053 (1965).(164) S. W. Benson, K. W. Egger, and D. M. Golden, *ibid.*, **87**, 468 (1965).(165) G. A. Crowder and B. R. Cook, *J. Phys. Chem.*, **71**, 914 (1967).(166) F. Halverson and R. J. Francel, *J. Chem. Phys.*, **17**, 694 (1949).(167) D. E. Milligan and M. E. Jacox, *ibid.*, **39**, 712 (1963).

(168) "Selected Values of Chemical Thermodynamic Properties," Circular 500, National Bureau of Standards, Washington, D. C., 1952.

Table XXIII
Heats of Formation of Oximes at 25° and 1 Atm

Oxime	ΔH_f° (cryst)	Ref	$t_B, ^\circ\text{C}$	ΔH_f° (estd) ^a	ΔH_v° (estd) ^a	ΔH_f° (g)	ΔH_f° (g, estd)
CH ₃ CH=NOH	-18.2	168	115	3	10.5	-4.7	-6.3
CH=NOH	-21.1	168	~280	4	17	0	+1.9
CH=NOH							
	-81.6	181	~350	4	20	-7.6	-8.1
Octenal 1-monoxime	-56.8	182	} ~250 {	4	16.5	-36.3	-36.0
Octenal 2-monoxime	-59.3			4	16.5	-38.8	-38.5
Octenal 3-monoxime	-55.4			4	16.5	-36.9	-37.7
Octenal 4-monoxime	-59.8			4	16.5	-39.3	-37.7

^a See text.

obtain the group values for the C-nitroso compounds. It should be pointed out that only the secondary (C-NO) bond strength is known with any assurance, but it would be surprising if other C-NO bond strengths were significantly different. We have, however, made small adjustments to reflect reasonable differences in these bond strengths.

$\Delta H_f^\circ(\text{NH}_2\text{OH}, \text{g})$ has not been measured directly and the value quoted in the table was obtained by combining the value for $\Delta H_f^\circ(\text{NH}_2\text{OH}, \text{aq}) = -23.5$ kcal/mole¹⁸⁰ with an estimate for the heat of solvation of -14.2 kcal/mole. The latter was obtained by taking the mean of the heats of solvation of N_2H_4 (-14.5 kcal/mole)¹⁶⁸ and H_2O_2 (-13.9 kcal/mole).¹⁶⁸ This procedure offers the most reasonable method of allowing for solvent interaction in hydroxylamine solution.

The heats of formation of C-methyl- and O-methylhydroxylamines have never been measured but have been estimated here by consideration of the effects of substituting a methyl group for hydrogen among a series of isoelectronic molecules. The following table for $\Delta H_f^\circ(\text{CH}_3\text{XY}) - \Delta H_f^\circ(\text{HXY})$ summarizes these numbers (in kcal/mole). All values for ΔH_f° used in compiling this table are observed values listed in this paper and sections III and IV, except that for $\Delta H_f^\circ(\text{CH}_3\text{OOH})$, which is a group additivity value of -30.9 kcal/mole and is probably slightly more reliable. If the figures in parentheses are omitted, trends can nevertheless be seen both across the rows and down the columns of this table, and the two figures in parentheses were inserted to be consistent with these trends. From these two values, $\Delta H_f^\circ(\text{CH}_3\text{NHOH})$ and $\Delta H_f^\circ(\text{NH}_2\text{OCH}_3)$ were obtained.

Y	X		
	-CH ₂ -	-NH-	-O-
-CH ₃	-4.6	+1.0	+4.1
-NH ₂	-5.5	-0.5	(+3.1) ^a
-OH	-8.1	(-2.7) ^a	+1.6

^a Obtained by interpolation.

Heats of formation of several solid oximes are known from combustion measurements, and these are listed in Table XXIII.^{181,182} Unfortunately, their heats of sublimation are not


(180) "Selected Values of Chemical Thermodynamic Properties, Series I," Technical Note 270-1, National Bureau of Standards, Washington, D. C., 1965.

(181) C. R. Brent, *Dissertation Abstr.*, **24**, 3108 (1964).


(182) G. Geiseler, M. Rätzsch, K. Ebster, and E. Zeigel, *Ber. Bunsenges. Physik. Chem.*, **70**, 221 (1966).

known with any assurance. Attempts to measure these latter have usually been performed by the effusion technique. However, in most cases no checks were made to discover whether the vapors emerging from the effusion cell were actually those of the oximes, or those of their decomposition fragments. If, as we suspect, the latter was often the case, then serious underestimates of heats of sublimation have been made. Thus, we have disregarded quoted values for heats of sublimation and attempted to estimate them. Our estimates are shown in columns 4 and 5 of Table XXIII. The heats of fusion are typical of organic solids of relatively high molecular weight, while the heats of vaporization were derived from Trouton's rule ($S_T = 23$ eu) and the boiling points (some of which were guessed), as listed in the table. This procedure is admittedly rather crude, but the comparison between $\Delta H_f^\circ(\text{g})$ derived from this procedure, and $\Delta H_f^\circ(\text{g, estd})$ derived from the group value, $\text{N}_1-(\text{OH})$ in Table XXXVI, indicates that the scatter is less than ± 2 kcal/mole. Thus, only if all the values of $\Delta H_f^\circ(\text{g})$ are systematically in error is the oxime group value for ΔH_f° worse than ± 2 kcal/mole.

In Table XXXVI the group contribution of $\text{C}-(\text{N})_2(\text{H})_2$ to ΔH_f° has been derived from the symmetrical triazine shown. The other groups occurring in this ring are all known and the ring strain has been taken by analogy with




as $+2.8$ kcal/mole. In any case, since the group of interest occurs three times in the molecule, errors in estimation of the ring strain will be reduced by a factor of 3 in the group value. For this reason the value ought to be reasonably reliable. The heat of formation of 1,3,5-trimethylhexahydro-1,3,5-triazine has been derived from measurements of the equilibrium constant for



in the temperature range 98–164°, by Gowenlock and

Thomas.¹⁸³ Unfortunately, in their paper, there is an error where by K_{eq} has been wrongly described as K_p but, from examination of the results, it is evidently K_c . Consequently, a quoted value for ΔH° at 35 kcal/mole is, in reality, a value for ΔE° . We have recast their results in the proper units where we find that $\Delta H^\circ = 36.6$ kcal/mole and also that $\Delta S^\circ = 83.5$ eu (at $\sim 400^\circ\text{K}$). However, there is a more serious criticism of these results, which becomes evident when they are compared with the one for the similar equilibrium



Using group values in section III, we calculate that at 400°K , $\Delta S^\circ \simeq 99$ eu for this system. It has been clearly demonstrated¹⁸⁴ that substitutions of $-\text{N}<$ for $-\text{CH}<$ in such molecules scarcely affect entropy values, at or near to room temperature. Moreover, where entropy changes are concerned, any small differences between the above equilibrium and that studied by Gowenlock and Thomas ought to be self-cancelling. Therefore, we would expect that in the former equilibrium ΔS° ought to be $\simeq 99$ eu, also. This argument throws some doubt onto what Gowenlock and Thomas were actually observing. Since they, unfortunately, only made pressure measurements, they could not be certain that side reactions were not occurring. We have taken the point of view that while the observed variation of K_p with temperature may be somewhat in error, the over-all magnitude of K_p in the middle of their temperature range (400°K) is approximately correct. With this assumption, and a value of $\Delta S^\circ \simeq 99$ eu, it can be shown that $\Delta H^\circ = 42.8$ kcal/mole (at 400°K). Furthermore, this value is not particularly sensitive to errors in K_p , since an error of a factor of 10 in the latter would only lead to an error of $\simeq 1.8$ kcal/mole in ΔH° . At room temperature, $\Delta H^\circ = 42.5$ kcal/mole, and when combined with a group additivity estimate of $\Delta H_f^\circ(\text{CH}_3\text{—N=CH}_2) = 17.3$ kcal/mole (which may be obtained directly from Table XXXV), $\Delta H_f^\circ(1) = 9.4$ kcal/mole is obtained.

This section prepared by S. W. Benson and R. Walsh.

VI. Halogen-Containing Compounds

The data for halogen-containing groups, as for the nitrogen groups, come principally from compounds containing no other type of substituent, and conformational corrections differ from those applicable to hydrocarbons. Analysis of the rotational isomers of *n*-propyl chloride,^{185,186} bromide,¹⁸⁶ and fluoride,¹⁸⁷ for example, has indicated that the *gauche* isomer is somewhat more stable (~ 0.1 kcal/mole) than the *trans* isomer. Consequently, no correction for *gauche* interactions between alkyl substituents and halogens will be used. Similarly, no correction will be made for *gauche* interactions between halogens and

amino, alkoxy, thiol, and vinyl substituents. On the other hand, analysis of the 1,2-dihaloethanes of chlorine,^{188,189} bromine,¹⁹¹ and fluorine¹⁹⁰ indicates a value $+1.0$ kcal/mole for chlorine–chlorine and bromine–bromine *gauche* interactions, but none for fluorine–fluorine. We shall adopt a correction of 1.0 kcal/mole for *gauche* interactions between chlorine, bromine, and iodine, and no correction for interactions between halogens and fluorine.¹⁹¹ Little evidence is available with respect to *cis* corrections in olefins. The *cis* isomer of 1,2-dichloroethylene is 0.4 kcal/mole more stable than the *trans*.¹⁹² We will assume no *cis* correction is necessary for halogens or for halogen–alkyl groups and make the usual assumption that *cis*-alkyl groups are 1.0 kcal/mole less stable than *trans*, including perhaloalkyl groups.¹⁹³ Ring strains are assumed to be the same as for the unsubstituted rings. This is probably in error for cyclopropane rings. *ortho* corrections are made as indicated by the data.

Data and Estimated Values. The group values for haloalkanes are summarized in Table XXXVII and a comparison with observed data, from which the groups were obtained, is made in Table XXIV. The values for haloalkenes, -alkynes, and -arenes are given in Table XXXVII and compared with experiment in Table XXV. The heats of formation of all fluorine-substituted compounds have been corrected to conform with $\Delta H_f^\circ_{298}(\text{HF,g}) = -64.8$ kcal/mole,¹³³ and extensive use has been made of Stull's compilation of vapor pressure data¹⁹⁴ to determine $\Delta H_v^\circ(25^\circ)$, when necessary.

Although the data for haloalkanes is somewhat sparse, the estimates agree to ± 2 kcal/mole in ΔH_f° , ± 1 gibbs/mole in S°_{298} , and ± 0.5 gibbs/mole in C_p° . Kolesov, *et al.*,¹⁹⁵ have reported $\Delta H_f^\circ_{298}(\text{CF}_3\text{CH}_3) = -175.9$ kcal/mole, 8 kcal/mole more stable than predicted by groups; however, they experienced incomplete combustion ($\sim 95\%$) and only analyzed for HF and CO_2 . Formation of $\text{CF}_2=\text{CH}_2$, CO , CH_2F_2 , etc., would all lead to low values of ΔH_f° , so that their results are not included. Rodgers¹⁹⁶ has suggested that nearest-neighbor interactions only may prove insufficient to adequately predict ΔH_f° of mixed fluorocarbon–hydrocarbon compounds, and there appears adequate evidence of this in haloalkenes. Thus, $\text{CF}_2=\text{CH}_2$ and $\text{CH}_2=\text{CCl}_2$ are 9.3 and 3.9 kcal/mole more stable than the estimates from group contributions (Table XXV). Compounds composed of groups of widely different polarity may show deviations resulting from dipole–dipole interactions. Such an effect might be expected to stabilize $\text{CF}_3\text{—CH}_3$ also. Additional evidence of discrepancies in the alkene groups comes from the equilibrium data of Schlag and Peatman¹⁹⁷ (1) and Chesick¹⁹⁸ (2):

- (183) B. G. Gowenlock and K. E. Thomas, *J. Chem. Soc., B*, 409 (1966).
 (184) S. W. Benson, "Techniques of Thermochemical Kinetics," John Wiley and Sons, Inc., New York, N. Y., 1968.
 (185) T. N. Sarachman, *J. Chem. Phys.*, **39**, 469 (1963).
 (186) C. Komaki, I. Ichishima, K. Kuratani, T. Miyazawa, T. Shumanouchi, and S. Mizushima, *Bull. Chem. Soc. Japan*, **28**, 330 (1955).
 (187) E. Hirota, *J. Chem. Phys.*, **37**, 283 (1962).

- (188) S. Mizushima "Structure of Molecules and Internal Rotation," Academic Press, New York, N. Y., 1954, p 41.
 (189) G. Allen, P. N. Brier, and G. Lane, *Trans. Faraday Soc.*, **63**, 824 (1967).
 (190) P. Klaboe and J. R. Nielsen, *J. Chem. Phys.*, **33**, 1764 (1960).
 (191) The data are somewhat mixed on this point. See: (a) R. E. Kagarise, *ibid.*, **29**, 680 (1958); (b) P. Klaboe and J. R. Nielsen, *ibid.*, **32**, 899 (1960).
 (192) K. S. Pitzer and J. L. Hollenberg, *J. Am. Chem. Soc.*, **76**, 1493 (1954).
 (193) E. W. Schlag and E. W. Kaiser, Jr., *ibid.*, **87**, 1171 (1965).
 (194) D. R. Stull, *Ind. Eng. Chem.*, **39**, 517 (1949).
 (195) V. P. Kolesov, A. M. Martynov, and S. M. Skurator, *Russ. J. Phys. Chem.*, **39**, 223 (1965).
 (196) A. S. Rodgers, *J. Phys. Chem.*, **71**, 1996 (1967).
 (197) E. W. Schlag and W. B. Peatman, *J. Am. Chem. Soc.*, **86**, 1676 (1964).
 (198) J. P. Chesick, *ibid.*, **88**, 4800 (1966).


Table XXIV: Haloalkanes

Compound	ΔH_f°		Δ	Ref	n, σ	S°		Δ	Ref	C_p^{300}		Δ	C_p^{800}		Δ	Ref
	Obsd	Estid				Obsd	Estid			Obsd	Estid		Obsd	Estid		
CF ₃ CCl ₃	...				3 ²	86.4	88.5	-2.0	199	28.1	29.0	-0.9	39.6	39.9	-0.3	199
CF ₂ ClCF ₂ Cl	-212.6	-212.6	[0] ^a	196	2	...				28.9	27.4	+1.5	...			200
CF ₃ CF ₃	-316.8	-316.8	0.0	133	3 ² × 2	79.4	79.3	0.1	201	25.5	25.4	0.1	38.3	38.6	0.3	201
CCl ₃ CCl ₃	-35.3	-35.4	[0]	196	3 ² × 2	94.8	95.1	0.3	202	32.7	32.6	0.1	41.3	41.2	0.1	202
CBBr ₂ CBBr ₂	...				3 ² × 2	105.6	105.6	[0]	202	33.4	33.4	[0]	39.9	39.8	[0]	202
CF ₃ CHClBr	...				2 × 3	87.1	87.1	[0]	203	25.1	25.1	[0]	37.2	37.2	[0]	203
CF ₃ CHCl ₂	...				3	84.1	84.0	0.1	199	24.4	24.8	-0.4	37.3	37.6	-0.3	199
CF ₃ CH ₂ Cl	...				3	78.2	78.1	0.1	204	21.4	21.6	-0.2	34.8	34.6	-0.2	204
CF ₃ CH ₂ Br	...				3	80.7	81.1	-0.4	204	21.8	21.8	0.0	35.1	34.8	0.3	204
CF ₃ CH ₂ I	...				3	82.5	82.8	-0.3	204	21.9	21.9	[0]	35.1	35.1	[0]	204
CF ₃ CH ₂ F	...				3	76.0	75.7	0.3	204	20.9	20.8	0.1	34.5	34.5	0.0	204
CF ₃ CH ₃	...				3 ²	68.5	68.5	0.0	205	18.9	18.9	0.0		206
CHF ₂ CH ₂ F	(-159.5)	-161.1	+1.6	196	1	80.6	81.5	-0.9	208	21.3	21.0	0.3	33.3	33.2	0.1	208
CHCl ₂ CH ₂ Cl	...				3 ²	77.2	76.4	0.8	209	22.4	22.5	-0.1	33.3	33.6	-0.3	209
CH ₂ CCl ₂	...				3	67.3	67.3	[0]	210	16.1	16.1	[0]		210
CH ₃ CHF ₂	(-121)	-119.4	-1.6	196	3	72.9	71.9	1.0	212	18.2	18.3	-0.1	30.9	30.9	0.0	212
CH ₃ CHCl ₂	(-31)	-29	[0] ^c	211	3		
CH ₂ ClCH ₂ Cl	-31.1	-31.4	0.3	213, 214	3		
CH ₃ CH ₂ F	...				3	63.3	63.6	-0.3	215	14.1	14.3	-0.2		215
CH ₃ CH ₂ Cl	-25.7	-25.8	0.1	216	3	66.1	66.0	0.1	217	15.1	15.1	0.0	28.3	28.3	0.0	217
CH ₃ CH ₂ Br	-14.8	-15.6	0.8	216	3	68.6	69.0	-0.4	217	15.5	15.3	0.2	28.6	28.5	0.1	217
CF ₃ CF ₂ CF ₂ Cl	...				3 ²	97.3	97.4	[0]	218	37.4	37.3	[0]		218
CF ₃ CF ₂ CH ₃	...				3	86.3	86.3	[0]	218	28.8	28.8	[0]		218
CH ₃ CH ₂ CH ₂ F	-66.8	-66.8	[0]	207	3		218
CH ₃ CHFCH ₃	-68.6	-68.6	[0]	207	3		218
CH ₃ CH ₂ CH ₂ Cl	-31.2	-30.9	-0.3	216	3	76.4	75.4	1.0	217	20.3	20.6	-0.3	39.2	39.4	-0.2	217
CH ₃ CHClCH ₃	-32.9	-33.0	[0]	216	3	74.1	74.1	[0]	219	21.3	21.3	[0]		217
CH ₃ CCl ₂ CH ₃	(-41.7)	-39.7	[0] ^c	211	3		219
CH ₃ CH ₂ CH ₂ Br	-19.9	-20.6	0.7	216	3	79.2	78.4	0.8	217	20.7	20.8	-0.1	39.4	39.6	-0.2	217
CH ₃ CHBrCH ₃	-22.7	-23.6	0.9	216	3		
CH ₃ CH ₂ CHBrCH ₂ Br	-24.4	-23.8	-0.6	220	3		
(CH ₃) ₃ CCl	-43.0	-43.0	[0]	219	3 ⁴	77.0	77.0	[0]	219	27.9	27.9	[0]		219
CH ₃ CH ₂ CHBrCH ₃	-28.7	-28.6	-0.1	220, 221	3		
CH ₃ CH ₂ CH ₂ CH ₂ Br	-31.0	-30.5	-0.5	220	3		
CH ₃ CH ₂ CH ₂ CH ₂ CH ₂ Br	-801.9	-801.8	0.1	222	3		
<i>n</i> -C ₇ F ₁₆	-687.3	-687.3 ^d	0.0	222	3		
CF ₃ CCl ₂ F ₁₁	-783.5	-783.5 ^d	0.0	222	3		
CF ₃ CF ₂ CCl ₂ F ₁₁	-26.3	-26.6 ^d	0.3	220	3		
1,2-Br ₂ CCl ₂ H ₁₀	-35.9	-35.5	-0.4	220	3		
<i>n</i> -C ₈ H ₁₆ Br	-40.7	-40.5	-0.2	220	3		
<i>n</i> -C ₇ H ₁₄ Br	-46.3	-45.5	-0.8	220	3		
<i>n</i> -C ₈ H ₁₇ Br	15.9	15.9	[0]	224	2	83.3	83.6	-0.3	224	19.2	18.4	0.8		224
CH ₂ ICH ₂ I	10.6	10.95	-0.35	224	2		
CH ₂ ICH ₂ I	8.6	8.6	[0]	224	2,3	94.3	94.3	[0]	224	(24.1)	24.1	[0]		224
CH ₃ CHICH ₂ I	-17.2	-17.2	[0]	225	3 ⁴	82.5	82.5	[0]	225		225
(CH ₂) ₂ Cl	-52.0	-52.0	[0]	223	3		
(CH ₂) ₂ CHCH ₂ CH(NF ₂)CH ₂ (NF ₂)	-52.0	-52.0	[0]	223	3		


^a [0] indicates group value determined by these data. ^b Parentheses indicate questionable data. ^c ΔH_f° 's from ref 211 are generally 2 ± 2 kcal/mole too stable, and group values have been correspondingly adjusted. ^d Calculated assuming strain-free ring.

Table XXV: Haloalkenes, Alkynes, and Arenes

Compound	n, σ	R_{e_j}	ΔH_f°		Δ	S°		Δ	$C_{p,300}^\circ$		Δ	$C_{p,800}^\circ$		Δ	Ref
			Obsd	Estd		Obsd	Estd		Obsd	Estd		Obsd	Estd		
$\text{CFCl}=\text{CCl}_2$	1		81.7	81.9	-0.2	21.8	21.7	0.1	28.8	28.8	0.0	28.8	28.8	0.0	226
$\text{CF}_2=\text{CClBr}$	1		82.4	82.4	[0]	21.8	21.8	[0]	28.5	28.5	[0]	28.5	28.5	[0]	227
$\text{CFCl}=\text{CFCl}$	2		78.2	78.2	0.0	21.0	20.6	0.4	28.4	28.4	0.0	28.4	28.4	0.0	226
$\text{CF}_2=\text{CBr}_2$	2		87.0	83.5	3.5	22.2	22.0	0.2
$\text{CF}_2=\text{CFCl}$	1		77.0	77.1	-0.1	20.0	20.0	0	28.0	28.0	0.0	28.0	28.0	0.0	229
$\text{CF}_2=\text{CFBr}$	1		79.8	79.8	[0]	20.5	20.5	[0]	28.1	28.1	[0]	28.1	28.1	[0]	230
$\text{CF}_2=\text{CF}_2$	2 ²	231	71.7	71.8	-0.1	19.3	19.4	-0.1	27.6	27.6	0.0	27.6	27.6	0.0	231
$\text{CCl}_2=\text{CCl}_2$	2 ²	196	81.3	81.4	-0.1	22.7	22.8	-0.1	29.2	29.2	0.0	29.2	29.2	0.0	232
$\text{CBr}_2=\text{CBr}_2$	2 ²		92.4	92.4	0.0	24.5	24.6	-0.1	29.7	29.8	-0.1	29.7	29.8	-0.1	232
$\text{CHF}=\text{CCl}$	1		75.1	74.9	0.2	18.3	18.2	0.1	26.5	26.4	0.1	26.5	26.4	0.1	233
$\text{CF}_2=\text{CHBr}$	1		75.2	75.6	-0.4	17.6	17.8	-0.2	26.2	26.2	0.0	26.2	26.2	0.0	234
$\text{CF}_2=\text{CHF}$	1		69.9	70.1	-0.2	16.5	16.5	0.0	25.6	25.6	0.0	25.6	25.6	0.0	236
$\text{CHCl}=\text{CCl}_2$	1	235	77.6	77.5	0.1	19.3	19.3	0.0	26.9	26.9	0.0	26.9	26.9	0.0	237
$\text{CHBr}=\text{CBr}_2$	1		85.7	85.9	-0.2	20.5	20.4	0.1	238
$\text{CH}_2=\text{CFCl}$	1		67.6	67.4	0.2	15.3	15.4	-0.1	24.4	24.3	0.1	24.4	24.3	0.1	239
$\text{CH}_2=\text{CF}_2$	1	
$\text{CH}_2=\text{CCl}_2$	2	235	69.0	68.3	0.7	16.1	16.5	0.4	24.7	24.7	0.0	24.7	24.7	0.0	241
$\text{CHCl}=\text{CHCl}$ (av)	2	240	69.3	69.4	-0.1	15.8	15.8	0.0	24.6	24.6	0.0	24.6	24.6	0.0	192
$\text{CHBr}=\text{CHBr}$ <i>cis</i>	2		75.8	75.2	0.6	16.5	16.2	0.3	238
$\text{CH}_2=\text{CHCl}$	1	242	63.2	63.0	0.2	12.9	13.0	-0.1	22.4	22.4	0.0	22.4	22.4	0.0	243, 244
$\text{CH}_2=\text{CHBr}$	1	245	66.0	65.9	0.1	13.4	13.2	0.2	22.5	22.5	0.0	22.5	22.5	0.0	243, 244
$\text{CH}_2=\text{CHI}$	1		68.1	68.1	[0]	13.9	13.9	[0]	22.7	22.7	[0]	22.7	22.7	[0]	244
$\text{CH}_3\text{C}\equiv\text{CCI}$	3		68.0	68.0	[0]	17.2	17.2	[0]	27.0	27.0	[0]	27.0	27.0	[0]	246
$\text{CH}_3\text{C}\equiv\text{CBr}$	3		70.7	70.7	[0]	17.6	17.6	[0]	27.1	27.1	[0]	27.1	27.1	[0]	246
$\text{CH}_3\text{C}\equiv\text{CI}$	3		72.4	72.4	[0]	17.8	17.8	[0]	27.2	27.2	[0]	27.2	27.2	[0]	246
$\text{C}_6\text{H}_6\text{F}$	2	247	72.4	72.2	0.4	22.7	22.5	0.2	47.6	47.5	0.1	47.6	47.5	0.1	248
<i>o</i> - $\text{C}_6\text{H}_4\text{F}_2$	2	249	77.3	77.0	0.3	25.6	25.6	0.0	49.5	49.8	-0.3	49.5	49.8	-0.3	251
<i>m</i> - $\text{C}_6\text{H}_4\text{F}_2$	2	249	76.7	77.0	-0.3	25.5	25.6	-0.1	49.7	49.8	-0.1	49.7	49.8	-0.1	251
<i>p</i> - $\text{C}_6\text{H}_4\text{F}_2$	2 ²	249	75.6	75.6	0.0	25.7	25.6	0.1	49.7	49.8	-0.1	49.7	49.8	-0.1	251
<i>p</i> - $\text{FC}_6\text{H}_4\text{CH}_3$	3 × 2	252	81.3	81.4	-0.1	27.9	28.2	-0.3	58.6	58.1	0.5	58.6	58.1	0.5	252
$\text{C}_6\text{H}_5\text{CF}_3$	3 × 2	253	89.2	89.2	[0]	31.4	31.4	[0]	62.8	62.8	[0]	62.8	62.8	[0]	253
4,4'- $\text{FC}_6\text{H}_4\text{C}_6\text{H}_4\text{F}$	2	254
2,2'- $\text{FC}_6\text{H}_4\text{C}_6\text{H}_4\text{F}$	2	254
$\text{C}_6\text{H}_5\text{Cl}$	2	255	75.2	75.2	[0]	23.6	23.6	[0]	47.9	47.9	[0]	47.9	47.9	[0]	248
<i>o</i> - $\text{C}_6\text{H}_4\text{Cl}_2$	2	255	75.2	75.2	[0]	23.6	23.6	[0]	47.9	47.9	[0]	47.9	47.9	[0]	248
<i>m</i> - $\text{C}_6\text{H}_4\text{Cl}_2$	2	255	75.2	75.2	[0]	23.6	23.6	[0]	47.9	47.9	[0]	47.9	47.9	[0]	248
<i>p</i> - $\text{C}_6\text{H}_4\text{Cl}_2$	2	255	75.2	75.2	[0]	23.6	23.6	[0]	47.9	47.9	[0]	47.9	47.9	[0]	248
<i>p</i> - $\text{ClC}_6\text{H}_4\text{Et}$	2	255	75.2	75.2	[0]	23.6	23.6	[0]	47.9	47.9	[0]	47.9	47.9	[0]	248
<i>p</i> - $\text{ClC}_6\text{H}_4\text{Et}$	2	255	75.2	75.2	[0]	23.6	23.6	[0]	47.9	47.9	[0]	47.9	47.9	[0]	248
2,5'- $\text{Cl}_2\text{C}_6\text{H}_3\text{CH}=\text{CH}_2$	2	240	21.6	21.6	-0.2	240	240	-0.2	240	240	-0.2	240	240	-0.2	240
2,2'- $\text{ClC}_6\text{H}_4\text{C}_6\text{H}_4\text{Cl}$	2	254	30.4	30.5	-0.1	254	254	-0.1	254	254	-0.1	254	254	-0.1	254
4,4'- $\text{ClC}_6\text{H}_4\text{C}_6\text{H}_4\text{Cl}$	2	254	28.8	29.3	-0.5	254	254	-0.5	254	254	-0.5	254	254	-0.5	254
$\text{C}_6\text{H}_5\text{Br}$	2	256	27.2	27.2	[0]	27.2	27.2	[0]	27.2	27.2	[0]	27.2	27.2	[0]	256
$\text{C}_6\text{H}_5\text{CH}_2\text{Br}$	2	257	15.1	15.1	[0]	15.1	15.1	[0]	15.1	15.1	[0]	15.1	15.1	[0]	257
$\text{C}_6\text{H}_5\text{I}$	2	258, 259	40.5	40.5	[0]	40.5	40.5	[0]	40.5	40.5	[0]	40.5	40.5	[0]	258, 259
$\text{C}_6\text{H}_5\text{CH}_2\text{I}$	2	260	30.4	30.4	[0]	30.4	30.4	[0]	30.4	30.4	[0]	30.4	30.4	[0]	260
C_6F_6	2	261	-224.0	-226.8	2.8	-224.0	-226.8	2.8	-224.0	-226.8	2.8	-224.0	-226.8	2.8	261
C_6Cl_6	2	240	-8.3	-9.6	1.3	-8.3	-9.6	1.3	-8.3	-9.6	1.3	-8.3	-9.6	1.3	240



$$\begin{aligned}
 \Delta H_{298}^\circ &= 11.7 \text{ kcal/mole} \\
 \Delta S_{298}^\circ &= 9.6 \text{ gibbs/mole}
 \end{aligned}$$



$$\begin{aligned}
 \Delta H_{298}^\circ &= 0.4 \text{ kcal/mole} \\
 \Delta S_{298}^\circ &= 4.9 \text{ gibbs/mole}
 \end{aligned}$$

Writing equilibria 1 and 2 into group notation and subtracting yields

$$[\Delta(1) - \Delta(2)]/2 = [C_d(C_d)(F) - C_d(C)(F)] - [C_d(C_d)(C) - C_d(C)_2] \quad (3)$$

where we have assumed identical ring strains. Both quantities in brackets on the right-hand side (rhs) of eq 3 are expected to be approximately equal, *i.e.*, $\text{rhs}(3) \cong 0$. This is true for hydrocarbons (H replaces F in eq 3 for which $\text{rhs}(3) = -0.15$ kcal/mole from enthalpies and $+0.3$ gibbs/mole from entropies. Equilibria 1 and 2, however, yield $\text{rhs}(3) = 5.2$ kcal/mole and 2.35 gibbs/mole for enthalpies and entropies, respectively. These large differences are hard to rationalize with group additivity.

As a result of the above-mentioned contradictions in group additivity for haloalkenes, we feel that the group contributions to ΔH_{298}° for these compounds should be used with caution, particularly when combining groups with large differences in polarity. Estimates of S_{298}° and C_p° , on the other hand, are seen in Table XXV to be accurate to ± 1.0 and ± 0.5 gibbs/mole, respectively.

The estimates of ΔH_f° , S_{298}° , and C_p° for haloarenes are found accurate to ± 2.0 kcal/mole, ± 1.0 gibbs/mole, and ± 0.5 gibbs/mole, respectively.

This section prepared by S. W. Benson and A. S. Rodgers.

- (199) N. W. Luft, *J. Chem. Phys.*, **22**, 153 (1954).
 (200) J. J. Martin, *J. Chem. Eng. Data*, **5**, 334 (1960).
 (201) Reference 135, Sept 1965.
 (202) R. A. Carney, E. A. Piotrowski, A. G. Meister, J. H. Braun, and F. F. Cleveland, *J. Mol. Spectry.*, **7**, 209 (1961).
 (203) R. Theimer and J. R. Nielsen, *J. Chem. Phys.*, **27**, 887 (1957).
 (204) C. R. Ward and C. H. Ward, *J. Mol. Spectry.*, **12**, 289 (1964).
 (205) J. R. Nielsen, H. H. Claasen, and D. C. Smith, *J. Chem. Phys.*, **18**, 1471 (1950).
 (206) H. W. Thompson and R. B. Temple, *J. Chem. Soc.*, 1428 (1948).
 (207) J. R. Lacher, A. Kianpour, and J. D. Park, *J. Phys. Chem.*, **60**, 1454 (1956).
 (208) R. H. Harrison and K. A. Kobe, *J. Chem. Phys.*, **26**, 1411 (1957).
 (209) K. S. Pitzer and J. L. Hollenberg, *J. Am. Chem. Soc.*, **75**, 2219 (1953).
 (210) D. C. Smith, R. A. Saunders, J. R. Nielsen, and E. E. Ferguson, *J. Chem. Phys.*, **20**, 847 (1952).
 (211) L. Smith, L. Bjellerup, S. Krook, and H. Westermark, *Acta Chem. Scand.*, **7**, 65 (1953).
 (212) J. C. M. Li and K. S. Pitzer, *J. Am. Chem. Soc.*, **78**, 1077 (1956).
 (213) J. B. Conn, G. B. Kistiakowsky, and E. A. Smith, *ibid.*, **60**, 2764 (1938).
 (214) G. C. Sinke and D. R. Stull, *J. Phys. Chem.*, **62**, 397 (1958).
 (215) E. Catalano and K. S. Pitzer, *ibid.*, **62**, 873 (1958).
 (216) J. Davies, J. R. Lacher, and J. D. Park, *Trans. Faraday Soc.*, **61**, 2413 (1965).
 (217) J. H. S. Green and D. J. Holden, *J. Chem. Soc.*, 1794 (1962).
 (218) C. L. Angell, *J. Chem. Eng. Data*, **9**, 341 (1964).
 (219) K. E. Howlett, *J. Chem. Soc.*, 1784 (1955).
 (220) L. Bjellerup, *Acta Chem. Scand.*, **15**, 121 (1961).

VII. Sulfur, Boron, Phosphorus Compounds and Organometallic Compounds

This section extends the group additivity rules to include organometallic (Sn, Hg, Zn, Cd, Ge, Al, Pb, Ti, V, and Cr) and organononmetallic (S, B, P) compounds. Table XXXVIII lists the best group values and ring corrections. The group values in the table are based on the most stable conformation. We found *gauche* corrections only necessary for the organoboron compounds. In fact, organosulfur compounds indicate no *gauche* interactions. This suggests that these conformational interactions are only important for the first-row elements.

- (221) J. R. Lacher, T. J. Billings, D. E. Campion, K. R. Lea, and J. D. Park, *J. Am. Chem. Soc.*, **74**, 5291 (1952).
 (222) W. D. Good, D. R. Douslin, D. W. Scott, A. George, J. L. Lacina, J. P. Dawson, and G. Waddington, *J. Phys. Chem.*, **63**, 1133 (1959).
 (223) W. D. Good, D. R. Douslin, and J. P. McCullough, *ibid.*, **67**, 1312 (1963).
 (224) S. W. Benson and A. Amano, *J. Chem. Phys.*, **36**, 3464 (1962).
 (225) S. W. Benson and A. Amano, *ibid.*, **37**, 197 (1962).
 (226) D. E. Mann and E. K. Plyler, *ibid.*, **23**, 1989 (1955).
 (227) J. R. Nielsen and R. Theimer, *ibid.*, **30**, 98 (1959).
 (228) J. R. Nielsen and R. Theimer, *ibid.*, **30**, 103 (1959).
 (229) D. E. Mann, N. Acquista, and E. K. Plyler, *ibid.*, **21**, 1949 (1953).
 (230) D. E. Mann, N. Acquista, and E. K. Plyler, *ibid.*, **22**, 1199 (1954).
 (231) Reference 133, Sept 1964.
 (232) D. E. Mann, J. H. Meal, and E. K. Plyler, *J. Chem. Phys.*, **24**, 1018 (1956).
 (233) C. J. Muellemann, K. Ramaswamy, F. F. Cleveland, and S. Sundaram, *J. Mol. Spectry.*, **11**, 262 (1963).
 (234) R. Theimer and J. R. Nielsen, *J. Chem. Phys.*, **27**, 264 (1957).
 (235) V. P. Kolesov, A. M. Martynov, S. M. Shetekher, and S. M. Skuratov, *Russ. J. Phys. Chem.*, **36**, 1118 (1962).
 (236) D. E. Mann, N. Acquista, and E. K. Plyler, *J. Chem. Phys.*, **22**, 1586 (1954).
 (237) G. Allen and H. T. Berstein, *Can. J. Chem.*, **32**, 1044 (1954).
 (238) J. C. Evans and H. J. Berstein, *ibid.*, **33**, 1171 (1955).
 (239) D. E. Mann, N. Acquista, and E. K. Plyler, *J. Chem. Phys.*, **23**, 2122 (1955).
 (240) G. C. Sinke and D. R. Stull, *J. Phys. Chem.*, **62**, 397 (1958).
 (241) D. L. Hildebrand, R. A. McDonald, W. R. Kramer, and D. R. Stull, *J. Chem. Phys.*, **30**, 930 (1959).
 (242) J. R. Lacher, H. B. Gottlieb, and J. D. Park, *Trans. Faraday Soc.*, **58**, 2348 (1962).
 (243) C. W. Gullikson and J. R. Nielsen, *J. Mol. Spectry.*, **1**, 155 (1957).
 (244) R. E. Richards, *J. Chem. Soc.*, 1931 (1948).
 (245) J. R. Lacher, A. Kianpour, P. Montgomery, H. Knedler, and J. D. Park, *J. Phys. Chem.*, **61**, 1125 (1957).
 (246) P. N. Daykin, S. Sundaram, and F. F. Cleveland, *J. Chem. Phys.*, **37**, 1087 (1962).
 (247) W. D. Good and D. W. Scott, *Pure Appl. Chem.*, **2**, 77 (1961).
 (248) D. H. Whiffen, *J. Chem. Soc.*, 1350 (1956).
 (249) W. D. Good, J. L. Lacina, D. W. Scott, and J. P. McCullough, *J. Phys. Chem.*, **66**, 1529 (1962).
 (250) D. W. Scott, J. F. Messerly, S. S. Todd, I. A. Hossenlopp, A. Osborn, and J. P. McCullough, *J. Chem. Phys.*, **38**, 532 (1963).
 (251) J. H. S. Green, W. Kynaston, and H. M. Paisley, *J. Chem. Soc.*, 473 (1963).
 (252) D. W. Scott, J. F. Messerly, S. S. Todd, I. A. Hossenlopp, D. R. Douslin, and J. P. McCullough, *J. Chem. Phys.*, **37**, 867 (1962).
 (253) D. W. Scott, D. R. Douslin, J. F. Messerly, S. S. Todd, I. A. Hossenlopp, T. C. Kincheloe, and J. P. McCullough, *J. Am. Chem. Soc.*, **81**, 1015 (1959).
 (254) N. K. Smith, G. Gorin, W. D. Good, and J. P. McCullough, *J. Phys. Chem.*, **68**, 940 (1964).
 (255) W. N. Hubbard, J. W. Knowlton, and H. M. Huffman, *ibid.*, **58**, 396 (1954).
 (256) K. Hartley, H. Pritchard, and H. Skinner, *Trans. Faraday Soc.*, **47**, 254 (1951), combined with $\Delta H_f^\circ(\text{C}_6\text{H}_6)_{\text{liq}}$.
 (257) S. J. Ashcroft, A. S. Carson, and J. B. Pedley, *ibid.*, **59**, 2713 (1963).
 (258) L. Smith, *Acta Chem. Scand.*, **10**, 884 (1956).
 (259) L. S. Carson, E. M. Carson, and B. Wilmshirst, *Nature*, **170**, 320 (1952).
 (260) R. Walsh, D. M. Golden, and S. W. Benson, *J. Am. Chem. Soc.*, **88**, 650 (1966).
 (261) J. D. Cox, H. A. Gundry, and A. J. Head, *Trans. Faraday Soc.*, **60**, 653 (1964).

This reduction in next-nearest-neighbor interaction is coincident with a 20–30% increase in the bonding distance. Bond lengths in ångströms for the first- and second-row elements are, respectively, C–B, 1.56; C–C, 1.54; C–N, 1.47; C–O, 1.47; C–F, 1.38; and C–Al, 1.92; C–Si, 1.87; C–P, 1.87; C–S, 1.82; C–Cl, 1.77.

The utilization of these group values is not restricted solely to the obvious predicting of thermochemical properties of unmeasured compounds. The concept of group additivity of molecular properties allows a check of the internal consistency of the measured thermochemical data of related compounds. This is a plausible test of the reliability of the experimental thermochemical data. As an example, the inconsistencies in the gaseous heat of formation of organosilicon compounds are made apparent by the failure to construct a reproducible set of group values. This inconsistency is evident in the values of the groups $[C-(Si)(C)(H)_2] - [C-(Si)(H)_3]$ and $[Si-(C)(H)_3] + [C-(Si)(C)(H)_2]$. The former group has the values: +18.08 kcal/mole in $Si(CH_3)_4 - Si(C_2H_5)_4$; +17.08 kcal/mole in $SiH(CH_3)_3 - SiH(C_2H_5)_3$; +12.58 kcal/mole in $SiH_2(CH_3)_2 - SiH_2(C_2H_5)_2$; +13.90 kcal/mole in $SiH_3CH_3 - SiH_3C_2H_5$; and the latter group has the values: -17.92 kcal/mole in $C_2H_5SiH_3$; -8.94 kcal/mole in $(CH_3)_2CHCH_2SiH_3$; -11.02 kcal/mole in $CH_3(CH_2)_3SiH_3$. Similarly, a correlation of the heat of formation of $(CH_3)_nM$ and $(C_2H_5)_nM$ and $(C_2H_5)_4M$ compounds, where $M = Si, Ge, Sn, Pb$, by Telnoi and Rabinovitch²⁶² demonstrates that the heat of formation of tetramethylsilicon is incorrect.^{262a} O'Neal and King²⁶³ found cases where the reported thermochemical data of the organosilicon compounds deviated appreciably from the less stringent bond additivity rule. They concluded that large errors in the heat of combustion are not unusual, and additional thermochemical data of greater accuracy are needed.

A. ORGANOSULFUR COMPOUNDS

1. Alkylthiols

The uncertainties in the predicted ΔH_f° , S° , and C_p° are ± 0.2 kcal/mole, ± 0.2 eu, and ± 0.1 gibbs/mole, respectively. The uncertainty in the heat capacity at high temperatures (800°K) increases to ± 0.5 gibbs/mole.

2. Thiaalkanes

The uncertainty in the predicted ΔH_f° , S° , and C_p° are ± 0.3 kcal/mole, ± 0.2 eu, and ± 0.3 gibbs/mole, respectively. The uncertainty in the heat capacity at high temperatures (800°K) increases to ± 0.5 gibbs/mole. The disagreement between calculated and observed values of S° and C_p° for the compound $(CH_3)_2CHSCH(CH_3)_2$ (see Table XXVI^{264–270})

(262) V. I. Telnoi and I. B. Rabinovich, *Russ. J. Phys. Chem.*, **39**, 1108 (1965).

(262a) Expressing the heat of formation of $(CH_3)_nM$ and $(CH_3CH_2)_nM$ in terms of their groups and combining, we find the relation $\Delta H_f^\circ[(CH_3CH_2)_nM] = \Delta H_f^\circ[(CH_3)_nM] + 4\{[C-(M)(C)(H)_2] - [C-(M)(H)_3] + [C-(C)(H)_3]\}$. If we invoke bond additivity rules, the linear correlation $\Delta H_f^\circ[(CH_3CH_2)_nM] = \Delta H_f^\circ[(CH_3)_nM] + \text{constant}$ is obtained. The error in this relation should not be larger than ± 5 kcal/mole (see ref 2).

(263) H. E. O'Neal and M. A. King, *Inorg. Chem.*, **5**, 435 (1966).

suggests that there is an error in the observed values.

3. Aromatic Sulfides

The uncertainty in the predicted ΔH_f° is ± 0.3 kcal/mole.

4. Thiaalkanes

The uncertainty in the predicted ΔH_f° is ± 0.2 kcal/mole.

5. Dithiaalkanes

The uncertainties in the predicted ΔH_f° , S° , and C_p° are ± 0.5 kcal/mole, ± 0.4 eu, and ± 0.2 gibbs/mole, respectively. At higher temperatures (800°K), the uncertainty in the heat capacity increases to ± 0.7 gibbs/mole.

6. Sulfoxides

The uncertainty in the predicted ΔH_f° is ± 1.0 kcal/mole. This is larger than the reported experimental precision, suggesting systematic errors.

7. Sulfones

The uncertainty in the predicted ΔH_f° is ± 2.0 kcal/mole for aliphatic, ± 2.8 kcal/mole for alkene, and ± 1.9 kcal/mole for aromatic sulfones. Consequently, the inconsistencies in the heats of formation are considerably larger than the reported experimental precision. The gas-phase heats of formation were derived from the standard heats by combining with the relevant heats of fusion, vaporization, and sublimation. In the case of the sulfones, these latter quantities have been estimated by empirical methods. It is quite possible that the inconsistencies in the group values arise from these crude estimates.

8. Vinyl Sulfones

The group additivity rule indicates that the reported heat of formation of *p*-tolylprop-1-ene sulfone (see Table XXVI) is inconsistent with the heats of formation of the other members of the homologous series.

9. Thiolacetic Acid

The uncertainty in the predicted ΔH_f° is ± 0.9 kcal/mole.

10. Thiophenes and Thiolenes

The uncertainties in the predicted ΔH_f° , S° , C_p° , and C_p° for the thiophenes are ± 0.1 kcal/mole, ± 0.8 eu, ± 0.4 gibbs/mole, and 0.6 gibbs/mole, respectively.

(264) H. Mackle and P. A. G. O'Hare, *Tetrahedron*, **19**, 961 (1963).

(265) D. W. Scott and J. M. McCullough, U. S. Bureau of Mines Bulletin 595, U. S. Government Printing Office, Washington, D. C., 1961.

(266) M. Mansson and S. Sunner, *Acta Chem. Scand.*, **16**, 1863 (1962).

(267) D. W. Scott and G. A. Crowder, *J. Chem. Phys.*, **46**, 1054 (1967).

(268) J. P. McCullough, H. L. Finke, W. N. Hubbard, S. S. Todd, J. F. Messerly, D. R. Douslin, and G. Waddington, *J. Phys. Chem.*, **65**, 784 (1961).

(269) W. D. Good and B. L. DePrater, *ibid.*, **70**, 3606 (1966).

(270) H. Mackle and R. T. B. McClean, *Trans. Faraday Soc.*, **59**, 669 (1963).

(271) H. Mackle, D. V. McNally, and W. Steel, Paper 9, ref 105 above.

(272) S. Sunner, *Acta Chem. Scand.*, **9**, 837 (1955).

(273) S. Sunner, *ibid.*, **9**, 847 (1955).

(274) G. B. Guthrie, Jr., D. W. Scott, and G. Waddington, *J. Am. Chem. Soc.*, **76**, 1488 (1954).

(275) S. Sunner, *Svensk. Chem. Tidskr.*, **58**, 71 (1946).

(276) R. E. Dinny and E. L. Pace, *J. Chem. Phys.*, **32**, 805 (1960).

(277) A. P. Claydon and C. T. Mortimer, *J. Chem. Soc.*, 3212 (1962).

(278) J. V. Davies and S. Sunner, *Acta Chem. Scand.*, **16**, 1870 (1962).

(279) D. W. Scott, *J. Chem. Eng. Data*, **4**, 246 (1959).

Table XXVI
Organosulfur Compounds



Compound	$\Delta H_{f, 298}^{\circ}$		Δ	Ref	n, σ	S_{388}°		Δ	Ref	$C_{p, 300}^{\circ}$		Δ	$C_{p, 800}^{\circ}$	Δ	Ref	
	Obsd	Estd				Obsd	Estd			Obsd	Estd					
CH ₃ SH	-5.46	-5.46	[0.0]	264	3	60.96	60.96	[0.0]	265	12.05	12.05	[0.0]	20.32	20.32	[0.0]	265
CH ₃ CH ₂ SH	-5.27		+0.19	265	3	70.77	70.84	+0.07	265	17.44	17.43	+0.01	31.83	32.58	-0.75	265
	-10.99	-11.11	+0.12	264												
	-10.87		+0.24	265												
CH ₃ (CH ₂) ₂ SH	-16.19	-16.06	-0.13	264	3	80.40	80.26	+0.14	265	22.75	22.93	-0.18	43.60	43.65	-0.05	265
	-16.06		[0.0]	265												
	-21.02	-21.01	-0.01	264												
CH ₃ (CH ₂) ₃ SH	-20.89		+0.12	265	3	89.38	89.68	[0.0]	265	28.37	28.43	-0.06	55.68	54.72	+0.96	265
	-25.88	-25.96	+0.08	264												
	-26.70		-0.74	269												
CH ₃ (CH ₂) ₄ SH	-25.94		+0.02	269	3	99.28	99.10	+0.18	295	33.91	33.93	-0.02	66.78	65.79	+0.99	268
	-30.82	-30.91	+0.09	269												
	-35.65	-35.86	+0.21	269												
CH ₃ (CH ₂) ₅ SH	-50.57	-50.71	+0.14	269	3 ²	77.51	77.88	-0.37	264	23.04	23.09	-0.05	43.26	43.24	+0.02	264
	-18.19	-18.18	-0.01	264												
	-23.21	-23.09	-0.12	264												
(CH ₃) ₂ CHCH ₂ SH	-23.12	-23.13	+0.01	264	3 ²	87.65	87.30	+0.35	267	28.64	28.59	+0.05	54.29	54.31	-0.02	264
	-23.12		+0.01	264												
	-26.14	-26.17	+0.03	264												
(CH ₃) ₃ C ₂ SH	-27.41	28.04	-0.63	264	3 ⁴	92.48	92.43	+0.05	264	34.46	34.50	-0.04	66.28	66.15	+0.13	264
	-30.34	-31.12	+0.78	264												
	-2.23	-2.06	-0.17	266												
HS(CH ₂) ₂ SH	-7.03	-7.01	-0.02	266
HS(CH ₂) ₃ SH	-12.03	-11.96	-0.07	266
HS(CH ₂) ₄ SH	-16.87	-16.91	+0.04	266
	-11.42	-11.52	+0.10	264	...	86.38	86.39	-0.01	264	25.94	25.21	+0.73	58.61	58.55	+0.06	264
	-22.90	-22.77	-0.13	264	...	87.37	87.31	+0.06	267	32.13	31.81	+0.32	75.19	73.37	+1.82	267
	-22.85		-0.08	294
	26.66	26.66	[0.0]	264	...	80.51	80.51	[0.0]	264	25.22	25.22	[0.0]	51.59	51.59	[0.0]	264
	21.9	21.90	[0.0]	264
	-8.94	-8.65	-0.29	264, 265	3 ² × 2	68.32	68.25	+0.07	264, 265	17.77	17.37	+0.40	31.59	31.45	+0.14	264, 265
CH ₃ CH ₂ S	-14.22	-14.30	+0.08	264, 265	3 ²	79.62	79.50	+0.12	264, 265	22.82	22.75	+0.07	42.93	43.71	+0.78	264, 265
CH ₃ S(CH ₂) ₂ CH ₃	-19.51	-19.25	-0.26	264, 265	3 ²	88.84	88.92	-0.08	264, 265	28.17	28.26	-0.09	54.45	54.78	-0.33	264, 265
(CH ₃) ₂ CHSCH ₃	-21.58	-21.37	-0.21	264, 265	3 ³	85.87	86.52	-0.65	264, 265	28.82	28.41	+0.41	54.89	54.37	-0.52	264, 265
CH ₃ S(CH ₂) ₃ CH ₃	-24.39	-24.20	-0.19	264, 265	3 ²	98.43	98.34	+0.09	264, 265	33.79	33.75	+0.04	66.53	65.85	+0.68	264, 265
CH ₃ SC(CH ₃) ₃	-24.20		[0.0]	264
	-28.93	-29.36	+0.43	264, 265	3 ³	89.21	89.49	-0.28	264, 265	34.66	34.22	+0.44	66.74	66.21	+0.53	264, 265
	-29.10	-29.15	+0.05	264, 265
CH ₃ S(CH ₂) ₄ CH ₃	-19.92	-19.95	+0.03	264, 265	3 ² × 2	87.96	88.01	-0.05	264, 265	28.09	28.13	-0.04	54.91	55.97	-1.06	264, 265

Table XXVII
Organophosphorus Compounds^a

Compound	$\Delta H_f^\circ_{298}$		Δ	S°_{298}		Δ	Ref	n, σ
	Obsd	Estd		Obsd	Estd			
(CH ₃) ₃ P	-23.2	-23.2	[0]					
	-23.0		+0.2					
(CH ₃ CH ₂) ₃ P	-32.8	-30.6	-2.2					
(C ₆ H ₅) ₃ P	72.4	72.4	[0]					
	76.0		+3.6					
(CH ₃ O) ₃ P	-165.0	-167.5	+2.0					
(CH ₃ CH ₂ O) ₃ P	-191.9	-193.0	+1.1					
((CH ₃) ₂ CHO) ₃ P	-221.8	-218.8	-3.0					
(CH ₃) ₃ PO	-103.0	-103.0	[0]					
(CH ₃ CH ₂ CH ₂) ₃ PO		-128.1						
(CH ₃ CH ₂ CH ₂) ₃ P	-44.9	-45.5	+0.6					
(CH ₃ CH ₂ CH ₂ CH ₂) ₃ PO		-141.1						
(CH ₃ CH ₂ CH ₂ CH ₂) ₃ P	-59.0	-58.5	-0.5					
(C ₆ H ₅) ₃ PO	3.5	3.5	[0]					
CH ₃ (PO)(OCH ₂ CH ₂) ₂	-228.1	-228.1	[0]					
CH ₃ CH ₂ (PO)(OCH(CH ₃) ₂) ₂	-248.7	-248.7	0					
(CH ₃ CH ₂ O) ₂ PO	-282.4	-282.4	[0]					
((CH ₃) ₂ CHO) ₃ PO	-308.2	-308.2	0					
CH ₃ (PO)Cl ₂	-133.9	-133.1	-0.8	81.2	81.2	[0]		3
CH ₃ (PO)F ₂				75.0	75.0	[0]		3
CH ₃ (PO)FCl				80.4	80.4	[0]		2.3
CH ₃ CH ₂ (PO)Cl ₂	-135.5	-136.5	+1.0					
CH ₃ (PO)(OH) ₂	-241.1	-239.6	-1.5					
CH ₃ CH ₂ (PO)(OH) ₂	-241.4	-243.0	+1.6					
CH ₃ CH ₂ PCl ₂	-62.6	-62.7	+0.1					
((CH ₃) ₂ CHO)(CH ₃)(PO)(OH)		-242.4						
((CH ₃) ₂ CHO)(CH ₃)(PO)(Cl)	-190.5	-190.5	[0]					
(CH ₃ CH ₂ O) ₂ (PO)OH		-288.2						
(CH ₃ CH ₂ O) ₂ (PO)F	-286.3	-286.3	[0]					
(CH ₃ CH ₂ O) ₂ (PO)O(PO)(OCH ₂ CH ₃) ₂	-500.8	-500.8	[0]					
(CH ₃ CH ₂ O) ₂ P(OH)	-209.7	-209.7	[0]					
((CH ₃ CH ₂) ₂ N) ₃ P	-49.4	-50.5	+1.1					
((CH ₃) ₂ N) ₃ P	-32.0	-30.7	-1.3					
((CH ₃) ₂ N) ₃ PO	-114.1	-111.7	-2.4					
((CH ₃ CH ₂) ₂ N) ₃ PO	-148.8	-151.3	-2.5					
(CH ₃) ₃ P:N(CH ₂ CH ₃)	-20.4	-20.4	[0]					
(C ₆ H ₅) ₃ P:N(CH ₂ CH ₃)	40.0	40.0	[0]					
(Cl ₂ P:N) ₃	-175.9	-174.9	-1.3					
(Cl ₂ P:N) ₄	-231.7	-233.2	+1.1					
((CH ₃) ₂ P:N) ₂	-107.1	-107.1	-0.1					
((C ₆ H ₁₁ O) ₂ P:N) ₃	-565.0	-564.9	-0.1					
(C ₆ H ₅) ₂ P:N) ₄	58.7	58.8	-0.1					

^a All data from ref 280.

((CH₃)₂CHO)₂BH, which justifies ascribing the disparity in (CH₃)₃B to experimental error.

D. ORGANOMETALLIC COMPOUNDS

The uncertainties in the predicted $\Delta H_f^\circ_{298}$ values (kcal/mole) are as follows: R₄Sn, ± 1.7 ; R₃SnX, ± 1.6 ; R₂SnX₂, ± 1.2 ; RSnX₃, ± 0.7 ; R₃SnH, ± 1.3 ; R₃Sn(CH=CH₂), ± 1.4 ; R₃SnSnR₃, ± 2.8 .

The group values clearly indicate that an inconsistency exists between the thermochemical data of (CH₃)₃SnSn(CH₃)₃ and (CH₃CH₂)₃SnSn(CH₂CH₃)₃, suggesting that the heats of formation of the ethyl compounds should be reduced by 16–18 kcal/mole (see Table XXIX^{281, 289–296}). This amounts to less

than 1% systematic error in the measured heats of combustion, which is entirely possible.

The gas-phase standard heats of formation of the compounds (CH₃CH₂CH₂CH₂)₃SnCl and (CH₂=CH)₄Sn were calculated from group values, which were combined with the observed heats of redistribution reaction to obtain the heat of formation of the related compounds, (CH₃CH₂CH₂CH₂)₃SnCl₃ and (CH₂=CH)₃SnCl.

1. Zinc

The uncertainty in the predicted $\Delta H_f^\circ_{298}$ of the tetraalkylzinc compounds is ± 2.5 kcal/mole.

(289) G. A. Nash, H. A. Skinner, and W. F. Stack, *Trans. Faraday Soc.*, **61**, 640 (1964).

(290) D. J. Coleman and H. A. Skinner, *ibid.*, **62**, 1721 (1966).

(291) W. F. Stack, G. A. Nash, and H. A. Skinner, *ibid.*, **61**, 2122 (1965).

(292) J. V. Davies, A. E. Pope, and H. A. Skinner, *ibid.*, **59**, 2233 (1963).

(293) D. C. Bradley and M. T. Hillyer, *ibid.*, **62**, 2367, 2374, 2382 (1966).

(294) A. E. Pope and H. A. Skinner, *ibid.*, **60**, 1402 (1964).

(295) K. Hartley, H. O. Pritchard, and H. A. Skinner, *ibid.*, **47**, 254 (1951); **46**, 1019 (1950).

(296) C. T. Mortimer, H. O. Pritchard, and H. A. Skinner, *ibid.*, **48**, 220 (1952).

Table XXVIII
Organoboron Compounds

Compound	(gauche)	$\Delta H_f^{\circ 298}$		Δ	Ref
		Obsd	Estd		
(CH ₃) ₃ B	(0)	-29.3	-29.3	[0]	281
(CH ₃ CH ₂) ₃ B	(0)	-36.5	-36.0	-0.5	281
(CH ₃ CH ₂ CH ₂ CH ₂) ₃ B	(0)	-67.7	-65.7	-2.0	281
((CH ₃) ₂ CHCH ₂) ₃ B	(3)	-66.9	-67.1	+0.2	281
((CH ₃ CH ₂)(CH ₃)CH) ₃ B	(9)	-58.7	-63.9	+5.2	281
(CH ₃ (CH ₂) ₄ CH ₂) ₃ B	(0)	-94.8	-95.4	+0.6	281
(CH ₃ (CH ₂) ₅ CH ₂) ₃ B	(0)	-109.4	-110.3	+0.9	281
(CH ₃ (CH ₂) ₆ CH ₂) ₃ B	(0)	-124.5	-125.1	+0.6	281
(CH ₃ CH ₂ CH ₂ CH ₂) ₂ BCl	(0)	-86.7	-86.7	[0]	281
(CH ₃ CH ₂ CH ₂ CH ₂) ₂ BBr	(0)	-71.3	-71.3	[0]	281
(CH ₃ CH ₂ CH ₂ CH ₂) ₂ BI	(0)	-53.3	-53.3	[0]	281
CH ₃ BF ₂	(0)	-199	-203	+4	281
CH ₃ CH ₂ BF ₂	(0)	-209	-205	+4	281
(CH ₃) ₂ CHBF ₂	(0)	-212	-212	0	281
CH ₂ =CHBF ₂	(0)	-171	-171	[0]	281
(CH ₃ O) ₂ BH	(0)	-139.2	-139.2	[0]	283
((CH ₃) ₂ CHO) ₂ BH	(0)	-174.4	-173.3	-1.1	282
(CH ₃ CH ₂ O) ₂ BCl	(0)	-197.0	-195.8	-1.2	284
(CH ₃ CH ₂ O)BCl ₂	(0)	-149.9	-149.2	-0.7	284
(CH ₃ CH ₂ O) ₃ B	(0)	-240.8	-239.6	-1.2	281
(CH ₃) ₃ BO ₃	(0)	-215.7	-215.7	[0]	281
(CH ₃ CH ₂) ₃ BO ₃	(0)	-240.8	-245.5	+4.7	281
(CH ₃ CH ₂ CH ₂) ₃ BO ₃	(0)	-260.9	-260.4	-0.5	281
(CH ₃ CH ₂ CH ₂ CH ₂) ₃ BO ₃	(0)	-279.4	-275.2	-4.2	281
(CH ₃ CH ₂ CH ₂ CH ₂) ₂ BOH	(0)	-130.6	-130.6	[0]	281
((CH ₃) ₂ N) ₃ B	(0)	-65.9	-65.9	[0]	286
((CH ₃) ₂ N) ₂ BCl	(0)	-85.3	-84.0	-1.3	285
(CH ₃) ₂ NBCl ₂	(0)	-99.1	-98.0	-1.1	285
(CH ₃ S) ₃ B	0	-37.4	-49.4	-12.0	288
(CH ₃ CH ₂ S) ₃ B	0	-68.2	-66.4	-1.8	288
(CH ₃ CH ₂ CH ₂ S) ₃ B	0	-80.0	-81.2	-1.2	288
(CH ₃ CH ₂ CH ₂ CH ₂ S) ₃ B	0	-94.2	-96.2	-2.0	288
(CH ₃ CH ₂ CH ₂ CH ₂ CH ₂ S) ₃ B	0	-110.7	-111.1	-0.4	288
(CS) ₃ B	0	45.1	45.1	[0]	288

^a Heat of vaporization corrected to 25°.

2. Organotitanium, -vanadium, and -chromium Compounds

The reported uncertainty in $\Delta H_f^{\circ 298}$ is about ± 2 kcal/mole. Group additivity suggests ± 9 kcal/mole for titanium alkoxides. *gauche* corrections were not employed. The deviation of the calculated heat of formation from the observed value could not be reconciled with next-nearest neighbor interactions. Apparently these errors are associated with the experimental values. A comparison of the steric interactions in the titanium alkyl oxides with the observed steric interactions in di-*t*-butyl ether indicates that the intramolecular strain must be nearly zero for these compounds.

In Table XXX it is immediately apparent that the heat of formation of [(CH₃CH₂)(CH₃)₂CO]₄Ti is in error by at least 16 kcal/mole. The other values fall into two sets: [CH₃CH₂O]₄Ti, [(CH₃)₂CHO]₄Ti, [CH₃(CH₃CH₂)CHO]₄Ti, [(CH₃)₃CO]₄Ti with an average value of -251 ± 2 kcal/mole and [CH₃(CH₂)₂O]₄Ti, [CH₃(CH)₂O]₄Ti, [CH₃(CH₂)₃O]₄Ti, [(CH₃)₂CHCH₂O]₄Ti with an average value of -263 ± 3 kcal/mole. This disparity probably originates from the estimates of the heats of vaporization and heats of depolymerization of the liquids, rather than the measured heats of formation of the liquids.

3. Mercury

The uncertainties in the predicted $\Delta H_f^{\circ 298}$ are listed as follows: R₂Hg, ± 2.7 kcal/mole; RHgX, ± 0.9 kcal/mole.

E. MISCELLANEOUS COMPOUNDS

Table XXXI²⁹⁷⁻³⁰⁵ includes data for compounds containing unique groups.

(297) B. J. McBride, S. Heimel, J. G. Ehlers, and S. Gordon, NASA Report SP-3001.

(298) M. G. Krishna-Pillai and F. F. Cleveland, *J. Mol. Spectry.*, **6**, 465 (1961).

(299) H. W. Thomson, *Trans. Faraday Soc.*, **37**, 251 (1941).

(300) J. T. Brandenburg, Ph.D. Dissertation, University of Nebraska, 1958; *Dissertation Abstr.*, **18**, 1994 (1958).

(301) K. Ramaswamy, K. Sathianandan, and F. F. Cleveland, *J. Mol. Spectry.*, **9**, 107 (1962).

(302) R. F. Porter and S. K. Gupta, *J. Phys. Chem.*, **68**, 2732 (1964).

(303) H. J. Spangenberg and M. Pfeiffer, *Z. Physik. Chem. (Leipzig)*, **232**, 343 (1966).

(304) H. J. Spangenberg, *ibid.*, **232**, 271 (1966).

(305) S. R. Gunn and J. H. Kindsvater, *J. Phys. Chem.*, **70**, 1750 (1966).

Table XXIX
Organometallic Compounds

Metal	Compound	$\Delta H_f^{\circ}{}_{298}$		Δ	Ref	
		Obsd	Estd			
Tin	(CH ₃) ₄ Sn	-4.1	-4.1	[0]	292	
		-4.6		-0.5	281	
	(CH ₃) ₃ Sn(CH ₂ CH ₃)	-6.5 ^a	-6.3	-0.2		
		-7.1		-0.8	292	
	(CH ₃) ₃ Sn(CH(CH ₃) ₂)	-11.2	-10.8	-0.4	290	
	(CH ₃) ₃ Sn(C)CH ₃) ₃	-16.1	-16.1	[0]	290	
	(CH ₃) ₃ SnCH ₂ C ₆ H ₅	20.2	20.2	[0]	292	
		20.8 ^a		-1.1		
	(CH ₃) ₃ SnC ₆ H ₅	26.7 ^a	26.7	[0]		
		26.1		-1.7	292	
	(CH ₃) ₃ SnCl	-41.8	-40.0	-1.8	289	
	(CH ₃) ₂ SnCl ₂	-70.6	-69.4	-1.2	289	
	CH ₃ SnCl ₃	-99.8	-99.6	-0.2	289	
	(CH ₃) ₂ SnBr	-32.2	-32.0	-0.2	289	
		-33.3		-1.3		
		-33.6		-1.6		
	(CH ₃) ₃ SnI	-20.3	-20.3	[0]	289	
		-17.8		+2.5		
	(CH ₃) ₃ (CH ₂ =CH)Sn	22.1	22.4	-1.4	292	
		22.4 ^a		[0]		
	(CH ₃) ₃ SnSn(CH ₃) ₃	-7.6	-7.7	+0.1	281	
		-5.6		+2.1	292	
	(CH ₃ CH ₂) ₄ Sn	-10.7	-12.8	+2.1	281	
		-8.1		+4.7	292	
	(CH ₃ CH ₂) ₃ SnCl	-46.2	-46.6	+0.4	281, 289	
	(CH ₃ CH ₂) ₃ SnCl ₃	-102.0	-101.8	-0.2	281, 289	
	(CH ₃ CH ₂) ₃ SnSn(CH ₂ CH ₃) ₃	-37.0	-20.8	-16.2	281, 289	
	(CH ₃ CH ₂ CH ₂) ₄ Sn	-34.6	-36.6	+2.0	281, 289	
		-33.2		+3.4		
	(CH ₃ CH ₂ CH ₂) ₃ SnH	-18.8	-16.8	-2.0	291	
	(CH ₃) ₂ CH) ₄ Sn	-29.3	-27.9	-1.4	290	
	(CH ₃ CH ₂ CH ₂ CH ₂) ₄ Sn	-52.9	-52.4	-0.5	281	
		-55.9 ^a		-3.5	292	
	(CH ₃ CH ₂ CH ₂ CH ₂) ₃ SnH	-32.2	-31.7	-0.5	291	
	(CH ₃ CH ₂ CH ₂ CH ₂) ₃ SnBr	-66.0 ^a	-68.4	+2.4	292	
		-65.5		+2.9	281	
			-76.28			
	(CH ₃ CH ₂ CH ₂ CH ₂) ₃ SnCl					
	(CH ₃ CH ₂ CH ₂ CH ₂) ₃ SnCl ₃	-113.5 ^b	-111.7	-1.8	289	
	(C ₆ H ₅) ₄ Sn	114.3	114.3	[0]	294	
	(CH ₂ =CH) ₄ Sn		96.32		289	
	(CH ₂ =CH) ₃ SnCl	36.8 ^b	36.9	[0]	289	
	(CH ₂ =CH) ₂ SnCl ₂	-20.6 ^b	-20.6	[0]	289	
	(CH ₂ =CH)SnCl ₃	-67.1 ^b	-67.1	[0]	289	
	Lead	(CH ₃) ₄ Pb	32.6	32.6	[0]	281
		(CH ₃ CH ₂) ₄ Pb	25.8	25.8	[0]	281
	Chromium	((CH ₃) ₃ CO) ₄ Cr	-305	-305	[0]	293
	Zinc	(CH ₃) ₂ Zn	13.1	13.1	[0]	281
		(CH ₃ CH ₂) ₂ Zn	13.0	9.6	+3.4	281
		10.2		+0.6	281	
	(CH ₃ CH ₂ CH ₂) ₂ Zn	-2.2 ^a	-0.3	-1.9	281	
	(CH ₃ CH ₂ CH ₂ CH ₂) ₂ Zn	-12.2 ^a	-10.2	-2.0	281, 289	
Titanium	(CH ₃ CH ₂ O) ₄ Ti	-325	-325	0	293	
	(CH ₃ CH ₂ CH ₂ O) ₄ Ti	-354	-345	-9	293	
	((CH ₃) ₂ CHO) ₄ Ti	-360	-360	0	293	
	(CH ₃ CH ₂ CH ₂ CH ₂ O) ₄ Ti	-377	-365	-12	293	
	((CH ₃) ₂ CHCH ₂ O) ₄ Ti	-381	-373	-9	293	
	((CH ₃) ₂ (CH ₃ CH ₂ CHO) ₄ Ti	-382	-379	-2	293	
	((CH ₃) ₃ CO) ₄ Ti	-395	-398	+3	293	
	(CH ₃ CH ₂ CH ₂ CH ₂ CH ₂ O) ₄ Ti	-403	-385	-18	293	
	((CH ₃) ₃ CCH ₂ O) ₄ Ti	-438	-418	-20	293	
	((CH ₃ CH ₂) ₂ N) ₄ Ti	-100	-100	[0]	293	
	Vanadium	((CH ₃) ₃ CO) ₄ V	-328	-328	[0]	293
	Cadmium	(CH ₃) ₂ Cd	26.2	26.2	[0]	281
(CH ₃ CH ₂) ₂ Cd		25.5	25.5	[0]	281	

Table XXIX (Continued)

Metal	Compound	ΔH_f°		Δ	Ref
		Obsd	Estd		
Aluminum	(CH ₃) ₃ Al	-21.0	-21.0	[0]	281
	(CH ₃ CH ₂) ₃ Al	-19	-19	[0]	281
Germanium	(CH ₃ CH ₂) ₄ Ge	-34.6	-34.9	+0.3	281
	(CH ₃ CH ₂ CH ₂) ₄ Ge	-54.9	-54.7	-0.2	281
Mercury	(CH ₃) ₂ Hg	22.9	22.3	+0.6	295
		22.3		[0]	281
	CH ₃ HgCl	-12.6	-12.9	+0.3	295
		-12.3		+0.6	281
	CH ₃ HgBr	-4.7	-5.2	+0.5	295
		-4.4		+0.8	281
	CH ₃ HgI	5.2	5.7	-0.5	295
		5.3		-0.4	281
	(CH ₃ CH ₂) ₂ Hg	21.9	17.0	+4.9	295
		17.2		+0.2	281
	(CH ₃ CH ₂)HgCl	-14.2	-15.6	+1.4	294
		-15.8		-0.2	281
	(CH ₃ CH ₂)HgBr	-6.1	-7.88	+1.8	295
		-7.7		+0.2	281
	(CH ₃ CH ₂)HgI	4.7	3.0	+1.7	295
		3.1		+0.1	281
	(CH ₃ CH ₂ CH ₂) ₂ Hg	12.4 ^a	7.08	+5.3	296
		6.7		-0.4	281
	(CH ₃ CH ₂ CH ₂)HgCl	-18.6	20.5	+1.9	296
		-20.6		-0.1	281
	(CH ₃ CH ₂ CH ₂)HgBr	-10.6	-12.8	+2.2	296
		-12.7		+0.1	281
	(CH ₃ CH ₂ CH ₂)HgI	0.8	-1.93	+2.7	296
		-1.2		-0.7	281
	((CH ₃) ₂ CH) ₂ Hg	10.4	9.42	+1.0	296
		9.4		[0]	281
	((CH ₃) ₂ CH)HgCl	-20.1	-19.4	-0.7	296
		-19.7		-0.3	281
	((CH ₃) ₂ CH)HgBr	-13.0	-11.7	-1.3	296
		-12.6		-0.9	281
((CH ₃) ₂ CH)HgI	-1.4	-0.76	-0.6	296	
	-1.1		-0.3	281	
(C ₆ H ₅) ₂ Hg	93.8	93.8	[0]	281	
	90.9		-2.9	295	
	90.5		-3.3	295	
(C ₆ H ₅)HgCl	24.6	24.6	[0]	281	
	27.4		+2.8	295	
(C ₆ H ₅)HgBr	32.6	32.8	-0.2	295	
	32.8		[0]	281	
(C ₆ H ₅)HgI	42.9	42.6	+0.3	295	
	42.6		[0]	281	

• Heat of vaporization corrected to 25°. ^b See text

Table XXX

Gaseous Heats of Formation of Titanium Alkylxides

Compound	ΔH_f° , kcal/mole for [Ti-(O) ₄] + 4[O-(Ti)(C)]
[CH ₃ CH ₂ O] ₄ Ti	-251
[CH ₃ (CH ₂) ₂ O] ₄ Ti	-260
[CH ₃ (CH ₂) ₃ O] ₄ Ti	-263
[CH ₃ (CH ₂) ₄ O] ₄ Ti	-269
[(CH ₃) ₂ CHO] ₄ Ti	-251
[(CH ₃) ₂ CHCH ₂ O] ₄ Ti	-259
[(CH ₃)(CH ₃ CH ₂)CHO] ₄ Ti	-254
[(CH ₃) ₂ CO] ₄ Ti	-248
[(CH ₃ CH ₂)(CH ₃) ₂ CO] ₄ Ti	-285

This section prepared by S. W. Benson and G. R. Haugen.

VIII. Guide to the Use of the Group Tables (Tables XXXIII-XXXVIII)

(a) All values of ΔH_f° are in kcal/mole; S° and C_p° are in cal/(deg mole).

(b) Values of properties calculated from the tables may be corrected to any temperature by the approximations² (parameters in these equations refer to formation reactions).

$$\Delta H_{T_2}^\circ = \Delta H_{T_1}^\circ + \overline{\Delta C_p}^\circ (T_2 - T_1)$$

$$\Delta S_{T_2}^\circ = \Delta S_{T_1}^\circ + \overline{\Delta C_p}^\circ \ln (T_2/T_1)$$

where

$$\overline{\Delta C_p}^\circ = 0.5(\Delta C_{p^\circ T_2} + \Delta C_{p^\circ T_1})$$

Table XXXI (Continued)

Compound	ΔH_f° 298	S_{298}°	C_p°						Rej
			300°	400°	500°	600°	800°	1000°	
$\text{Si}_3\text{O}_4(\text{CH}_3)_{12}$	-667.0								263
$\text{Si}_2(\text{OCH}_3)_6$	-448.0								263
$\text{HC}\equiv\text{CSiH}_3$		64.27	17.31	20.56	23.05	25.05	28.14	30.35	306
H_3SiGeH_3	27.8								305

Table XXXII

Corrections

Group type	gauche	cis	ortho	Ring or steric strain	Other
Alkanes	ΔH_f°
Alkenes	ΔH_f°	$\Delta H_f^\circ, S^\circ, C_p^\circ$
Alkynes
Aromatics	$\Delta H_f^\circ, S^\circ, C_p^\circ$
Polyaromatics	$\Delta H_f^\circ, S^\circ, C_p^\circ$	$\Delta H_f^\circ, S^\circ, C_p^\circ$	Conjugation corrections to enthalpy
Ether oxygen	ΔH_f°
Ditertiary ethers	ΔH_f°	...
Aliphatic nitrogen	ΔH_f°
Pyridine	ΔH_f° ^a
Halogen (except F)	ΔH_f° ^b	...	ΔH_f°
Fluorine	ΔH_f°
Carbon-boron	ΔH_f°

^a Also for *para* substituents. ^b Only for halogen against halogen.

(c) σ is the symmetry number and includes internal, as well as external, symmetry. In these calculations, care must be taken to avoid duplication of rotations as can occur when external and internal rotors share a common axis, *e.g.*, in $\text{CH}_3\text{-CH}_3$ for which $\sigma_{\text{int}} = 3$ and $\sigma_{\text{ext}} = 3 \times 2$, *i.e.*, $\sigma_{\text{tot}} = 3 \times 3 \times 2 = 18$. If σ_{int} is calculated as 3×3 , then σ_{ext} is only 2; otherwise duplicate rotomers were included in the calculation. For the general molecule, where the above complication is absent, the total symmetry number is evaluated as

$$\sigma_{\text{tot}} = \sigma_{\text{ext}} \prod_{i=1}^i (\sigma_{\text{int}})_i$$


where $(\sigma_{\text{int}})_i$ is the rotational degeneracy of the *i*th rotor, *e.g.*, 3 for methyl, 2 for $-\text{BF}_2$, etc., and σ_{ext} is the rotational degeneracy of the molecule as a whole, *e.g.*, 2 for acetone, $\text{CH}_3(\text{C}=\text{O})\text{-CH}_3$. The correction $-R \ln \sigma_{\text{tot}}$ must be added to a calculated S° value.

(d) If the molecule is optically active, $R \ln n$ must be added to its entropy estimate, where n is the total number of stereoisomers of equal energy.

(e) Conformational corrections are applied by adding once for each interaction in the molecule, the value of the appropriate correction to the property to be calculated. If no correction is listed, such interaction as may exist is already included in the relevant groups in as many cases as possible.

The nature of the estimation of the corrections has already been discussed under the appropriate group type, and their values are listed in the relevant group tables. A summary of


the types of correction associated with various classes of compounds is given in Table XXXII. Care should, however, be exercised when calculating properties of any molecule where large groups, several carbon atoms apart, are forced closely together, *e.g.*, in



Predicted values for such compounds may be in error by precisely the value of the distant interaction, *i.e.*, in the above case, the interaction between the *t*-butyl group and the ring hydrogen or methyl. The predicted value thus provides a useful model to estimate the magnitude of such interactions which, as a rough guide, may be expected to arise when atoms can approach to less than the sum of their van der Waals radii. This may be estimated from a molecular model.

(f) Examples

Acetone



$$\begin{aligned} \Delta H_f^\circ &= 2 \times \text{C}-(\text{H})_3(\text{C}) + 1 \times \text{CO}-(\text{C})_2 \\ &= 2 \times -10.08 - 31.5 = \\ &\quad -51.66 \text{ kcal/mole } (-51.70 \text{ obsd}) \\ S^\circ &= 2 \times \text{C}-(\text{H})_3(\text{C}) + 1 \times \text{CO}-(\text{C})_2 - R \ln \sigma \\ \sigma_{\text{ext}} &= 2, \sigma_{\text{int}} = 3 \times 3; \therefore \sigma_{\text{tot}} = 2 \times 3^2 = 18 \\ S^\circ &= 2 \times 30.41 + 15.01 - R \ln 18 = \\ &\quad 70.1 \text{ cal/mole } (70.5 \text{ obsd}) \end{aligned}$$

$$C_p^\circ(300^\circ\text{K}) = 2 \times C-(\text{H})_3(\text{C}) + 1 \times \text{CO}-(\text{C})_2 \\ = 2 \times 6.19 + 5.59 = 17.97 \text{ cal/mole}$$

Triethylamine

This has the groups

$$3 \times C-(\text{C})(\text{H})_3 + 3 \times C-(\text{N})(\text{C})(\text{H})_2 + \text{N}-(\text{C})_3 \\ \Delta H_f^\circ = 3 \times -10.08 = -30.24 \\ + 3 \times -6.6 = -19.8 \\ \underline{-50.04}$$

$$+ 24.4 = +24.4$$

$$+ 3 \text{ gauche interactions} \\ \text{at } 0.8 \text{ each} = +2.4$$

$$\therefore \Delta H_f^\circ = -50.04 + 26.8 = -23.24 \text{ kcal/mole} \\ \Delta H_f^\circ \text{ obsd} = -23.9 \text{ kcal/mole}$$

1,1-Dimethylhydrazine

This has the groups

 $\sigma_{\text{rot}} = 3 \times 3 = 9$ and two optical isomers exist

$$S^\circ = 2 \times 30.41 = +60.82$$

$$1 \times -13.8 = -13.8$$

$$1 \times +29.13 = +29.13$$

$$\underline{+89.95}$$

$$-R \ln(3 \times 3/2) = -2.98$$

$$\underline{-16.78}$$

i.e.

$$S^\circ = +89.95 - 16.78 = 73.17 \text{ cal/mole (obsd } 72.82 \text{ cal/mole)}$$

This section prepared by S. W. Benson and F. R. Cruickshank.

Together with Dr. S. W. Benson, the authors of the various sections of this joint project were: hydrocarbons, D. M. Golden and H. E. O'Neal; oxygen compounds, R. Shaw; nitrogen compounds, R. Walsh; halogen compounds, A. S. Rodgers; sulfur, boron, and organometallic compounds, G. R. Haugen; sections I, II, VIII, inclusion of recent data, and construction of the whole as a review, F. R. Cruickshank.

Table XXXIII

Hydrocarbon Groups

Group	$\Delta H_f^\circ_{298}$	S°_{298}	C_p°						
			300°	400°	500°	600°	800°	1000°	1500°
C-(H) ₃ (C)	-10.08	30.41	6.19	7.84	9.40	10.79	13.02	14.77	17.58
C-(H) ₂ (C) ₂	-4.95	9.42	5.50	6.95	8.25	9.35	11.07	12.34	14.25
C-(H)(C) ₃	-1.90	-12.07	4.54	6.00	7.17	8.05	9.31	10.05	11.17
C-(C) ₄	0.50	-35.10	4.37	6.13	7.36	8.12	8.77	8.76	8.12
C _d -(H) ₂	6.26	27.61	5.10	6.36	7.51	8.50	10.07	11.27	13.19
C _d -(H)(C)	8.59	7.97	4.16	5.03	5.81	6.50	7.65	8.45	9.62
C _d -(C) ₂	10.34	-12.7	4.10	4.61	4.99	5.26	5.80	6.08	6.36
C _d -(C _d)(H)	6.78	6.38	4.46	5.79	6.75	7.42	8.35	8.99	9.98
C _d -(C _d)(C)	8.88	-14.6	(4.40)	(5.37)	(5.93)	(6.18)	(6.50)	(6.62)	(6.72)
[C _d -(C _B)(H)]	6.78	6.4	4.46	5.79	6.75	7.42	8.35	8.99	9.98
C _d -(C _B)(C)	8.64	(-14.6)	(4.40)	(5.37)	(5.93)	(6.18)	(6.50)	(6.62)	(6.72)
[C _d -(C _d)(H)]	6.78	6.4	4.46	5.79	6.75	7.42	8.35	8.99	9.98
C-(C _d)(C)(H) ₂	-4.76	9.8	5.12	6.86	8.32	9.49	11.22	12.48	14.36
C-(C _d) ₂ (H) ₂	-4.29	(10.2)	(4.7)	(6.8)	(8.4)	(9.6)	(11.3)	(12.6)	(14.4)
C-(C _d)(C _B)(H) ₂	-4.29	(10.2)	(4.7)	(6.8)	(8.4)	(9.6)	(11.3)	(12.6)	(14.4)
C-(C _d)(C)(H) ₂	-4.73	10.3	4.95	6.56	7.93	9.08	10.86	12.19	14.20
C-(C _B)(C)(H) ₂	-4.86	9.3	5.84	7.61	8.98	10.01	11.49	12.54	13.76
C-(C _d)(C) ₂ (H)	-1.48	(-11.7)	(4.16)	(5.91)	(7.34)	(8.19)	(9.46)	(10.19)	(11.28)
C-(C _d)(C) ₂ (H)	-1.72	(-11.2)	(3.99)	(5.61)	(6.85)	(7.78)	(9.10)	(9.90)	(11.12)
C-(C _B)(C) ₂ (H)	-0.98	(-12.2)	(4.88)	(6.66)	(7.90)	(8.75)	(9.73)	(10.25)	(10.68)
C-(C _d)(C) ₃	1.68	(-34.72)	(3.99)	(6.04)	(7.43)	(8.26)	(8.92)	(8.96)	(8.23)
C-(C _B)(C) ₃	2.81	(-35.18)	(4.37)	(6.79)	(8.09)	(8.78)	(9.19)	(8.96)	(7.63)
C _t -(H)	26.93	24.7	5.27	5.99	6.49	6.87	7.47	7.96	8.85
C _t -(C)	27.55	6.35	3.13	3.48	3.81	4.09	4.60	4.92	6.35
C _t -(C _d)	29.20	(6.43)	(2.57)	(3.54)	(3.50)	(4.92)	(5.34)	(5.50)	(5.80)
C _t -(C _B)	(29.20)	6.43	2.57	3.54	3.50	4.92	5.34	5.50	5.80
C _B -(H)	3.30	11.53	3.24	4.44	5.46	6.30	7.54	8.41	9.73
C _B -(C)	5.51	-7.69	2.67	3.14	3.68	4.15	4.96	5.44	5.98
C _B -(C _d)	5.68	-7.80	3.59	3.97	4.38	4.72	5.28	5.61	5.75
[C _B -(C _t)]	5.7	-7.80	3.59	3.97	4.38	4.72	5.28	5.61	5.75
C _B -(C _B)	4.96	-8.64	3.33	4.22	4.89	5.27	5.76	5.95	(6.05)
C _a	34.20	6.0	3.9	4.4	4.7	5.0	5.3	5.5	5.7

Table XXXIII (Continued)
Next-Nearest Neighbor Corrections















Group	ΔH_f° ₂₉₈	S° ₂₉₈	C_p°						
			300°	400°	500°	600°	800°	1000°	1500°
Alkane <i>gauche</i> correction	0.80								
Alkene <i>gauche</i> correction	0.50								
<i>cis</i> correction	1.00 ^a	<i>b</i>	-1.34	-1.09	-0.81	-0.61	-0.39	-0.26	0
<i>ortho</i> correction	0.57	-1.61	1.12	1.35	1.30	1.17	0.88	0.66	-0.05
Corrections to be Applied to Ring Compound Estimates									
Ring (σ)	ΔH_f° ₂₉₈	S° ₂₉₈	C_p°						
			300°	400°	500°	600°	800°	1000°	1500°
Cyclopropane (6)	27.6	32.1	-3.05	-2.53	-2.10	-1.90	-1.77	-1.62	(-1.52)
Methylenecyclopropene	40.9								
Cyclopropene (2)	53.7	33.6							
Cyclobutane (8)	26.2	29.8	-4.61	-3.89	-3.14	-2.64	-1.88	-1.38	-0.67
Cyclobutene (2)	29.8	29.0	-2.53	-2.19	-1.89	-1.68	-1.48	-1.33	-1.22
Cyclopentane (10)	6.3	27.3	-7.50	-6.49	-5.40	-4.37	-2.93	-1.93	-0.37
Cyclopentene (2)	5.9	25.8	-5.98	-5.35	-4.89	-4.14	-2.93	-2.26	-1.08
Cyclopentadiene	6.0								
Cyclohexane (6)	0	18.8	-6.40	-4.60	-3.30	-1.60	0.82	1.98	3.19
Cyclohexene (2)	1.4	21.5	-4.28	-3.04	-1.98	-1.43	-0.29	0.08	0.81
Cyclohexadiene-1,3	4.8								
Cyclohexadiene-1,4	0.5								
Cycloheptane (1)	6.4	15.9							
Cycloheptene	5.4								
Cycloheptadiene-1,3	6.6								
Cycloheptatriene-1,3,5 (1)	4.7	23.7							
Cyclooctane (8)	9.9	16.5							
<i>cis</i> -Cyclooctene	6.0								
<i>trans</i> -Cyclooctene	15.3								
Cyclooctatriene-1,3,5	8.9								
Cyclooctatetraene	17.1								
Cyclononane	12.8								
<i>cis</i> -Cyclononene	9.9								
<i>trans</i> -Cyclononene	12.8								
Spiropentane (4)	63.5	67.6							
Bicyclo[1.1.0]butane (2)	67.0	69.2							
Bicyclo[2.1.0]pentane	55.3								
Bicyclo[3.1.0]hexane	32.7								
Bicyclo[4.1.0]heptane	28.9								
Bicyclo[5.1.0]octane	29.6								
Bicyclo[6.1.0]nonane	31.1								

^a When one of the groups is *t*-butyl, *cis* correction = 4.00; when both are *t*-butyl, *cis* correction = ~10.00; and when there are two *cis* corrections around one double bond, the total correction is 3.00. ^b +1.2 for but-2-ene, 0 for all other 2-enes, and -0.6 for 3-enes.

Table XXXIV
Oxygen-Containing Groups



Group	ΔH_f°	S°	C_p°						
			300°	400°	500°	600°	800°	1000°	1500°
CO-(CO)(C)	-29.2								
CO-(O)(C _a)	-33.5								
CO-(O)(C _B)	-46.0								
CO-(O)(C)	-33.4	14.78	5.97	6.70		8.02	8.87	9.36	
[CO-(O)(H)] ^a	-29.5	34.93	7.03	7.87	8.82	9.68	11.16	12.20	
CO-(C _a)(H)	-31.7								
CO-(C _B) ₂	-39.1								
CO-(C _B)(C)	-37.6								
[CO-(C _B)(H)] ^b	-31.7								
CO-(C) ₂	-31.5	15.01	5.59	6.32	7.09	7.76	8.89	9.61	
CO-(C)(H)	-29.6	34.93	7.03	7.87	8.82	9.68	11.16	12.20	
CO-(H) ₂	-27.7	53.67	8.47	9.38	10.46	11.52	13.37	14.81	
O-(CO) ₂	-50.9								
O-(CO)(O)	-19.0								
[O-(CO)(C _a) ^c]	-41.3								
O-(CO)(C)	-41.3	8.39							
O-(CO)(H) _i	-60.3	24.52	3.81	4.98	5.80	6.34	7.19	7.75	
O-(O)(O) _i	(19.0)	(9.4)	(3.7)	(3.7)	(3.7)	(3.7)	(4.2)	(4.2)	(4.8)
O-(O)(C)	(-4.5)	(9.4)	(3.7)	(3.7)	(3.7)	(3.7)	(4.2)	(4.2)	(4.8)
O-(O)(H)	-16.27	27.85	5.17	5.79	6.28	6.66	7.15	7.51	8.17

Table XXXIV (Continued)

Group	ΔH_f°	S°	C_p°						
			300°	400°	500°	600°	800°	1000°	1500°
O-(C _d) ₂	-32.8								
O-(C _d)(C)	-31.3								
O-(C _B) ₂	-19.3								
O-(C _B)(C)	-22.6								
[O-(C _B)(H) ^a]	-37.9	29.1	4.3	4.5	4.8	5.2	6.0	6.6	
O-(C) ₂	-23.7	8.68	3.4	3.7	3.7	3.8	4.4	4.6	
O-(C)(H)	-37.88	29.07	4.33	4.45	4.82	5.23	6.02	6.61	
C _d -(CO)(O)	6.3								
C _d -(CO)(C)	9.4								
[C _d -(CO)(H) ^a]	7.7								
[C _d -(O)(C _d) ^f]	8.9								
[C _d -(O)(C) ^g]	10.3								
[C _d -(O)(H) ^h]	8.6		4.16	5.03	5.81	6.50	7.65	8.45	9.62
C _B -(CO)	9.7								
C _B -(O)	-1.8	-10.2	3.9	5.3	6.2	6.6	6.9	6.9	
C-(CO) ₂ (H) ₂	-7.2								
C-(CO)(C) ₃	1.58								
C-(CO)(C) ₂ (H)	-1.83 ⁱ	-12.0							
C-(CO)(C)(H) ₂	-5.0	9.6	6.2	7.7	8.7	9.5	11.1	12.2	
[C-(CO)(H) ₃ ^j]	-10.08	30.41	6.19	7.84	9.40	10.79	13.02	14.77	17.58
C-(O) ₂ (C) ₂	-16.8								
C-(O) ₂ (C)(H)	-17.2								
C-(O) ₂ (H) ₂	-17.7								
C-(O)(C _B)(H) ₂	-6.6	9.7							
C-(O)(C _d)(H) ₂	-6.9								
C-(O)(C) ₃	-6.60	-33.56	4.33	6.19	7.25	7.70	8.20	8.24	
C-(O)(C) ₂ (H)	-7.00	-11.00	4.80	6.64	8.10	8.73	9.81	10.40	
C-(O)(C)(H) ₂	-8.5	10.3	4.99	6.85	8.30	9.43	11.11	12.33	
[C-(O)(H) ₃ ^k]	-10.08	30.41	6.19	7.84	9.40	10.79	13.02	14.77	17.58
Strain									
Ether oxygen <i>gauche</i>	0.3								
Ditertiary ethers	8.4								
	27.6	31.4	-2.0	-2.8	-3.0	-2.6	-2.3	-2.3	
	26.4	27.7	-4.6	-5.0	-4.2	-3.5	-2.6	+0.2	
	6.7								
	2.2								
	3.5								
	5.4								
	3.4								
	-6.2								
	2.5								
	6.0								
	3.4								
	1.1								
	1.4								
	4.6								

^a CO-(O)(H) ≡ CO-(C)(H), assigned. ^b CO-(C_B)(H) ≡ CO-(C_d)(H), assigned. ^c O-(CO)(C_d) ≡ O-(CO)(C), assigned. ^d O-(C_B)(H) ≡ O-(C)(H), assigned. ^e C_d-(CO)(H) ≡ 1/2[C_d-(C_d)(H) + C_d-(C)(H)], assigned. ^f C_d-(O)(C_d) ≡ C_d-(C_d)(C), assigned. ^g C_d-(O)(C) ≡ C_d-C₂, assigned. ^h C_d-(O)(H) ≡ C_d-(C)(H), assigned. ⁱ Estimated values. ^j C-(CO)(H)₂ ≡ C-(C)(H)₃, assigned. ^k C-(O)(H)₃ ≡ C-(C)(H)₃, assigned.

Table XXXV (Continued)

Group	ΔH_f°	S°	C_p°						
			300°	400°	500°	600°	800°	1000°	1500°
	3.4								
	8.5								

^a N_I = double-bonded nitrogen in imines; $N_I-(C_B)$ = pyridine N. N_A = double-bonded nitrogen in azo compounds. ^b No *cis* corrections applied to imines or azo compounds. ^c *gauche* corrections of +0.8 kcal/mole to ΔH_f° applied just as for hydrocarbons. ^d Estimates by authors; see text under appropriate section. ^e For *ortho* or *para* substitution in pyridine add -1.5 kcal/mole per group. ^f $C-(N_A)(H)_3 \equiv C-(N)(H)_2 \equiv C-(C)(H)_3$ by convention.

Table XXXVI

Approximate Group Contributions to ΔH_f° at 25° and 1 Atm: Miscellaneous Compounds

Group	ΔH_f°	Data Sources and References
$N-(NO_2)(C)_2$	17.8	From $\Delta H_f^\circ((CH_3)_2NNO_2) = -0.3$ kcal/mole and $\Delta H_f^\circ((C_2H_5)_2NNO_2) = -13.8$ kcal/mole ^a
$C-(N)_3(C)(H)_2$	70	From $\Delta H_f^\circ(C_2H_5N_3) = 59$ kcal/mole (ref 170)
$N-(N)(C_B)_2$	36.3	From $\Delta H_f^\circ((C_6H_5)_2NN(C_6H_5)_2) = 144.5$ kcal/mole (ref 119)
N_I-N_I	21.6	From $\Delta H_f^\circ(C_6H_5CH=NN=CHC_6H_5) = 104.2$ kcal/mole (ref 138)
N_I-C_B	14.1	From $\Delta H_f^\circ(C_6H_5CH=NC_6H_5) = 61.4$ kcal/mole (ref 138)
C_B-N_I	-0.5	
$C-(NO)(C)(H)_2$	19.1	From $\Delta H_f^\circ(C_2H_5NO) = 9$ kcal/mole, $\Delta H_f^\circ(i-C_3H_7NO) = 2$ kcal/mole, $\Delta H_f^\circ(t-C_4H_9NO) = -7$ kcal/mole. These estimates were, in turn, based upon primary, secondary, and tertiary C-NO bond strengths of 38.5, 37, and 35.5 kcal/mole, respectively (see text)
$C-(NO)(C)_2(H)$	22.2	
$C-(NO)(C)_3$	23.2	
$N-(O)(H)_2$	4.8	From $\Delta H_f^\circ(NH_2OH) = -9.3$ kcal/mole $\Delta H_f^\circ(CH_3NHOH) = -12.0$ kcal/mole, and $\Delta H_f^\circ(NH_2OCH_3) = -6.2$ kcal/mole (see text)
$N-(O)(C)(H)$	12.2	
$O-(N)(H)$	-14.1	See text
$O-(N)(C)$	-0.9	
$N_I-(OH)-5$	-5.0	From $\Delta H_f^\circ(1) = 9.4$ kcal/mole (see text)
$C-(N)_2(H)_2$	-12.1	

^a From R. C. Cass, S. E. Fletcher, C. T. Mortimer, P. G. Quincey, and H. D. Springall, *J. Chem. Soc.*, 958 (1958).

Table XXXVII

Halogen-Containing Groups

Group	ΔH_f°	S°	C_p°						
			300°	400°	500°	600°	800°	1000°	
$C_d-(F)_2$	-77.5	37.3	9.7	11.0	12.0	12.7	13.8	14.5	
$C_d-(Cl)_2$	-1.8	42.1	11.4	12.5	13.3	13.9	14.6	15.0	
$C_d-(Br)_2$		47.6	12.3	13.2	13.9	14.3	14.9	15.2	
$C_d-(F)(Cl)$		39.8	10.3	11.7	12.6	13.3	14.2	14.7	
$C_d-(F)(Br)$		42.5	10.8	12.0	12.8	13.5	14.3	14.7	
$C_d-(Cl)(Br)$		45.1	12.1	12.7	13.5	14.1	14.7		
$C_d-(F)(H)$	-37.6	32.8	6.8	8.4	9.5	10.5	11.8	12.7	
$C_d-(Cl)(H)$	2.1	35.4	7.9	9.2	10.3	11.2	12.3	13.1	
$C_d-(Br)(H)$	12.7	38.3	8.1	9.5	10.6	11.4	12.4	13.2	
$C_d-(I)(H)$	24.5	40.5	8.8	10.0	10.9	11.6	12.6	13.3	
$C_t-(Cl)$		33.4	7.9	8.4	8.7	9.0	9.4	9.6	
$C_t-(Br)$		36.1	8.3	8.7	9.0	9.2	9.5	9.7	
$C_t-(I)$		37.9	8.4	8.8	9.1	9.3	9.6	9.8	
Arenes									
$C_B-(F)$	-42.8	16.1	6.3	7.6	8.5	9.1	9.8	10.2	
$C_B-(Cl)$	-3.8	18.9	7.4	8.4	9.2	9.7	10.2	10.4	
$C_B-(Br)$	10.7	21.6	7.8	8.7	9.4	9.9	10.3	10.5	
$C_B-(I)$	24.0	23.7	8.0	8.9	9.6	9.9	10.3	10.5	
$C-(C_B)(F)_2$	-162.7	42.8	12.5	15.3	17.2	18.5	20.1	21.0	
$C-(C_B)(Br)(H)_2$	-6.9								
$C-(C_B)(I)(H)_2$	8.4								

Table XXXVII (Continued)

Group	ΔH_f°	S°	C_p°					
			300°	400°	500°	600°	800°	1000°
Corrections for Next-Nearest Neighbors								
<i>ortho</i> (F)(F)	5.0	0	0	0	0	0	0	0
<i>ortho</i> (Cl)(Cl)	2.2							
<i>ortho</i> (alk)(halogen)	0.6							
<i>cis</i> (halogen) (halogen)	(0)							
<i>cis</i> (halogen)(alk)	(0)							
C-(F) ₃ (C)	-158.4	42.5	12.7	15.0	16.4	17.9	19.3	20.0
C-(F) ₂ (H)(C)	(-109.3)	39.1	9.9	12.0		15.1		
C-(F)(H) ₂ (C)	-51.8	35.4	8.1	10.0	12.0	13.0	15.2	16.6
C-(F) ₂ (C) ₂	-97.0	17.8	9.9	11.8	13.5			
C-(F)(H)(C) ₂	-48.4	(14.0)						
C-(F)(C) ₃	-43.9							
C-(F) ₂ (Cl)(C)	-106.3	40.5	13.7	16.1	17.5			
C-(Cl) ₃ (C)	-20.7	50.4	16.3	18.0	19.1	19.8	20.6	21.0
C-(Cl) ₂ (H)(C)	(-18.9)	43.7	12.1	14.0	15.4	16.5	17.9	18.7
C-(Cl)(H) ₂ (C)	-15.6	37.8	8.9	10.7	12.3	13.4	15.3	16.7
C-(Cl) ₂ (C) ₂	(-19.5)							
C-(Cl)(H)(C) ₂	-12.8	17.6	9.0	9.9	10.5	11.2		
C-(Cl)(C) ₃	-12.8	-5.4	9.3	10.5	11.0	11.3		
C-(Br) ₃ (C)		55.7	16.7	18.0	18.8	19.4	19.9	20.3
C-(Br)(H) ₂ (C)	-5.4	40.8	9.1	11.0	12.6	13.7	15.5	16.8
C-(Br)(H)(C) ₂	-3.4							
C-(Br)(C) ₃	-0.4	-2.0	9.3	11.0				
C-(I)(H) ₂ (C)	7.95	42.5	9.2	11.0	12.9	13.9	15.8	17.2
C-(I)(H)(C) ₂	10.7	22.2	(8.7)					
C-(I)(C) ₃	13.0	(0.0)	(9.7)					
C-(Cl)(Br)(H)(C)		45.7	12.4	14.0	15.6	16.3	17.9	19.0
N-(F) ₂ (C)	-7.8							
C-(Cl)(C)(O)(H)	-22.3	15.4	(9.0)	(9.9)	(10.5)	(11.2)		
Corrections for Next-Nearest Neighbors								
<i>gauche</i> (halogen) (alk)	0.0	0	0	0	0	0	0	0
<i>gauche</i> (halogen) (exclusive of F)	1.0							
<i>gauche</i> (F) (halogen)	0.0							

Table XXXVIII










Sulfur-Containing Groups

Group	ΔH_f°	S°	C_p°					
			300°	400°	500°	600°	800°	1000°
[C-(H) ₃ (S) ^a]	-10.08	30.41	6.19	7.84	9.40	10.79	13.02	14.77
C-(C)(H) ₂ (S)	-5.65	9.88	5.38	7.08	8.60	9.97	12.26	14.15
C-(C) ₂ (H)(S)	-2.64	-11.32	4.85	6.51	7.78	8.69	9.90	10.57
C-(C) ₃ (S)	-0.55	-34.41	4.57	6.27	7.45	8.15	8.72	8.10
C-(C _B)(H) ₂ (S)	-4.73							
C-(C _d)(H) ₂ (S)	-6.45							
[C _B -(S) ^b]	-1.8	10.20	3.90	5.30	6.20	6.60	6.90	6.90
[C _d -(H)(S) ^c]	8.56	8.0	4.16	5.03	5.81	6.50	7.65	8.45
C _d -(C)(S)	10.93	-12.41	3.50	3.57	3.83	4.09	4.41	5.00
S-(C)(H)	4.62	32.73	5.86	6.20	6.51	6.78	7.30	7.71
S-(C _B)(H)	11.96	12.66	5.12	5.26	5.57	6.03	6.99	7.84
S-(C) ₂	11.51	13.15	4.99	4.96	5.02	5.07	5.41	5.73
S-(C)(C _d)	9.97							
S-(C _d) ₂	-4.54	16.48	4.79	5.58	5.53	6.29	7.94	9.73
S-(C _B)(C)	19.16							
S-(C _B) ₂	25.90							
S-(S)(C)	7.05	12.37	5.23	5.42	5.51	5.51	5.38	5.12
S-(S)(C _B)	14.5							
S-(S) ₂	3.04	13.36	4.66	4.97	5.13	5.23	5.33	5.39
[C-(SO)(H) ₃ ^d]	-10.08	30.41	6.19	7.84	9.40	10.79	13.02	14.77
C-(C)(SO)(H) ₂	-7.72							
C-(C) ₃ (SO)	-3.05							
C-(C _d)(SO)(H) ₂	-7.35							

Table XXXVIII (Continued)

Group	ΔH_f°	S°	C_p°						
			300°	400°	500°	600°	800°	1000°	
[C _B -(SO) ^q]	2.3								
SO-(C) ₂	-14.41	18.10	8.88	10.03	10.50	10.79	10.98	11.17	
SO-(C _B) ₂	-12.0								
[C-(SO ₂)(H) ₃]	-10.08	30.41	6.19	7.84	9.40	10.79	13.02	14.77	
C-(C)(SO ₂)(H) ₂	-7.68								
C-(C) ₂ (SO ₂)(H)	-2.62								
C-(C) ₃ (SO ₂)	-0.61								
C-(C _d)(SO ₂)(H) ₂	-7.14								
C-(C _B)(SO ₂)(H) ₂	-5.54								
[C _B -(SO ₂) ^q]	2.3								
C _d -(H)(SO ₂)	12.53								
C _d -(C)(SO ₂)	14.47								
SO ₂ -(C) ₂	-69.74	20.90	11.52						
SO ₂ -(C)(C _B)	-72.29								
SO ₂ -(C _B) ₂	-68.58								
[SO ₂ -(C _d)(C _B) ^h]	-68.58								
SO ₂ -(C _d) ₂	-73.58								
SO ₂ -(SO ₂)(C _B)	-76.25								
[CO-(S)(C) ⁱ]	-31.56	15.43	5.59	6.32	7.09	7.76	8.89	9.61	
S-(H)(CO)	-1.41	31.20	7.63	8.09	8.12	8.17	8.50	8.24	
C-(S)(F) ₃		38.9							
S-(C)(CN)	37.18	41.06	9.51	10.44	11.22	11.86	12.85	13.50	
[CS-(N) ₂]	-31.56	15.43	5.59	6.32	7.09	7.76	8.89	9.61	
N-(CS)(H) ₂	12.78	29.19	6.07	7.28	8.18	8.91	10.09	10.98	
[S-(S)(N) ^h]	-4.90								
N-(S)(C) ₂	29.9								
[SO-(N) ₂]	-31.56								
N-(SO)(C) ₂	16.0								
[SO ₂ -(N) ₂ ^m]	-31.56								
N-(SO ₂)(C) ₂	-20.4								

Organosulfur Compounds Ring Corrections

Ring (σ)									
(a)  (2)	17.7	29.47	-2.85	-2.59	-2.66	-3.02	-4.32	-5.82	
(b)  (2)	19.37	27.18	-4.59	-4.18	-3.91	-3.91	-4.60	-5.70	
(c)  (2)	1.73	23.56	-4.90	-4.67	-3.68	-3.66	-4.41	-5.57	
(d)  (2)	0	17.46	-6.22	-4.26	-2.24	-0.69	0.86	1.29	
(e)  (2)	3.89								
(f)  (2)	5.07								
(g)  (1)	5.07								
(h)  (2)	5.74								
(i)  (2)	1.73	23.56	-4.90	-4.67	-3.68	-3.66	-4.41	-5.57	

Organophosphorus Groups^p

Group	ΔH_f°	S°	Remarks
[C-(P)(H) ₃]	-10.08	30.41	C-(P)(H) ₃ ≡ C-(C)(H) ₃ , assigned
C-(P)(C)(H) ₂	-2.47		
[C-(PO)(H) ₂]	-10.08	30.41	C-(PO)(H) ₂ ≡ C-(C)(H) ₂ , assigned
C-(PO)(C)(H) ₂	-3.4		
[C-(P:N)(H) ₃]	-10.08	30.41	C-(P:N)(H) ₃ ≡ C-(C)(H) ₃ , assigned
C-(N:P)(C)(H) ₂	19.4		
[C _B -(P)]	-1.8		C _B -(P) ≡ C _B -(O), assigned
[C _B -(PO)]	2.3		C _B -(PO) ≡ C _B -(CO), assigned
[C _B -(P:N)]	2.3		C _B -(P:N) ≡ C _B -(CO), assigned
P-(C) ₃	7.04		

Table XXXVIII (Continued)

Group	ΔH_f°	S°	Remarks
P-(C)(Cl) ₂	-50.1		
P-(C _B) ₃	28.3		
P-(O) ₃	-66.8		
[P-(N) ₃]	-66.8		P-(N) ₃ ≡ P-(O) ₃ , assigned
PO-(C) ₃	-72.8		
PO-(C)(F) ₂		46.77	
PO-(C)(Cl)(F)		50.80	
PO-(C)(Cl) ₂	-123.0	52.97	
PO-(C)(O)(Cl)	-112.6		
PO-(C)(O) ₂	-99.5		
PO-(O) ₃	-104.6		
PO-(O) ₂ (F)	-167.7		
PO-(C _B) ₃	-52.9		
[PO-(N) ₃]	-104.6		PO-(N) ₃ ≡ PO-(O) ₃ , assigned
[O-(C)(P)]	-23.5		O-(C)(P) ≡ O-(C) ₂ , assigned
O-(H)(P)	-58.7		
[O-(C)(PO)]	-40.7		O-(C)(PO) ≡ O-(C)(CO), assigned
O-(H)(PO)	-65.0		
O-(PO) ₂	-54.5		
[O-(P:N)(C)]	-40.7		O-(P:N)(C) ≡ O-(C)(CO), assigned
N-(P)(C) ₂	32.2		
N-(PO)(C) ₂	17.8		
[P:N-(C) ₃ (C)]	0.50		P:N-(C) ₃ (C) ≡ C-(C) ₄ , assigned
P:N-(C _B) ₃ (C)	-25.7		
P:N-(N:P)(C) ₂ (P:N)	-15.5		
P:N-(N:P)(C _B) ₃ (P:N)	-22.9		
P:N-(N:P)(Cl) ₂ (P:N)	-58.2		
P:N-(N:P)(O) ₂ (P:N)	-43.4		

Organoboron Groups^a

Group	ΔH_f°	Remarks
[C-(B)(H) ₃]	-10.08	C-(B)(H) ₃ ≡ C-(C)(H) ₃ , assigned
C-(B)(C)(H) ₂	-2.22	
C-(B)(C) ₂ (H)	1.1	
[C-(BO ₃)(H) ₃]	-10.08	C-(BO ₃)(H) ₃ ≡ C-(C)(H) ₃ , assigned
C-(BO ₃)(C)(H) ₂	-2.2	
C _d -(B)(H)	15.6	
B-(C) ₃	0.9	
B-(C)(F) ₂	-187.9	
B-(C) ₂ (Cl)	-42.7	
B-(C) ₂ (Br)	-26.9	
B-(C) ₂ (I)	-8.9	
[B-(C) ₂ (O)]	29.3	B-(C) ₂ (O) ≡ N-(C) ₂ (N), assigned
[B-(C _d)(F) ₂]	-192.9	B-(C _d)(F) ₂ ≡ B-(C)(F) ₂ , assigned
[B-(O) ₃]	24.4	B-(O) ₃ ≡ B-(N) ₃ , assigned
B-(O) ₂ (Cl)	-19.7	
B-(O)(Cl) ₂	-61.2	
B-(O) ₂ (H)	19.9	
B-(N) ₃	24.4	B-(N) ₃ ≡ N-(C) ₃ , assigned
B-(N) ₂ (Cl)	-23.8	
B-(N)(Cl) ₂	-67.9	
BO ₃ -(C) ₃	-208.7	
O-(B)(H)	-115.5	
O-(B)(C)	-69.43	
N-(B)(C) ₂	-9.93	
[B-(S) ₃]	24.4	B-(S) ₃ ≡ B-(O) ₃ , assigned
S-(B)(C)	-14.5	
S-(B)(C _B)	-7.8	

Organometallic Groups^a

Metal	Group	ΔH_f°	Remarks
Tin	[C-(Sn)(H) ₃]	-10.08	C-(Sn)(H) ₃ ≡ C-(C)(H) ₃ , assigned
	C-(Sn)(C)(H) ₂	-2.18	
	C-(Sn)(C) ₂ (H)	3.38	
	C-(Sn)(C) ₃	8.16	
	C-(Sn)(C _B)(H) ₂	-7.77	

Table XXXVIII (Continued)

Metal	Group	ΔH_f°	Remarks
	[C _B -(Sn)]	5.51	C _B -(Sn) \equiv C _B -(C), assigned
	[C _d -(Sn)(H)]	8.77	C _d -(Sn)(H) \equiv C _d -(C)(H), assigned
	Sn-(C) ₄	36.2	
	Sn-(C) ₃ (Cl)	-9.8	
	Sn-(C) ₂ (Cl) ₂	-49.2	
	Sn-(C)(Cl) ₃	-89.5	
	Sn-(C) ₃ (Br)	-1.8	
	Sn-(C) ₃ (I)	9.9	
	Sn-(C) ₃ (H)	34.8	
	[Sn-(C _d) ₄]	36.2	Sn-(C _d) ₄ \equiv Sn-(C) ₄ , assigned
	Sn-(C _d) ₃ (Cl)	-8.2	
	Sn-(C _d) ₂ (Cl) ₂	-50.7	
	Sn-(C _d)(Cl) ₃	-82.2	
	Sn-(C) ₃ (C _d)	37.6	
	Sn-(C _B) ₄	26.26	
	Sn-(C) ₃ (C _B)	34.93	
	Sn-(C) ₃ (Sn)	26.4	
Lead	[C-(Pb)(H) ₃]	-10.08	C-(Pb)(H) ₃ \equiv C-(C)(H) ₃ , assigned
	C-(Pb)(C)(H) ₂	-1.70	
	Pb-(C) ₄	72.9	
Chromium	[O-(Cr)(C)]	-23.5	O-(Cr)(C) \equiv O-(Ti)(C), assigned
	Cr-(O) ₄	-64.0	
Zinc	[C-(Zn)(H) ₃]	-10.08	C-(Zn)(H) ₃ \equiv C-(C)(H) ₃ , assigned
	C-(Zn)(C)(H) ₂	-1.78	
	Zn-(C) ₂	33.3	
Titanium	[O-(Ti)(C)]	-23.5	O-(Ti)(C) \equiv O-(P)(C), assigned
	Ti-(O) ₄	-157	
	[N-(Ti)(C) ₂]	39.1	N-(Ti)(C) ₂ \equiv N-(P)(C) ₂ , assigned
	Ti-(N) ₄	-123	
Vanadium	[O-(V)(C)]	-23.5	O-(V)(C) \equiv O-(Ti)(C), assigned
	V-(O) ₄	-87.0	
Cadmium	[C-(Cd)(H) ₃]	-10.08	C-(Cd)(H) ₃ \equiv C-(C)(H) ₃ , assigned
	C-(C _d)(C)(H) ₂	-0.30	
	C _d -(C) ₂	46.4	
Aluminum	[C-(Al)(H) ₃]	-10.08	C-(Al)(H) ₃ \equiv C-(C)(H) ₃ , assigned
	C-(Al)(C)(H) ₂	0.70	
	Al-(C) ₃	9.2	
Germanium	C-(Ge)(C)(H) ₂	-7.7	
	[Ge-(C) ₄]	36.2	Ge-(C) ₄ \equiv Sn-(C) ₄ , assigned
	Ge-(Ge)(C) ₃	15.6	
Mercury	[C-(Hg)(H) ₃]	-10.08	C-(Hg)(H) ₃ \equiv C-(C)(H) ₃ , assigned
	C-(Hg)(C)(H) ₂	-2.68	
	C-(Hg)(C) ₂ (H)	3.62	
	[C _B -(Hg)]	-1.8	C _B -(Hg) \equiv C _B -(O), assigned
	Hg-(C) ₂	42.5	
	Hg-(C)(Cl)	-2.82	
	Hg-(C)(Br)	4.88	
	Hg-(C)(I)	15.78	
	Hg-(C _B) ₂	64.4	
	Hg-(C _B)(Cl)	9.9	
	Hg-(C _B)(Br)	18.1	
	Hg-(C _B)(I)	27.9	

^a C-(S)(H)₃ \equiv C-(C)(H)₃, assigned. ^b C_B-(S) \equiv C_B-(O), assigned. ^c C_d-(S)(H) \equiv C_d-(O)(H), assigned. ^d C-(SO)(H)₃ \equiv C-(CO)(H)₃, assigned. ^e C_B-(SO) \equiv C_B-(CO), assigned. ^f C-(SO)₂(H)₃ \equiv C-(SO)(H)₃, assigned. ^g C_B-(SO)₂ \equiv C_B-(CO), assigned. ^h SO₂-(C_d)(C_B) \equiv SO₂-(C_B)₂, assigned. ⁱ CO-(S)(C) \equiv CO-(C)₂, assigned. ^j CS-(N)₂ \equiv CO-(C)₂, assigned. ^k S-(S)(N) \equiv O-(O)(C), assigned. ^l SO-(N)₂ \equiv CO-(C)₂, assigned. ^m SO₂-(N)₂ \equiv SO-(N)₂, assigned. ⁿ Assume ring corrections for g and f are the same. ^o Assume ring corrections for i and c are the same. ^p No *gauche* corrections across the X-P, X-PO, and X-P:N bonds (X represents C, O, N). ^q *gauche* corrections across the C-B bond are +0.8 kcal/mole. ^r No *gauche* corrections across C-M bond.