

STEREOCHEMICAL STUDIES OF NITROGEN BRIDGEHEAD COMPOUNDS BY SPECTRAL MEANS


T. A. CRABB, R. F. NEWTON, AND D. JACKSON

Department of Chemistry, Portsmouth Polytechnic, Hampshire, England

Received March 20, 1970 (Revised Manuscript Received July 13, 1970)

Contents

I. Introduction	109
A. Scope	109
B. Quinolizidine	110
II. Spectroscopic Methods	110
A. Infrared	110
B. Nuclear Magnetic Resonance	111
III. Stereochemical Determinations	112
A. Fused 6/6 Ring Systems with No Additional Heteroatom	112
B. Fused 6/6 Ring Systems with Additional Heteroatoms	118
1. One Additional Oxygen Atom	119
2. One Additional Nitrogen Atom	120
3. One Additional Sulfur Atom	121
4. Two Nitrogen Atoms at a Ring Fusion	121
C. Fused 6/5 Ring Systems with No Additional Heteroatom	121
1. Indolizidine	121
2. Indolizidine Methiodides	123
D. Fused 6/5 Ring Systems with Additional Heteroatoms	123
E. Fused 5/5 Ring Systems	125



I. Introduction

A. SCOPE

Quinolizidine (1), indolizidine (2), and pyrrolizidine (3) form the nucleus of many compounds of biological and medicinal interest. For example, the quinolizidine moiety occurs in matrine (4), the yohimbine group of alkaloids (5), and the veratrum alkaloid cevine (6), while the indolizidine nucleus is found in the skeleton of the solanum alkaloid solanidine (7) and the veratrum alkaloid rubijervine. The necine alkaloids, *e.g.*, laburnine (8), are built around the pyrrolizidine ring. The stereochemistry of these classes of alkaloids will not be dealt with here since this has been reviewed elsewhere.¹⁻⁵

Heterocyclic systems with bridgehead nitrogen atoms have been the subject of an extensive review,⁶ but since this was published a considerable number of papers have appeared

dealing with stereochemical studies of saturated heterocyclic systems, and it is this aspect of the subject which forms the basis of the present review. In particular, the stereochemistry of the simpler derivatives of 1, 2, and 3 together with related compounds possessing additional heteroatoms will be discussed with reference to spectroscopic methods of determining their configurations and preferred conformations. The bulk of the data will refer to the establishment of the *cis* or *trans* nature of the fusion between the two rings, but evidence for the configurations, *i.e.*, the *cis* or *trans* relationship of a substituent with respect to the bridgehead hydrogen, will also be presented. The extensive work on kinetic methods of determining stereochemistry has not been reviewed and will be mentioned only to provide the necessary background to a discussion of certain groups of compounds. Spectroscopic and stereochemical data relating to bridged systems such as quinuclidine are not included in this review.

(1) G. A. Morrison, *Fortschr. Chem. Org. Naturst.*, **25**, 269 (1967).
 (2) N. K. Kochetkov and A. M. Likhoshester, *Advan. Heterocycl. Chem.*, **5**, 315 (1965).
 (3) R. H. F. Manske and H. L. Holmes, "The Alkaloids," Academic Press, New York, N. Y., 1950.
 (4) E. L. Eliel, N. L. Allinger, S. J. Angyal, and G. A. Morrison, "Conformational Analysis," Interscience, New York, N. Y., 1965.
 (5) L. B. Bull, C. C. J. Culvenor, and A. T. Dick, "The Pyrrolizidine Alkaloids," North-Holland Publishing Co., Amsterdam, 1969.
 (6) W. L. Mosby, "Heterocyclic Systems with Bridgehead Nitrogen Atoms," Interscience, New York, N. Y., 1961.

B. QUINOLIZIDINE

The major difference between quinolizidine and its carbocyclic analog is the replacement of the axial C-H bond of the bridgehead carbon atom of the latter by the nitrogen lone pair of electrons in the former. The presence of the conformationally mobile nitrogen atom at the bridgehead permits ready interconversion between the trans-fused and cis-fused conformations (Figure 1). Barton and Cookson⁷ predicted that the





Figure 1. Equilibrium between *trans*- and *cis*-quinolizidine.

nitrogen lone pair would have a steric requirement approaching that of a methyl group, an idea supported by several experimental results.^{8,9} Other workers, however, have interpreted these observations in terms of a small steric requirement for the nitrogen lone pair. Many of the papers relevant to this topic were mentioned in a recent review,¹⁰ and the subject demands a separate treatment which falls outside the scope of the present work. The consensus of opinion now favors a negligible steric requirement for the nitrogen lone pair except under certain conditions, and, assuming this to be the case, Aaron¹¹ estimated that quinolizidine is more stable in the trans-fused ring conformation relative to the cis-fused ring conformation than is decalin by *ca.* 2.1 kcal/mol. This conclusion was arrived at by attributing the energy difference between the systems to the two nonbonded 1,3-diaxial hydrogen interactions (2×1.04 kcal/mol)¹² which destabilize *trans*-decalin relative to *cis*-decalin and which disappear when the bridgehead C-H is replaced by a tertiary nitrogen atom in quinolizidine. Thus *trans*-quinolizidine is estimated to be more stable than *cis*-quinolizidine by *ca.* 4.6 kcal/mol. Recently Aaron¹³ has revised his estimate of ΔG° for the quinolizidine equilibrium and shown that it should be similar to the free energy difference between *cis*- and *trans*-decalin, since in the original paper only 1,3-diaxial hydrogen-hydrogen interactions were considered, the comparable 1,2-skew interactions being neglected. From a study of the kinetics of quaternization of monomethylquinolizidines using hexahydrojulolidines (9) as fixed models of the *cis*- and *trans*-fused ring conformations, Katritzky, Schofield, and coworkers¹⁴ deduced that ΔG° for



the *cis* \rightleftharpoons *trans* quinolizidine equilibrium was *ca.* -4.4 kcal/mol.

II. Spectroscopic Methods

A. INFRARED

Chemical methods of assigning stereochemistry are often tedious and several physical criteria have therefore been developed. The most widely used physical method has no doubt been infrared spectroscopy, and the stereochemistry and conformational preferences of a large number of quinolizidines and their derivatives have been assigned by this means.

The first correlation between the configuration about the ring fusion of a nitrogen bridgehead compound and the appearance of bands in the 2800-2700-cm⁻¹ region of the infrared spectrum was made by Wenkert¹⁵ in connection with studies on the stereochemistry of the yohimbines and related alkaloids. Comparison of the spectra of various pairs of C-3 epimers showed that all compounds possessing an α hydrogen at C-3 gave rise to two or more distinct peaks of medium intensity on the low wave number side of the symmetric C-H stretching band. Compounds possessing a 3β hydrogen showed only shoulders on the low wave number side of the main peak at 2860 cm⁻¹. During structural investigations of a large number of alkaloids containing the quinolizidine nucleus, Bohlmann¹⁶ found that those in which the quinolizidine moiety was trans-fused exhibited a characteristic series of bands in the infrared spectrum between 2700 and 2800 cm⁻¹; these bands, which he termed "trans bands," were absent when the quinolizidine nucleus was cis-fused. In order to investigate this phenomenon further, a number of simple quinolizidines and deuterated quinolizidines were synthesized, and it was shown¹⁷ that the appearance of the bands was due to the axial C-H bonds α to the nitrogen atom and trans to the nitrogen lone pair of electrons. Two such C-H bonds were found to be necessary for the bands to be observed; in cis-fused quinolizidines, only one such bond is present. Rosen¹⁸ and Becket¹⁹ have modified Wenkert's original infrared correlation, since only those alkaloids possessing in their preferred conformation(s) the C-3-H and at least one more C-H bond trans diaxial to the nitrogen lone pair will exhibit bands in the infrared spectrum between 2860 and 2700 cm⁻¹, while those alkaloids possessing in their preferred conformation(s) the C-3-H cis to the nitrogen lone pair will not. Wiewiorowski and Skolik²⁰ have undertaken a more quantitative study of the 2900-2500-cm⁻¹ region of the

(7) D. H. R. Barton and R. C. Cookson, *Quart. Rev., Chem. Soc.*, **10**, 44 (1956).

(8) M. J. Aroney and R. J. W. Le Fèvre, *J. Chem. Soc.*, 3002 (1958).

(9) M. J. Aroney, C. Y. Chen, R. J. W. Le Fèvre, and J. D. Saxby, *ibid.*, 4269 (1964).

(10) F. G. Riddell, *Quart. Rev., Chem. Soc.*, 364 (1967).

(11) H. S. Aaron, *Chem. Ind (London)*, 1338 (1965).

(12) L. F. Fieser and M. Fieser, "Advanced Organic Chemistry," Reinhold, 1961, p 559.

(13) H. S. Aaron and C. P. Ferguson, *Tetrahedron Lett.*, 6191 (1968).

(14) C. D. Johnson, R. A. Y. Jones, A. R. Katritzky, C. R. Palmer, K. Schofield, and R. J. Wells, *J. Chem. Soc.*, 6797 (1965).

(15) E. Wenkert and D. Roychaudhuri, *J. Amer. Chem. Soc.*, **78**, 6417 (1956).

(16) F. Bohlmann, *Angew. Chem.*, **69**, 541 (1957); *Chem. Ber.*, **91**, 2157 (1958).


(17) F. Bohlmann, D. Schumann, and M. Schulz, *Tetrahedron Lett.*, 173 (1964).

(18) (a) W. E. Rosen, *ibid.*, 481 (1961); (b) W. E. Rosen and J. N. Shoolery, *J. Amer. Chem. Soc.*, **83**, 4816 (1961).

(19) W. F. Trager, C. M. Lee, and A. H. Becket, *Tetrahedron*, **23**, 365 (1967).

(20) M. Wiewiorowski and J. Skolik, *Bull. Acad. Pol. Sci., Ser. Sci. Chim.*, **10**, 1 (1962).

infrared spectra of the C-15 lupin alkaloids. Accordingly, the infrared spectra of 15 alkaloids of the sparteine series were examined under identical conditions and the following conclusions were drawn. The area within which "trans bands" (termed "T bands" by these authors) occur is from just below 2800–2700 cm^{-1} as suggested by Bohlmann. Of the compounds examined, dilactams of type **10** and the majority of neutral



10

salts were the only compounds with no T bands, all the other compounds exhibiting a series of T bands from just below 2860 to 2500 cm^{-1} . The intensity of the T band was expressed in terms of the integrated area of the peak, and using this system it was found that the T-band area for a *cis*-quinolizidine system was normally only about 20% smaller than that for a *trans*-quinolizidine system. In *cis*-quinolizidines, however, the T bands were present as a stepwise descending set of maxima below the C–H stretching band at 2860 cm^{-1} , whereas in those compounds containing a *trans*-fused quinolizidine ring system, the main T bands were present as well-defined maxima at about 2800 and 2750 cm^{-1} . During a more recent study by the same authors²¹ on a number of sparteine-type quinolizidine alkaloids and their selectively α -deuterated derivatives, the following points emerged. It was suggested that the lower frequency of the T bands compared with the normal C–H region was due to charge delocalization of the nitrogen lone pair of electrons to the α axial C–H bonds. In *trans*-fused quinolizidines all three α axial C–H bonds would take part in this delocalization. Since the axial C–H bonds on both C-4 and C-6 have the same symmetry and force constants, vibrational coupling can occur and two bands appear at 2800 and 2761 cm^{-1} , respectively. The first is due to the asymmetric and the second to the symmetric stretching vibration. In *cis*-fused quinolizidines both the chair–chair and the chair–boat conformations were considered. The chair–chair conformation has only one α axial C–H bond, and this gives rise to a single band between 2840 and 2600 cm^{-1} . In the chair–boat conformation three α axial C–H bonds are present and give rise to two main bands at *ca.* 2808 and 2761 cm^{-1} . In a further paper by Wiewiorowski, *et al.*,²² the conditions necessary for the occurrence of Bohlmann bands were more precisely defined. It was found that in order for bands to occur, one hydrogen on a carbon atom alpha to a nitrogen atom situated *trans* and axial with respect to the nitrogen lone pair of electrons was necessary. The intensity and complexity of the absorption was found to be roughly proportional to the number of hydrogen atoms so situated. There have been several papers^{23,24} dealing with the theoretical aspects of "Bohlmann bands," and it has been suggested that they are similar in origin to the characteristic *N*-methyl bands due to Fermi resonance which appear in the same region.

B. NUCLEAR MAGNETIC RESONANCE

Next to infrared spectroscopy, the most widely used physical method in conformational studies on bridgehead nitrogen systems has been nmr spectroscopy. In an investigation of the nmr spectrum of quinolizidine, Hamlow²⁴ identified the signals at τ 7.2–7.4 as due to the C-4 and C-6 equatorial protons, and a broad shoulder at τ 8.0 on the side of the main peak as due to the corresponding axial protons. The accurate chemical shifts of these protons were determined by spin decoupling, and it was found that the chemical shift difference between the C-4, C-6 axial and equatorial protons was 0.93 ppm. When the nmr spectrum of quinolizidine was run in trifluoroacetic acid, a protonating solvent, the equatorial protons absorbed at about τ 6.5 and the axial protons at τ 7.0, the chemical shift difference of 0.5 ppm being comparable to that observed for the axial and equatorial protons of cyclohexane. In order to explain the difference in chemical shift between the methylene protons adjacent to nitrogen in quinolizidines, it was suggested that partial participation of the nitrogen lone pair in a σ^* C–H_{ax} orbital on the adjacent carbon atom occurs, and this leads to an increase in the electron density at the axial proton. The chemical shift difference should therefore be greatest when the nitrogen lone pair and the adjacent C–H bond have a *trans*-diaxial relationship with each other. The theory that the nitrogen lone pair of electrons is responsible for shielding the axial proton is supported by the decreased shielding of this proton in a protonating solvent. Hamlow²⁴ has suggested that it is the decrease in the bonding energy of the axial C–H bonds adjacent to the nitrogen atoms which is responsible for the appearance of bands in the 2800–2700- cm^{-1} region of the infrared spectra of quinolizidine. An explanation of the anti-axial shielding effect of a proton by a lone pair in terms of a dipole effect has also been offered.²⁵

Bohlmann¹⁷ showed that Hamlow's assignments for the C-4 protons in quinolizidine were correct by investigating the spectra of a range of deuterated and methyl-substituted quinolizidines. He found that the spectrum of quinolizidine exhibited a two-proton doublet at τ 7.2 and a two-proton multiplet at τ 7.8 to 8.2. Replacement of both the C-4 protons by deuterium reduced the intensity of each of these signals to one proton, and replacement of both the C-4 and C-6 protons by deuterium led to the complete disappearance of the signals. The nmr spectrum of *cis*-4,10-*H*-4-methylquinolizidine, which was assumed to exist predominantly in the *trans*-fused ring conformation, only showed a one-proton signal to low field at τ 6.85. When this 4-methyl compound was deuterated at the C-4 axial position, the nmr spectrum showed no change in the low-field signal.

Recently an alternative explanation for the large Δ value in quinolizidine has been put forward by Robinson,²⁶ who points out that previous workers²⁴ assumed that an equatorial methylene group on the nitrogen atom would have little effect on the chemical shift of protons on adjacent carbon atoms, whereas in fact the available unambiguous evidence^{27–29} suggests that a vicinal axial hydrogen is shielded by an equatorial methyl or methylene group. The large chemical shift difference between

(21) J. Skolik, P. J. Krueger, and M. Wiewiorowski, *Tetrahedron*, **24**, 5439 (1968).

(22) M. Wiewiorowski, O. E. Edwards, and M. D. Bratek-Wiewiorowska, *Can. J. Chem.*, **45**, 1447 (1967).

(23) T. Masamune and M. Takasugi, *Chem. Commun.*, 625 (1967).

(24) H. P. Hamlow, S. Okuda, and N. Nakagawa, *Tetrahedron Lett.*, 2553 (1964).

(25) R. F. Zürcher, "Progress in NMR Spectroscopy," Pergamon Press, London, 1967, Chapter 5.

(26) M. J. T. Robinson, *Tetrahedron Lett.*, 1153 (1968).

(27) J. B. Lambert, R. S. Keske, R. E. Carhart, and A. P. Jovanovich, *J. Amer. Chem. Soc.*, **89**, 3761 (1967).

(28) A. Segre and J. I. Musher, *ibid.*, **89**, 706 (1967).

(29) N. W. J. Pumphrey and M. J. T. Robinson, *Chem. Ind. (London)*, 1903 (1963).

protons adjacent to the nitrogen atom in *trans*-quinolizidine can then be explained in terms of shielding of the axial proton by the methylene group on the nitrogen atom, rather than shielding caused by the *trans* nitrogen lone pair. The relative importance of the two shielding mechanisms has recently been discussed by Lambert.³⁰



III. Stereochemical Determinations

A. FUSED 6/6 RING SYSTEMS WITH NO ADDITIONAL HETEROATOM

1. Quinolizidines

a. Methylquinolizidines

All the isomeric monomethylquinolizidines have been subjected to a thorough investigation^{14,31,32} and are now known to exist predominantly in the *trans*-fused ring conformation. The configurations and preferred conformations were assigned to the compounds on the basis of chemical and spectroscopic evidence. All the compounds, with the exception of the *trans*-4,10-*H*-4-methyl derivative (11), showed a prominent band at 2760–2750 cm^{-1} together with other bands in the Bohlmann region of the infrared and were therefore assigned *trans*-fused conformations. The center of the methyl doublet and the apparent $J_{\text{CH-Me}}$ was an excellent indication of the equatorial or axial orientation of the methyl group, it being found that the axial methyl group absorbed at lower field and showed a larger “*J*” than its corresponding epimer. This did not apply in the case of the 4-methyl compound mentioned above which was assigned a *cis*-ring fusion with an equatorial methyl group on the basis of the infrared and nmr evidence.



In a later paper,¹⁴ in which the rates of quaternization of the monomethylquinolizidines and of the hexahydrojulolidines were measured and compared, it was shown that this latter assignment was incorrect and that, in fact, *trans*-4,10-*H*-4-methylquinolizidine exists in a predominantly *trans*-fused ring conformation (12) at room temperature. It should be mentioned that there is a strong band at 2805 cm^{-1} in the infrared spectrum of this compound which is in fact probably a *trans* band. It seems therefore that the Bohlmann criterion must be used with caution when dealing with 4-substituted quinolizidines.

The configurations of a 4,7-dimethyl and a 1,4,7-trimethyl quinolizidine have been assigned³³ and both compounds shown to exist in *trans*-fused ring conformations. The chemical shifts of the methyl groups are in accord with the observations on the monomethylquinolizidines (Table I).

Table I

Nmr and Ir Data on Monomethylquinolizidines³²

Compound	Nmr methyl doublet Center of methyl doublet (τ)	Apparent J (Hz)	Ir Bohlmann region (cm^{-1})
	9.05	5.8	2805 2755 2680 2610
	9.16	1.6	2805 2755 2685 2605
	9.09	2.7	2805 2760 2675 2605
	9.01	6.9	2800 2760 2675 2600
	8.92	6.6	2795 2755 2680 2615
	9.18	4.8	2805 2760 2685 2625
	9.00	5.1	2795 2760 2625
	9.07	4.6	2805

b. Hydroxyquinolizidines

This has been an attractive field to explore because of the potential biological activity of these compounds. Particular use has been made of infrared spectroscopy in assigning configurations which in many cases can be made on the basis of the presence or absence of intramolecular H-bonded OH frequencies.

Aaron³⁴ has studied the isomeric 1-, 2-, and 3-hydroxyquinolizidines (Table II), all of which showed Bohlmann bands and therefore exist in *trans*-fused ring conformations. The configurations of the 1- and 3-hydroxy compounds were deduced from the presence of a band at 3526 cm^{-1} in the infrared spectra of the axial-hydroxy epimers of the 1- and 3-hydroxy compounds indicative of intramolecular hydrogen bonding. Configurational assignments were made on the two 2-hydroxyquinolizidines by studying the C-2 proton signals in the nmr spectra. In all the six alcohols the carbinol CH proton in the equatorially substituted hydroxy compound absorbed at higher field and gave broader signals than that observed in the axial-hydroxy epimers.

(30) J. B. Lambert and R. G. Keske, *Tetrahedron Lett.*, 2023 (1969).

(31) T. M. Moynahan and K. Schofield, *Proc. Chem. Soc. London*, 218 (1961).

(32) T. M. Moynahan, K. Schofield, R. A. Y. Jones, and A. R. Katritzky, *J. Chem. Soc.*, 2637 (1962).

(33) M. Kotake, I. Kawasaki, T. Okamoto, S. Maturani, S. Kasumoto, and T. Kaneko, *Bull. Chem. Soc. Jap.*, 35, 1337 (1962).


(34) H. S. Aaron, G. E. Wicks, Jr., and C. P. Rader, *J. Org. Chem.*, 29, 2248 (1964).

Table II
Ir Data on Hydroxyquinolizidines

Compound	Bohlmann region (cm ⁻¹)	Intramolecularly bonded		Ref	Compound	Bohlmann region (cm ⁻¹)	Intramolecularly bonded		Ref
		Free OH (cm ⁻¹)	OH (cm ⁻¹)				Free OH (cm ⁻¹)	OH (cm ⁻¹)	
	2800	...	3526	34		2800-2700	...	Intra	38
	2777	...	3526	34		2800-2700	...	Intra	38
	2795	3618	...	34		2800-2700	...	Intra	38
	2750	3618	...	34		2800-2700	...	Intra	38
	2802	3625	...	34, 35		2800-2700	3620	...	38
	2764	3625	...	34, 35		2800-2700	3620	...	38
	2801	3620	...	34, 35		2800-2700	3620	...	38
	2762	3620	...	34, 35		2800-2700	3620	...	38
	2797	...	3527	34		2800-2700	Free OH	...	35
	2757	...	3527	34		2800-2700	Free OH	...	35
	2800	3609	...	34		2800-2700	Free OH	...	35
	2762	3609	...	34		2800-2700	Free OH	...	35
	2800-2700	...	3480	36		Prominent bands 2800-2600	3620	...	39
	2800-2700	3620	...	36		Prominent bands 2800-2600	...	3520	39
	2820	...	3480	37		2800-2600	3605	None	40
	2770	...	3480	37		2800-2600	3605	3350	40
	2810	3620	...	37		2800-2600	3605	3350	40
	2770	3620	...	37		2800-2600	3605	3350	40

Other hydroxyquinolizidines (Table II) have been examined³⁵⁻⁴⁰ by similar techniques and exist predominantly in the conformations shown. Recently the infrared spectra of *trans*-1,10-*H*-1-hydroxyquinolizidine has been studied¹³ using long-path-length cells and 2% of the *cis*-fused conformation detected in the equilibrium mixture. Reexamination of 1-hydroxy-1-(phenyl *trans*-10-*H*)-quinolizidine shows it to exist

as a conformational mixture of 60% *trans*-fused (13) and 40% *cis*-fused ring conformations (14).



In the two 1-hydroxy-1-methylquinolizidines³⁶ the equatorial methyl group in one isomer absorbed at τ 8.91 and the axial methyl in the epimer at τ 8.78 in agreement with the data presented in Table I.

Möhrle^{38,41} has studied the 1-hydroxy-7-methyl- and the

(35) F. Bohlmann, E. Winterfeldt, O. Schmidt, and W. Reusche, *Chem. Ber.*, **94**, 1767 (1961).

(36) D. Temple and J. Sam, *J. Heterocycl. Chem.*, **5**, 441 (1968).

(37) J. D. England and J. Sam, *ibid.*, **3**, 482 (1966).

(38) H. Möhrle and C. Karl, *Arch. Pharm. (Weinheim)*, **301**, 530 (1968).

(39) R. van Hes, U. K. Pandit, and H. O. Huisman, *Recl. Trav. Chim. Pays-Bas*, **86**, 1255 (1967).


(40) J. Sam, J. D. England, and D. Temple, *J. Med. Chem.*, **12**, 144 (1969).

Table III
Nmr of 1-Hydroxy-7-methyl- and
1-Hydroxy-1,7-dimethylquinolizidines³⁸


Compound	No.	R = H	R = Me
	15	7 Me (ax) τ 8.9 ($J = 7$ Hz)	7 Me (ax) τ 8.92 ($J = 7.1$ Hz) 1 Me (eq) τ 8.92
	16	7 Me (eq) τ 9.17 ($J = 5.8$ Hz)	7 Me (eq) τ 9.16 ($J = 6$ Hz) 1 Me (eq) τ 8.89
	17	7 Me (ax) τ 8.91 ($J = 7$ Hz)	7 Me (ax) τ 8.92 ($J = 7$ Hz) 1 Me (ax) τ 8.75
	18	7 Me (eq) τ 9.16 ($J = 5.7$ Hz)	7 Me (eq) τ 9.16 ($J = 5.9$ Hz) 1 Me (ax) τ 8.78

1-hydroxy-1,7-dimethylquinolizidines. All were assigned predominantly *trans*-fused ring conformations on the basis of the presence of Bohlmann bands in the infrared, and use was made of the results of hydrogen-bonding studies, methyl chemical shift data, and the values of " $J_{\text{CH-Me}}$ " in assigning configurations (Table III). Isomers **17** and **18** showing intramolecular hydrogen bonding were assigned structures with the axial-hydroxy groups.

The four isomers of **19** have been obtained⁴² and their con-




figurations and conformations assigned on the basis of the Bohlmann region of the infrared spectra and the chemical shifts and band widths of the carbinol protons. The epimeric 2-hydroxy-2-phenylquinolizidines have been synthesized⁴⁰ and configurations and conformations assigned on the basis of the chemical shifts of the methyl groups of the corresponding acetates. One of the isomers (**20**) shows intramolecular hydrogen bonding at low concentration and this is attributed to the boat form (**21**).




c. Arylquinolizidines

Bohlmann⁴³ has reported the nmr spectra of *cis*- and *trans*-4,10-*H*-4-phenylquinolizidines. The spectrum of *cis*-4,10-*H*-4-

phenylquinolizidine (**22**, R = Ph) showed a two-proton




multiplet at τ 7.3 which was assigned to the equatorial C-6 proton and the axial C-4 proton since the latter should be deshielded by the phenyl group on C-4. The spectrum of *trans*-4,10-*H*-4-phenylquinolizidine, which had been assigned a *cis*-fused ring conformation on the basis of the lack of bands between 2800 and 2700 cm^{-1} in its infrared spectrum, showed a one-proton signal at τ 6.17 which was assigned to the C-4 proton, a signal at τ 7.00 which was assigned to the C-10 proton, and peaks at τ 7.37 and 7.61 assigned to the equatorial and axial C-6 protons, respectively. The correctness of these assignments was proved by deuteration experiments. From these figures it was deduced that *trans*-4,10-*H*-4-phenylquinolizidine exists in a predominantly *cis*-fused ring conformation (**24**, R = Ph). Thus the C-4 proton comes to low field



due to the 1,3-deshielding effects acting upon it across the ring fusion, and there is only a small chemical shift difference between the C-6 protons since in the *cis*-fused ring conformation the lone pair of the nitrogen atom bisects the C-6 methylene group. In addition the C-10 proton comes to low field since it is no longer *trans* and axial with respect to the nitrogen lone pair.

The isomers of 4-(α -furyl)- and 4-(α -thiophenyl)quinolizidine have been prepared.⁴⁴ One furanyl isomer showed bands in the infrared at 2780 and 2740 cm^{-1} , and a thienyl isomer showed similar bands at 2795 and 2760 cm^{-1} . These compounds were therefore assigned the *trans*-fused ring conformation **22**. The two remaining isomers showed no Bohlmann bands and were therefore considered to have adopted the *cis*-fused ring conformation **24**.

Deoxynupharidine (**25**) has the stereochemistry shown³³. It



has ⁴⁵⁻⁴⁷ bands in the 2700–2800- cm^{-1} region of the infrared and τ values (benzene solution) for the axial 7-methyl of 8.90 and for the equatorial 1-methyl of 9.11. The 7-*epi* compound shows methyl peaks at τ 9.14 and 9.32 (equatorial methyl), and the 1-*epi* compound shows both axial methyls to absorb at τ 8.81 and 8.91.

(41) H. Möhrle, C. Karl, and U. Scheidegger, *Tetrahedron*, **24**, 6813 (1968).

(42) T. Matsunaga, I. Kawasaki, and T. Kaneko, *Tetrahedron Lett.*, 2471 (1967).

(43) F. Bohlmann, D. Schumann, and C. Arndt, *ibid.*, 2705 (1965).

(44) K. Kotera, Y. Hamada, and R. Mitsui, *Tetrahedron*, **24**, 2463 (1968).

(45) Y. Arata, *Chem. Pharm. Bull. Jap.*, **13**, 907 (1965).


(46) Y. Arata, N. Hazama, and Y. Kojima, *Yakugaku Zasshi*, **82**, 326 (1962).

(47) I. Kawasaki, S. Matsutani, and T. Kaneko, *Bull. Chem. Soc. Jap.*, **36**, 1474 (1963).

A number of diphenyl- and phenylmethyl-substituted quinolizidines have been prepared⁴⁸ and assigned *cis*- and *trans*-fused ring conformations on the basis of Bohlmann bands.

d. Miscellaneous Quinolizidines

Both epimers of 4-hydroxymethylquinolizidine show Bohlmann bands in the infrared and are therefore assigned⁴⁹ the *trans*-ring fusion **22** ($R = CH_2OH$) (2750, 2780 cm^{-1}) and **23** ($R = CH_2OH$) (2790 and 2770 cm^{-1}), whereas only one of the 4-carbomethoxyquinolizidines shows Bohlmann bands (2800, 2770 cm^{-1}); the second isomer must therefore adopt the *cis*-fused conformation **24** ($R = COOEt$). Lamprolobine (**26**)⁵⁰ and lusitanin (**27**)⁵¹ both show *trans* bands.




The diastereoisomers of 3-(4-chlorobenzyl)quinolizidine were⁵² considered to possess *trans*-ring fusions because of the presence of Bohlmann bands in the infrared. The axial-equatorial nature of the substituent was confirmed by the position of the $^+N-CH_3$ signals in the nmr of the corresponding methiodides.

e. Oxoquinolizidines


Bohlmann bands are shown by 1-oxo- (2795, 2750, 2680 cm^{-1}), 2-oxo- (2785, 2740, 2680 cm^{-1}), and 3-oxo- (2780, 2735, 2645 cm^{-1}) quinolizidine, and circular dichroism measurements indicate that in isoctane solutions at room temperature the *trans* conformation represents 90% of the conformational equilibrium.^{53,54}

The lactam **28** is considered⁵⁵ to exist in the conformation shown. The C-6 and C-10 protons all absorb at lower field than in quinolizidine and the C-6 equatorial proton lying in the amide plane is particularly deshielded by the amide carbonyl (τ 5.37 compared to τ 7.2 in quinolizidine). The spectra of two diastereoisomeric lactam carbinols **29**, degradation products of securinine, have been described.^{56,57}




f. Cycloalkanoquinolizidines


Interest in matrine has stimulated research into the synthesis and stereochemistry of cycloalkanoquinolizidines. The most carefully studied systems have been the hexahydrojulolidines (**9**)^{14,58,59} (three possible stereoisomers **30**–**32**), the correspond-



ing cyclopentanoquinolizidines^{60–62} (**30**–**32** with ring C five-membered), **33** and **34**, and the perhydronaphoquinolizidines (**35**).⁶³ The three diastereoisomeric hexahydrojulolidines have



been shown to possess the stereochemistry shown by structures **36**, **37**, and **38**. *cis,trans*-Hexahydrojulolidine (**31**) was



readily distinguished⁵⁸ from **30** and **32** since it showed only weak bands in the 2800–2700- cm^{-1} region of the infrared. The *cis,cis* isomer **36** possesses a nitrogen lone pair of electrons shielded from an attacking reagent by the C ring and accordingly was found¹⁴ to react only slowly with methyl iodide. In fact, no reaction⁵⁹ occurred on refluxing **32** with methyl iodide in ether whereas the *trans,trans* isomer **30** reacted readily under the same conditions. By these techniques and by studying their rate of reaction⁵⁸ with mercuric acetate, the three isomers were characterized. The stereochemistry of the three isomeric

(48) F. Bohlmann, E. Winterfeldt, G. Boroschewski, R. Mayer-Mader, and B. Gatscheff, *Chem. Ber.*, **96**, 1792 (1963).

(49) Y. Arata, Y. Asaoka, and M. Kassai, *Yakugaku Zasshi*, **82**, 1523 (1962).

(50) S. R. Johns, N. K. Hart, and J. A. Lambertson, *Aust. J. Chem.*, **21**, 1619 (1968).

(51) K. Wicky and E. Steinegger, *Pharm. Acta Helv.*, **40**, 658 (1965).

(52) I. Matsuo, K. Sugimoto, and S. Ohki, *Chem. Pharm. Bull. Jap.*, **16**, 1680 (1966).

(53) S. F. Mason, K. Schofield, and R. J. Wells, *Proc. Chem. Soc. London*, 337 (1963).

(54) S. F. Mason, G. W. Vane, K. Schofield, R. J. Wells, and J. S. Whitehurst, *J. Chem. Soc. B*, 553 (1967).

(55) F. Bohlmann and D. Schumann, *Tetrahedron Lett.*, 2435 (1965).

(56) S. Saito, K. Kotera, N. Shigematsu, A. Ide, N. Sugimoto, Z. Horii, M. Hanaoka, Y. Yamawaki, and Y. Tamura, *Tetrahedron*, **19**, 2085 (1963).

(57) Z. Horii, Y. Yamawaki, M. Hanaoka, Y. Tamura, S. Saito, and H. Yoshikawa, *Chem. Pharm. Bull. Jap.*, **13**, 22 (1965).

(58) F. Bohlmann and C. Arndt, *Chem. Ber.*, **91**, 2167 (1958).

(59) K. Tsuda and S. Saeki, *Chem. Pharm. Bull. Jap.*, **6**, 391 (1958).


(60) K. Schofield and R. J. Wells, *J. Chem. Soc.*, 621 (1967).

(61) L. Mandell, J. U. Piper, and K. P. Singh, *J. Org. Chem.*, **28**, 3440 (1963).

(62) N. J. Leonard and W. J. Middleton, *J. Amer. Chem. Soc.*, **74**, 5114 (1952).

cylopentanoquinolizidines (30–32, ring C five-membered) has been established⁶⁰ by similar techniques, the *trans,trans* compound showing weaker Bohlmann bands than does *trans,trans*-hexahydrojulolidine. Both 33 and 34 (tentative assignment) show prominent Bohlmann bands.⁶¹


Four isomers of the perhydronaphthoquinolizidine 35 have been obtained and their stereochemistry deduced⁶² as 39–42 on the basis of their rates of methylation, dehydrogenation, reaction with aluminum chloride, and from the appearance of Bohlmann bands in the ir spectra of all four isomers indicating a *trans* A:B ring fusion. 39 and 42 were readily methylated whereas reaction occurred only slowly with 40 and not at all



with the isomer 41 in which the nitrogen lone pair is shielded from attack. 40 and 42 were converted to the more stable isomer 39 on treatment with aluminum chloride.


An 8-hydroxy derivative of 32 was characterized⁶⁴ by the appearance of an intramolecularly hydrogen-bonded OH frequency in the infrared. Pairs of various epimeric 3-substituted derivatives of 32 and 30 have been assigned⁶⁵ their structures from their manifestation of Bohlmann bands and from the chemical shifts of the protons on carbon atoms α to the nitrogen.

The three isomers of perhydrobenzo[*c*]quinolizine (43) have been assigned⁶⁶ the stereostructures 43a–c on the basis of the




presence or absence of Bohlmann bands in the infrared, their method of synthesis, and their rates of methylation ($c > b > a$).

Infrared studies led to the assignment⁶⁷ of structures 44 and 45 to the hydroxy compounds related to securinine. 44 and 46 showed Bohlmann bands, and 45 and 46 showed absorption due to intramolecularly hydrogen-bonded hydroxy groups at 3505 and 3509 cm^{-1} , respectively. The free OH stretching frequency of 44 was at 3620 cm^{-1} . The *cis*-fused




quinolizidine moiety has been shown⁶⁸ to be present in 47 by the absence of Bohlmann bands, and the presence of a hydrogen-bonded hydroxy group is shown by the absorption in the infrared at 3503 cm^{-1} .

The stereochemistry of various hydroxylated derivatives of 48 and related compounds has been studied by Kupchan.^{69–72}



The pair of compounds 49 and 50⁷³ provide excellent examples of the use of Bohlmann's infrared criterion in making stereochemical assignments in compounds related to the Lycopodium alkaloids, 49 showing marked bands which are only hinted at in 50.




g. Benzoquinolizidines

In conformational studies on benzoquinolizidines of type 51 ($R = H, R' = H$), the chemical shift of the angular C-11b proton has been of considerable use. It was originally found that in methyl neoreserpat¹⁸ (which adopts a conformation 53


(63) S. Saeki, *Chem. Pharm. Bull. Jap.*, **9**, 226 (1961).
 (64) F. Bohlmann and O. Schmidt, *Chem. Ber.*, **97**, 1354 (1964).
 (65) F. Bohlmann, D. Habeck, E. Poetsch, and D. Schumann, *ibid.*, **100**, 2742 (1967).
 (66) S. Ohki, M. Akiba, H. Shimada, and K. Keenihira, *Chem. Pharm. Bull. Jap.*, **16**, 1889 (1968).

(67) Z. Horii, M. Ikeda, Y. Yamawaki, Y. Tamura, S. Saito, and K. Kodera, *Tetrahedron*, **19**, 2101 (1963).
 (68) J. Parello, *Bull. Soc. Chim. Fr.*, 1117 (1968).
 (69) S. M. Kupchan, G. F. Flouret, and C. A. Matuszak, *J. Org. Chem.*, **31**, 1707 (1966).
 (70) S. M. Kupchan, A. D. J. Balon, and C. G. DeGrazin, *ibid.*, **31**, 1713 (1966).
 (71) S. M. Kupchan and C. G. DeGrazin, *ibid.*, **31**, 1716 (1966).
 (72) S. M. Kupchan, J. H. Block, and A. C. Isenberg, *J. Amer. Chem. Soc.*, **89**, 1189 (1967).
 (73) W. A. Ayer and G. G. Iverach, *Can. J. Chem.*, **42**, 2514 (1964).



51


such that the angular proton and the nitrogen lone pair are trans diaxial) the angular proton absorbed at τ 6.8 whereas the angular proton in methyl reserpate (**52**) which is gauche to



52

53


the lone pair absorbed at τ 6.2. A similar difference in chemical shifts was observed¹⁵ for ajmalicine and 3-isoajmalicine. Use has been made^{74,75} of this variation in chemical shift of the angular proton with orientation with respect to the nitrogen lone pair in the conformational analysis of benzoquinolizidines (**51**). Thus the configurations and preferred conformations **54** and **55** (R = *p*-chlorophenyl) were assigned to the



54

55


isomeric *cis*- and *trans*-2-(*p*-chlorophenyl)benzoquinolizidines since **54** exhibited Bohlmann bands in the infrared and the angular 11b proton absorbed at τ 6.2, whereas **55** showed no Bohlmann bands but its 11b proton appeared as a one-proton triplet (J_{aa} and J_{ee}) at τ 6. The alternative *cis*-fused conformation **56** was adopted by 2-keto-4,4'-dimethyl-9,10-dimethoxy-



56

benzoquinolizidine. This compound showed no Bohlmann bands and its angular proton gave rise to a quartet (J_{aa} and

J_{ee}) centered at τ 5.85. The stereochemistry of **51** (R' = MeO; R = Ph, C₆H₇, *p*-tolyl, *p*-chlorophenyl) and **57** (R = CH₃, Ph)




57


has been studied⁷⁶ using infrared data and rates of dehydrogenation. It would appear that the *cis*-2*H*,11*bH* compounds adopt the *trans*-fused ring conformation whereas some of the *trans*-2*H*,11*bH* compounds may exist as equilibrium mixtures of *cis*- and *trans*-fused ring conformations. The *trans*-2*H*,11*bH*-2-aryl compounds, however, undoubtedly adopt **55** as the preferred conformation. Bohlmann⁶⁵ has studied **51** (R' = OMe, R = COOCH₃, CH₂OH, CH₂OAc) and found that the compounds exhibiting Bohlmann bands show a τ value for the angular 11*bH* proton of τ 6.55 whereas in the other isomers showing no Bohlmann bands this proton absorbs at τ 6.19–6.32.

Both epimers of the hydroxybenzoquinolizidine **51** (R' = H, R = OH) have been shown⁶⁸ to exist in *trans*-fused ring conformations and the axial-equatorial orientation of the hydroxy group assigned from a study of the carbinol proton signals in the nmr. Derivatives of benzo[*c*]quinolizidines have been described by Kupchan.^{70,71}

Three of the four possible isomers of 8-azaestrone have been obtained and assigned configurations and preferred conformations (**58**, **59**, and **60a**).⁷⁷ **58** and **59** showed Bohlmann bands




58



59

whereas **60a** did not. The chemical shift of the angular protons were in agreement with this assignment. Some 12-keto derivatives of this system have been described⁷⁸ together with 220-MHz spectral data.



60a

60b

(74) M. Uskokovic, H. Bruderer, C. von Planta, T. Williams, and A. Brossi, *J. Amer. Chem. Soc.*, **86**, 3364 (1964).


(75) H. Bruderer, M. Baumann, M. Uskokovic, and A. Brossi, *Helv. Chim. Acta*, **47**, 1852 (1964).

(76) J. Gootjer, A. M. de Roos, and W. Th. Nauta, *Recl. Trav. Chim. Pays-Bas*, **85**, 491 (1966).

(77) R. E. Brown, D. M. Lustgarten, R. J. Stanaback, and R. I. Meltzer, *J. Org. Chem.*, **31**, 1489 (1966).

(78) N. S. Bhacca, A. I. Meyers, and A. H. Reine, *Tetrahedron Lett.*, 229 (1968).


The cyclopentenoquinolizidine **61** has been assigned⁷⁹ a trans-fused ring conformation on the basis of infrared and nmr evidence.



The four isomers of the inside yohimbanes (**62**) have had their stereochemistry assigned⁸⁰ on the basis of chemical and spectroscopic data.


h. Quinolizidine Methiodides

Quaternization of quinolizidine^{81,82} gives the trans-fused methiodide **63** whereas treatment of 2-(4'-ethoxybutyl)-1-methylpiperidine with hydriodic acid affords the cis-fused methiodide **64**. This latter isomer was also obtained^{81,82} by



catalytic reduction of **65**. From an examination of the nmr spectra of these isomers and of the methiodides of the mono-methylquinolizidines, it became apparent that the *N*-methyl group absorbs at lower field (τ 6.84–6.92) in the cis-fused salts than in the trans-fused analogs (τ 7.01–7.06).

Methylation of **66** gave two salts, assigned⁸⁸ the trans- and



cis-fused ring stereochemistry since their *N*-methyl resonances appeared at τ 6.90 and 6.65, respectively. The epimer of **66** gave only one methiodide ($\tau(N\text{-methyl})$ 6.72), and this was tentatively assigned⁸⁸ the cis-ring fusion. Methylation studies on the epimeric 1-hydroxy-, 1-hydroxy-1-methyl-, and 1-hydroxy-1-phenylquinolizidines have been carried out⁸⁶ and stereochemical assignments made. These are as shown in Table IV.

The hydroxyquinolizidines shown in Table III have been quaternized and the structures shown in Table IV proposed⁴¹ for the methiodides on the basis of the nmr data.

Table IV

Nmr Spectra of Quinolizidine Methiodides^{86,41}

Compound		Chemical shift <i>N</i> -Me (τ)
	R = H	6.88
	R = Me	6.93
	R = Ph	6.42
	R = H	6.77
	R = Me	6.78
	R = Ph	5.99
	R = H	6.54
	R = Me	6.55
	R = Ph	5.99
		6.38
		6.38
		6.32
		6.54
		6.15
		6.51
		6.38

B. FUSED 6/6 RING SYSTEMS WITH ADDITIONAL HETEROATOMS

The introduction of a second heteroatom into the quinolizidine ring system might be expected to influence conformational preferences since in certain situations dipole-dipole interactions will be significant and there will also be differences in the types of nonbonded interactions and in bond lengths and angles.

(79) A. I. Meyers and N. K. Ralhan, *J. Org. Chem.*, **28**, 2950 (1963).

(80) G. C. Morrison, W. A. Cetenko, and J. Shavel, *ibid.*, **32**, 2768 (1967).

(81) K. Schofield and R. J. Wells, *Chem. Ind. (London)*, 572 (1963).


(82) K. Schofield and R. J. Wells, *Aust. J. Chem.*, **18**, 1423 (1965).

(83) J. D. England, D. Temple, and J. Sam, *J. Med. Chem.*, **11**, 353 (1968).

1. One Additional Oxygen Atom

Of particular importance in the conformational analysis of 1,3-hetero systems has been the value of the geminal coupling constant (J_{gem})^{84,85} of the methylene protons situated between the heteroatoms. A molecular orbital theory⁸⁶ of geminal coupling constants shows that in such a system inductive removal of electrons from the symmetric bonding orbital of the CH_2 group and transfer of lone-pair electrons into the anti-symmetric bonding orbital both result in a positive contribution to the value of J_{gem} . This latter process is dependent upon the dihedral angle between the heteroatom lone-pair orbitals and the adjacent C-H bonds: a parallel arrangement^{87,88} of the lone-pair orbitals and a CH bond producing the most positive contribution to J_{gem} and an arrangement such that the lone pair bisects the CH_2 giving rise to no contribution from the lone pair.


Since the geminal coupling constants of the C-2 methylene protons of *cis*-10,6-*H*-10-methyl- and *trans*-10,6-*H*-10-methyl-3-oxa-1-azabicyclo[4.4.0]decane were⁸⁹ -8 and -10 Hz, respectively, they were assigned the preferred conformations shown (67 and 68a), the larger value of J_{gem} corresponding to




the parallel arrangement of lone pair and CH bonds, and the more negative J_{gem} consistent with 68a in which the nitrogen lone pair bisects the CH_2 . The *trans*-fused ring compound 67 also showed Bohlmann bands which were absent in the *cis*-fused ring compound 68. The *trans*-10,6-*H*-10-methyl-3-oxa-1-azabicyclo[4.4.0]decane adopts the *cis*-fused conformation 68a rather than the alternative *trans* conformation 68b not only to relieve nonbonded interactions involving the axial methyl group but also to alleviate the unfavorable interaction arising from the parallel arrangement of lone pairs of electrons on nitrogen and oxygen. The destabilizing effect of parallel lone pairs in 1,3-hetero systems has also been discussed by Eliel.⁹⁰ Both of the possible 4-methyl-3-oxa-1-azabicyclo[4.4.0]decanes and the two 4-phenyl compounds have been shown⁸⁹ to adopt predominantly *trans*-fused ring conformations since they all show Bohlmann bands and a geminal coupling constant for the C-2 methylene protons of *ca.* -8 Hz. It has generally been assumed that Bohlmann's infrared criterion originally deduced for quinolizidines is applicable to other

saturated systems with bridgehead nitrogen atoms, and all the available evidence supports this assumption.

Several other substituted 3-oxa-1-azabicyclo[4.4.0]decanes (69) have been prepared^{89,91,92} and stereochemical studies carried out with them, in some cases, in order to determine the




configurations of the alkaloids sedamin and sedridin. Condensation of the two racemic diastereoisomeric piperidylpropanols with aromatic aldehydes gave⁹¹ one isomer in each case, and these were assigned the *trans*-fused ring conformations 70 and 71 with the bulky aryl group equatorial in both isomers. The



axial-equatorial nature of the methyl group was deduced from the chemical shifts of the C-4 proton and from the values of the coupling constants with adjacent protons.

However, more recently two isomers have been reported⁹² from the reaction of (-)-sedridine, a piperidylpropanol, with *p*-nitrobenzaldehyde. One of the isomers was assigned an axial 4-methyl group on the basis of nmr spectroscopy, and the *cis*-fused ring conformation 72 was considered in best agreement



with the spectral evidence. The other isomer was assigned the *trans*-fused ring conformation 71 ($R = p$ -nitro) with an equatorial methyl group. The *cis*-fused ring conformation 72 seems very unlikely for one of these isomers, particularly as models show that the axial 4-methyl group in this conformation points backwards under the other six-membered ring and introduces an extremely severe nonbonded interaction. The Bohlmann infrared criterion would also be expected to differentiate between the two conformations 71 and 72, but it was stated that the infrared spectra of the two compounds were similar. A *trans*-fused ring conformation such as 71 with an axial *p*-nitrophenyl group would therefore appear to be a more probable structure for *trans*-2, *cis*-4,6-*H*-2-*p*-nitrophenyl-4-methyl-3-oxa-1-azabicyclo[4.4.0]decane.

(84) R. C. Cookson, T. A. Crabb, J. J. Frankel, and J. Hudec, *Tetrahedron, Suppl.*, **7**, 355 (1966).

(85) R. Cahill, R. C. Cookson, and T. A. Crabb, *Tetrahedron*, **25**, 4681 (1969).

(86) J. A. Pople and A. A. Bothner-By, *J. Chem. Phys.*, **42**, 1339 (1965).

(87) M. Anteunis, *Bull. Soc. Chim. Belg.*, **75**, 413 (1966).

(88) R. C. Cookson and T. A. Crabb, *Tetrahedron*, **24**, 2385 (1968).


(89) T. A. Crabb and R. F. Newton, *ibid.*, **24**, 4423 (1968).

(90) E. L. Eliel, *Kem. Tidskr.*, **81**, 22 (1969).


(91) H. C. Beyerman, L. Maat, A. Van Veen, and A. Zweistra, *Recl. Trav. Chim. Pays-Bas*, **74**, 1568 (1955).

(92) G. A. Cooke and G. Fodor, *Can. J. Chem.*, **46**, 1105 (1968).

trans-4,6-*H*-4-Phenyl-3-oxa-1-azabicyclo[4.4.0]decane has been shown⁹³ to exist predominantly in a *trans*-fused ring conformation with an axial 4-phenyl group (73).




Riddell⁹³ reported the conformation of 3-oxa-1-azabicyclo[3.3.1]nonane as being 74. This compound had a J_{gem} for the




C-2 methylene group of -10.5 Hz consistent with a conformation such that the nitrogen lone pair bisects the methylene group.

The four possible isomers of perhydrobenzo[*e*]pyrido[1,2-*c*]-1,3-oxazine (75, $n = 2$) have been obtained⁹⁴ and as-




signed the stereostructures 76-79. The *trans* A:B junction in 76, 77, and 78 was shown by Bohlmann bands and J_{gem} (C-6



methylene) of *ca.* -7.5 Hz and the *cis* A:B junction in 79 by a similar geminal coupling constant and the absence of Bohlmann bands. The nature of the B:C ring fusions was deduced from the chemical shift and couplings of the C-7a proton and the chemical shifts of the C-2 protons. Independent evidence for the assignments was obtained by studying the rates of methylation. The corresponding cyclopentano (75, $n = 1$) and cycloheptano (75, $n = 3$) compounds and some 5-methyl-substituted derivatives were also studied.


5-Oxa-1-azabicyclo[4.4.0]decane (80) showed two bands between 2800 and 2700 cm^{-1} in its infrared spectrum and was accordingly assigned⁹⁵ a *trans*-fused ring conformation. 6-

Methyl-5-oxa-1-azabicyclo[4.4.0]decane (81), assigned a *trans*-fused ring conformation on the same basis, might be expected to exist as an equilibrium mixture containing significant amounts of the *cis*-fused ring conformation, but, as discussed above, infrared data in the CH stretching region are not able to demonstrate the presence of such conformations. The tricyclic 3-oxa-7-azatricyclo[7.3.1.0]tridecane was found by infrared to exist predominantly in the *trans*-fused ring conformation 82.




2. One Additional Nitrogen Atom

The configurations and preferred conformations of a series of substituted 1,3-diazabicyclo[4.4.0]decanes (83) have been



deduced⁹⁶ from chemical and spectroscopic evidence. The 3-alkyl- and 3-aryl-1,3-diazabicyclo[4.4.0]decanes (83, R = CH_3 , C_2H_5 , $(\text{CH}_3)_2\text{C}$, C_6H_5) were all found to exist in predominantly *trans*-fused ring conformations with equatorial 3-nitrogen substituents ($J_{gem} = -8.4$ to -8.6 Hz and strong Bohlmann bands). The geminal coupling constant of -8.8 Hz for the C-2 methylene group showed that the *cis*-4,6-*H*-3,4-dimethyl and *cis*-4,6-*H*-3-methyl-4-ethyl epimers existed in the *trans*-fused ring conformation with an equatorial 3-methyl group (84), while the *trans*-4,6-*H*-3,4-dimethyl ($J_{gem} = -10.8$ Hz) and *trans*-4,6-*H*-3-methyl-4-ethyl compounds ($J_{gem} =$



-11.4 Hz) existed predominantly in the *trans*-fused ring conformation with an axial 3-methyl group (85). This is a clear example of the importance of syn-axial lone-pair interactions in influencing the position of conformational equilibrium in such systems. Both of the possible 3,5-dimethyl-1,3-diazabicyclo[4.4.0]decanes were assigned *trans*-fused ring conformations with equatorial 3-methyl groups on the basis of their nmr spectra ($J_{gem} = -8.4$ and -8.7 Hz). The configurations of the 3,10-dimethyl- and 3-*tert*-butyl-10-methyl-1,3-diazabicyclo[4.4.0]decanes were deduced from chemical evidence. The predominant conformation of 3-methyl- and 3-*tert*-butyl-*cis*-10,6-*H*-10-methyl-1,3-diazabicyclo[4.4.0]decane was found to be that with a *trans*-fused ring junction and an equatorial 3-


(93) F. G. Riddell and J. M. Lehn, *J. Chem. Soc. B*, 1224 (1968).

(94) T. A. Crabb and E. R. Jones, *Chem. Ind. (London)*, 1695 (1968); *Tetrahedron*, 26, 1217 (1970).


(95) N. J. Leonard and W. K. Musker, *J. Amer. Chem. Soc.*, 82, 5148 (1960).

(96) T. A. Crabb and R. F. Newton, *Tetrahedron*, 26, 701 (1970).


alkyl group. The corresponding *trans*-10,6-*H*-10-methyl compounds existed as conformational mixtures containing appreciable amounts of the *cis*- and *trans*-fused ring conformations (**86** and **87**) as shown by values of $J_{gem} = -9.8$ and -9.9 Hz.




Both the *syn* (**88**) and *anti* (**89**) perhydrodipyrido[1,2-*c*:




2',1'-*f*]pyrimidines have been obtained⁹⁷ and the C-6 methylene protons in the *syn* compound show a J_{gem} of -8.5 Hz and a difference in chemical shift of 1.0 ppm, consistent with the structure shown with both lone pairs *trans* to C-6 H_{ax} . The unsubstituted *anti* compound (**89**) is in rapid conformational equilibrium as indicated. The *syn* compound **88** has Bohlmann bands of much greater area than the *anti* compound **89** since it possesses two *trans*-fused quinolizidine rings.



1,4-Diazabicyclo[4.4.0]decane (**90**) has been synthesized⁹⁸




and assigned a *trans*-fused ring conformation on the basis of the infrared spectrum of its 4-phenylurea derivative which had bands at 2760 and 2785 cm^{-1} . The infrared spectra of 1,3-diaza-, 1,4-diaza-, and 1,5-diazabicyclo[4.4.0]decane all show Bohlmann bands, and these compounds were accordingly assigned⁹⁹ *trans*-fused ring conformations. The preferred conformations of **91** and **92** have been suggested from a




(97) P. J. Chivers and T. A. Crabb, *Tetrahedron*, **26**, 3369 (1970).
 (98) M. E. Freed and A. R. Day, *J. Org. Chem.*, **25**, 2108 (1960).
 (99) R. L. Peck and A. R. Day, *J. Heterocycl. Chem.*, **6**, 181 (1969).

study¹⁰⁰ of their infrared spectra. Two isomers of **93** have been obtained and their stereochemistry determined¹⁰¹ from a study of the C- CH_3 chemical shifts and apparent coupling constants. Both isomers showed Bohlmann bands.



3. One Additional Sulfur Atom


3-Thia-1-azabicyclo[4.4.0]decane has been assigned¹⁰² the *trans*-fused ring conformation on the basis of its infrared (strong Bohlmann bands) and nmr spectra ($J_{2ax,2eq} = -12.0$ Hz) (Table V). *cis*-7,6-*H*-7-Methyl-3-thia-1-azabicyclo[4.4.0]decane showed very weak Bohlmann bands in the infrared indicating the *cis*-fused conformation **95**. However, $J_{2ax,2eq}$ was



only slightly smaller (-12.8 Hz) than observed for **94**, and this has been explained in terms of deviation from chair geometry of the 1,3-hetero ring as a consequence of the long C-S bond.

4. Two Nitrogen Atoms at a Ring Fusion

The nmr spectra and conformations of the compounds **96**¹⁰³ and **97**¹⁰⁴ have been discussed in detail.



C. FUSED 6/5 RING SYSTEMS WITH NO ADDITIONAL HETEROATOM

1. Indolizidine

ΔG° for the *cis* \rightleftharpoons *trans* indolizidine equilibrium has been determined¹³ as -2.4 kcal/mol, a value very close to that for the quinolizidine equilibrium (-2.6 kcal/mol).¹³ The free energy difference between the carbocyclic analogs *cis*- and *trans*-hydrindane and *cis*- and *trans*-decalin are 2.7 and 0.3 kcal/mol, respectively, and the small free energy difference between *cis*- and *trans*-hydrindane has been explained in terms of strain involving the ring fusion which raises the energy of

(100) A. R. Day and H. B. Sullivan, *Diss. Abstr.*, **24**, 1838 (1963).













(101) K. G. R. Pachler, *J. Chem. Soc. B*, **7**, 760 (1968).

(102) T. A. Crabb and R. F. Newton, *Tetrahedron*, **26**, 3941 (1970).

(103) J. P. Kintzinger, J. M. Lehn, and J. Wagner, *Chem. Commun.*, **5**, 206 (1967).

(104) B. Junge and H. A. Staab, *Tetrahedron Lett.*, **8**, 709 (1967);

Table V
Nmr and Ir Spectra of 3-Hetero-1-azabicyclo[4.4.0]decenes

Compound	Nmr (C-2 protons) J_{gem} (Hz)	Δ^c (ppm)	Ir Bohlmann region cm^{-1} (ϵa)	Compound	Nmr (C-2 protons) J_{gem} (Hz)	Δ^c (ppm)	Ir Bohlmann region cm^{-1} (ϵa)
	-8.0 ^a	0.66	2813 (50) 2776 (68) 2762 (84) 2739 (44) 2725 (50) 2682 (52)		-8.6 ^b	2.00	2821 (51) 2790 (78) 2720 (43) 2680 (43) 2660 (40)
	-8.0 ^a	0.61	2811 (58) 2764 (94) 2735 (54) 2723 (39) 2687 (36)		-9.8 ^b	0.91	2821 (49) 2800 (55) 2760 (28) 2600 (26)
	-8.0 (benzene)	0 ^a	2811 (60) 2759 (94) 2737 (64) 2714 (46) 2682 (54) 2656 (30)		-8.8 ^b	0.96	2825 (71) 2795 (110) 2750 (60) 2732 (50) 2670 (45)
	-8.0 ^a	1.31	2812 (46) 2774 (74) 2758 (62) 2725 (72) 2698 (50) 2683 (34)		-10.8 (benzene)	0 ^b	2801 (50) 2775 (55) 2739 (35) 2670 (25)
	-10.0 ^a	0.60	2760 (26) 2725 (23)		-12.0 ^a	0.31	2782 (75) 2738 (32) 2720 (32)
	-8.5 ^b	1.27	2822 (51) 2785 (70) 2740 (41) 2720 (41) 2677 (46)		-12.8	0.65	2790 (14) 2720 (12)


^a 10% CCl₄ solution. ^b 10% CDCl₃ solution. ^c Difference in chemical shift between C-2 protons.

trans-hydrindane. In order to explain the difference in ΔG° values between the indolizidine and hydrindane systems, nmr evidence has been quoted¹⁰⁵ to suggest that the strain present in *trans*-hydrindane is relieved in *trans*-indolizidine by flattening of the bonds around the bridgehead nitrogen atom.

The Bohlmann regions of the infrared spectra of indolizidine (2), 3,3'-dideuterioindolizidine, and 9-deuterioindolizidine have been compared.¹⁰⁶ It was found that replacement of the C-9 axial hydrogen in indolizidine by deuterium reduced the intensity of the Bohlmann bands by 28%, whereas replacement of the pseudo axial C-3 hydrogen reduced the bands by 33%. This result was unexpected since the C-3 hydrogen atom is not strictly *trans* coplanar with the nitrogen atom lone pair of electrons.

The configurations and preferred conformations of the four 5,7-dimethylindolizidines have been assigned¹⁰⁷ on the basis of

their nmr (C-methyl chemical shifts) and infrared spectra. **98** and **99** show prominent bands in the Bohlmann region but



these are much reduced in **100**. The *trans*-fused conformation **100** has only two correctly orientated CH bonds necessary for strong Bohlmann bands compared with three in **98** and **99**, and so a drop in intensity would be expected for **100**. Isomer **101**

(105) T. A. Crabb and R. F. Newton, *Tetrahedron Lett.*, 1551 (1970).

(106) A. E. Theobald and R. G. Lingard, *Spectrochim. Acta*, **24**, 1245 (1968).

(107) B. Lüning and C. Lundin, *Acta Chem. Scand.*, **21**, 2136 (1967).

showed very little absorption in the 2800–2700-cm⁻¹ region indicative of the cis-fused conformation shown.

The 1-, 2-, 7-, and 8-hydroxyindolizidines have been studied by Aaron.^{108,109} All eight compounds (Table VI) show prominent Bohlmann bands indicating the trans-fused ring conformations **102**–**109**. **102**, **104**, and **108** show infrared absorptions characteristic of intramolecular hydrogen bonding and

Table VI

Nmr and Ir Spectra of Hydroxyindolizidines^{108,109}

Compound	No.	—Ir spectra (cm ⁻¹)—		Nmr spectra	
		OH	Bohlmann region	τ	W_H^a (Hz)
	102	3623, 3580	2790, 2750, 2725	6.10	11.5
	103	3632	2795, 2730	6.25	21
	104	3624, 3594	2785, 2720	5.90	17
	105	3626	2790, 2720	5.75	14
	106	3626	2800, 2745	5.98	7.4
	107	3624	2788, 2751, 2714	6.53	21
	108	3522	2789, 2736	6.32	6.5
	109	3627	2790, 2725	6.75	>20


^a Peak width at half-height.

using long-path-length cells **109** has been shown¹³ to exist in equilibrium with 3% of the cis-fused ring conformation with an axial intramolecularly hydrogen-bonded hydroxy group. The configurations of the remaining alcohols were assigned from the chemical shift and line width of the carbinol proton signals. In each case (with exception of **104** and **105**) the axial proton absorbed at higher field than the equatorial proton. However, in **104** and **105** the carbinol proton did not appear to be definitely pseudoaxial or pseudoequatorial.

The 1-, 2-, 7-, and 8-ketoindolizidines also exist^{109–111} in predominantly trans-fused ring conformations as shown by the presence of Bohlmann bands.


The carbinyl acetate proton of *N*-acetylsafraamine absorbs at τ 4.76 ($W_H = 13$ Hz). Comparison with the chemical shifts of the corresponding protons in the acetates of **102** (τ 4.79,

$W_H = 13$ Hz) and of **103** (τ , 5.24, $W_H = 21$ Hz) permitted¹¹² the assignment of configuration shown in **110**. The coupling




constants in *N*-acetylsafraamine hydrochloride (D₂O solution) were $J_{1,9} = 6.5$ Hz, $J_{5ax,6} = 2.8$ Hz, and $J_{5ax,5eq} = -13$ Hz.

A dimethylcyclopentenoindolizidine was assigned the stereochemistry of **111** on the basis of the presence of Bohlmann bands and the chemical shift and coupling constant (10 Hz) of the angular proton. **112**¹¹³ shows Bohlmann bands.



2. Indolizidine Methiodides

Methylation of indolizidine produced¹¹⁴ a mixture of the cis- and trans-fused methiodides **113** and **114** showing *N*-methyl



resonances at τ 6.88 and 7.18, respectively. The stereochemistry of the ring fusion was assigned from these τ values since very similar values had been observed for the corresponding quinoindolizidine methiodides.

The different τ values have been explained by Elvidge¹¹⁵ in terms of the shielding effects of neighboring C–C and C–N bonds. The isomeric 5,7-dimethylindolizidines have been methylated¹⁰⁷ and whereas **98** and **99** each give two isomeric methiodides with *N*-methyl chemical shifts in the normal range, **100** and **101** give only one each. The methiodide of **101** must be cis-fused but shows an unusually high-field *N*-methyl resonance at τ 7.04; this may be due to the shielding influence of the adjacent equatorial methyl group.

D. FUSED 6/5 RING SYSTEMS WITH ADDITIONAL HETEROATOMS

The value of the J_{gem} for the N–CH₂–X protons and the difference in chemical shift between these protons have been shown to be reliable criteria in determining the preferred conformations of **115** (X = O),¹¹⁶ **115** (X = S),¹¹⁷ **115** (X =

(108) C. P. Rader, R. L. Young, and H. S. Aaron, *J. Org. Chem.*, **30**, 1536 (1965).

(109) H. S. Aaron, C. P. Rader, and G. E. Wicks, *ibid.*, **31**, 3502 (1966).

(110) S. Yamada and T. Kuneida, *Chem. Pharm. Bull. Jap.*, **15**, 490 (1967).

(111) R. Cahill and T. A. Crabb, unpublished work.

(112) R. A. Gardiner and K. L. Rinehart, *J. Amer. Chem. Soc.*, **90**, 5639 (1968).

(113) J. P. Li, *J. Heterocycl. Chem.*, **5**, 703 (1968).

(114) W. L. Meyer and N. Sapianchiay, *J. Amer. Chem. Soc.*, **86**, 3343 (1964).













(115) J. A. Elvidge and I. M. Skvortsov, *J. Chem. Soc. B*, 1589 (1968).

(116) T. A. Crabb and R. F. Newton, *Chem. Ind. (London)*, 339 (1966); *J. Heterocycl. Chem.*, **3**, 418 (1966); *Tetrahedron*, **24**, 1997 (1968).


(117) T. A. Crabb and R. F. Newton, *ibid.*, **24**, 2485 (1968).

Table VII


Nmr and Ir Spectra of 8-Hetero-1-azabicyclo[4.3.0]nonanes

Compound	Nmr (C-9 protons)		Ir Bohlmann region cm^{-1} (ϵ)
	J_{gem} (Hz)	$\Delta H_{99'}$ (ppm)	
	-0.8	0.79	2808 (150) 2778 (146) 2762 (107) 2754 (96) 2727 (71) 2710 (46)
	-6.0	0.65	2785 (115) 2734 (30) 2715 (33) 2635 (21)
	-4.3	0.87	2796 (64) 2728 (43) 2600 (13)
	-3.6	0.77	2803 (81) 2777 (67) 2742 (77) 2708 (37)
	-5.0	0.19	2830 (57) 2777 (30)
	-8.5	0.22	2803 (60) 2722 (16)
	-5.7	0.18	2825 (60)
	-6.0	0.34	2803 (61) 2740 (45) 2710 (24)
	-2.4	0.58	2803 (97) 2776 (97) 2752 (62)
	-7.2	0.22	2785 (122) 2731 (19) 2718 (21) 2710 (21) 2634 (13)
	-4.1	0.72	2797 (82) 2770 (60) 2735 (51) 2670 (23)
	-3.8	0.72	2802 (84) 2774 (67) 2741 (74)

N-alkyl),¹¹⁸ and **115** (X = *N*-phenyl).¹¹⁹ Spectral data on these systems are summarized in Table VII. Trans-fused ring com-




(118) T. A. Crabb and R. F. Newton, *Tetrahedron*, **24**, 6327 (1968).(119) T. A. Crabb and R. F. Newton, *J. Heterocycl. Chem.*, **6**, 301 (1969).

pounds (**116**) show a very positive value of J_{gem} and a large difference in chemical shift for the C-9 protons whereas cis-fused ring compounds (**117**) show a smaller chemical shift




difference and a more negative J_{gem} . The very large value of $J_{99'} = -0.8$ Hz for **116** (X = O) has been explained¹¹⁸ in terms of flattening of the bonds involving the bridgehead nitrogen atom, thus increasing the overlap of the nitrogen lone-pair orbitals with the adjacent CH bonds.¹⁰⁵ The infrared spectra of trans-fused ring compounds exhibited strong absorptions in the Bohlmann region whereas the cis-fused ring compounds did not (Figure 2). The unsubstituted parent compounds (**115**, X = O and X = S) show intermediate values of J_{gem} and $\Delta(H9H9')$ indicating an equilibrium between **116** and **117** at room temperature. Since indolizidine exists at room temperature almost entirely in the trans conformation, the existence of **115** (R = H; X = O and S) as equilibrium mixtures containing a large proportion of cis-fused conformations shows the destabilizing influence of the 1,3-heteroatom lone pairs in the trans-fused conformation **116**. **115** (X = N-R) exists at room temperature in a predominantly trans-fused ring conformation (**116**, X = N-R) contrary to an earlier report.⁹⁸

The 4-phenylurea derivative of 1,4-diazabicyclo[4.3.0]nonane (**118**)⁹⁸ exhibited strong Bohlmann bands and was accordingly assigned a predominantly trans-fused ring conformation. Leonard⁹⁵ synthesized 7-oxa-1-azabicyclo[4.3.0]nonane (**119**) and 5-oxa-1-azabicyclo[4.3.0]nonane (**120**) and



assigned both predominantly trans-fused ring conformations on the basis of their infrared spectra.

The configurations and preferred conformations of a series of monomethyl-substituted perhydrodipyrido[1,2-*c*:2',1'-*e*]imidazoles (**121**) have been assigned¹²⁰ from a study of their



infrared and nmr spectra. With one exception, all the isomers obtained were assigned the trans,anti,trans (**122**) or the trans,

(120) P. J. Chivers, T. A. Crabb, and R. O. Williams, *Tetrahedron*, **24**, 6625 (1968).






Figure 2. Infrared spectra (CH stretching region) of some 8-oxa-1-azabicyclo[4.3.0]nonanes.

syn,trans (**123**) stereochemistry. These showed marked ab-



sorption in the Bohlmann region of the infrared, and compounds **122** showed either zero or very small chemical shift differences between the N-CH₂-N protons whereas **123** showed chemical shift differences of ca. 0.4 ppm. $J_{\text{H6H6}'}$ for **122** and **123** varied between -3.4 and -4.5 Hz. *trans*-1*H*,-11*bH*-1-Methyl-*syn*-perhydrodipyrido[1,2-*c*:2',1'-*e*]imidazole was assigned the conformation **124** since it showed a marked reduction in the area of Bohlmann bands and a $J_{\text{H6H6}'}$ of -8.0 Hz. A similar study¹²¹ of the perhydropyrido[1,2-*c*]pyrrolo-[2,1-*e*]imidazoles (**125**) permitted assignments of the structures **126** ($J_{\text{H6H6}'} = \text{ca. } -7$ Hz, marked Bohlmann bands), **127** ($J_{\text{H6H6}'} \text{ ca. } -4$ Hz, marked Bohlmann bands), and **128** ($J =$




ca. -6 Hz, weak Bohlmann bands) to the various isomers.

(121) P. J. Chivers, T. A. Crabb, and R. O. Williams, *Tetrahedron*, **25**, 2921 (1969).

E. FUSED 5/5 RING SYSTEMS

A detailed discussion of the stereochemistry and nmr spectrum of pyrrolizidine (**3**) has been given by Skvortsov and Elvidge.¹¹⁵ This compound exists mainly in the *cis*-fused ring conformation with pseudorotation occurring in each ring. The angular proton absorbs (CCl₄ solution) at τ 6.55 and the *exo* and *endo* protons adjacent to nitrogen absorb at τ 6.97 and 7.40, respectively; the geminal coupling constant for the N-CH₂ methylene protons is -9.75 Hz. The nmr spectrum of 3-*endo*-methylpyrrolizidine is temperature dependent, and at room temperature the compound must exist as a mixture of *cis*- and *trans*-fused ring conformations.

The 3-*exo*-methyl isomer exists in a *cis*-fused ring conformation at temperatures between -85 and $+100^\circ$. The methyl resonances in the *exo* and *endo* compounds were at τ 8.84 and 8.88 (benzene solution), respectively. The *N*-methyl resonance in pyrrolizidine methiodide occurs at τ 6.86 (D₂O) (see also ref 114). The angular proton in 1-methoxycarbonylpyrrolizidine (**129**)¹²² absorbs at τ 6.2.





The 1- and 2-hydroxypyrrolizidines¹⁰⁹ exist in the *cis*-fused conformation. All four isomers show a free OH stretching vibration at 3625 – 3630 cm⁻¹ and the *trans*-2,8-*H* compound

(122) B. Lüning and H. Trankner, *Acta Chem. Scand.*, **22**, 2324 (1968).

shows in addition a band at 3590 cm^{-1} due to a weak intramolecular hydrogen bond.

The stereochemistry of the pyrrolizidine alkaloids has been reviewed by Fodor,¹²³ and the nmr spectra of these alkaloids have been studied by Culvenor.⁵ The nmr spectra of **130**,¹²⁴ derivatives of **131**,¹²⁵ and of **132**¹²⁶ have been reported. The

**130****131**



(123) G. Fodor, *Recent Develop. Chem. Nat. Carbon Compounds*, **1**, 15 (1965).

(124) R. C. Fort and W. L. Semon, *J. Org. Chem.*, **32**, 3685 (1967).

(125) R. G. Hiskey and S. J. Dominianni, *ibid.*, **30**, 1506 (1965).

(126) J. L. Aubagnac, J. Elguero, and R. Jacquier, *Bull. Soc. Chim. Fr.*, **9**, 3869 (1968).

temperature dependence of the nmr spectrum of **133** has been

**132****133**

studied.¹⁰³ At -60° the ring methylenes appear as an AB quartet with $J = -9.6\text{ Hz}$, whereas at room temperature a singlet is observed. These changes may be attributed to the interconversion between the two cis-fused ring conformations by nitrogen inversion.