

HETEROCYCLIC SYSTEMS BEARING PHOSPHORUS SUBSTITUENTS. SYNTHESIS AND CHEMISTRY

DEREK REDMORE

Petrolite Corporation, Tretolite Division, St. Louis, Missouri 63119

Received November 30, 1970 (Revised Manuscript Received January 15, 1971)

Contents

I. Introduction	315
II. General Methods of Synthesis	315
A. Nucleophilic Displacement Reactions on Phosphorus by Heterocyclic Organometallic Reagents	315
1. Preparation of Phosphines, Phosphine Oxides, and Phosphine Sulfides	315
2. Preparation of Phosphinates, Phosphonates, and Phosphoranes	316
B. Nucleophilic Displacement Reactions by Phosphorus Nucleophiles	317
1. Arbuzov and Michaelis-Becker Reactions on Heterocyclic Halides	317
2. Other Nucleophilic Displacements by Phosphorus	320
C. Nucleophilic Addition Reactions	321
D. Cycloaddition Reactions	324
E. Condensation Reactions	326
1. Epoxyethylphosphorus Compounds	326
2. Five-Membered Ring Compounds	328
3. Six-Membered Ring Compounds	330
F. Radical Reactions	331
G. Electrophilic Substitution by Phosphorus	332
III. Chemistry	332
A. Carbon-Phosphorus Bond Cleavage	332
1. Wittig and Wadsworth-Emmons Reactions	332
2. Other C-P Bond Cleavages	333
B. Rearrangements	334
C. Other Reactions	335
IV. Properties	335
A. Spectra	335
1. Ultraviolet Adsorption Spectra	336
2. Nuclear Magnetic Resonance Spectra	336
B. Biological Activity	337

I. Introduction

This review is limited to those compounds in which the phosphorus substituent is attached to a carbon atom of the heterocyclic ring, so that compounds such as tris-1-arizidinyl phosphoramidate (commonly incorrectly called tris-aziridinyl phosphine) are excluded from consideration. Numerous heterocyclic systems are represented and several phosphorus groupings, including phosphines, phosphine oxides, sulfides, and selenides, phosphoranes, and phosphinic and phosphonic acid derivatives. The coverage is comprehensive through September 1970. Particular emphasis has been placed on synthetic methods,


chemistry, and properties which are peculiar to the heterocyclic phosphorus derivatives.

II. General Methods of Synthesis

A. NUCLEOPHILIC DISPLACEMENT REACTIONS ON PHOSPHORUS BY HETEROCYCLIC ORGANOMETALLIC REAGENTS

1. Preparation of Phosphines, Phosphine Oxides, and Phosphine Sulfides

Nucleophilic displacement on phosphorus esters and halides by organometallic reagents is a very facile process leading to tertiary phosphines or phosphine oxides.¹ The results obtained



using heterocyclic Grignard or lithium reagents do not differ from those obtained with aliphatic or aryl derivatives except that yields tend to be lower. This is due to the susceptibility of the heterocyclic organometallics to side reactions, such as self-condensation. The addition of phenyllithium to pyridine, for example, is a well-known reaction.

Phosphine oxides can be prepared by the reaction of a Grignard reagent on phosphorus trichloride or bromide followed by oxidation or, alternatively, from the nucleophilic displacement by the same reagent on phosphoryl chloride or a dialkyl phosphorochloridate. It appears that the single-stage procedure gives superior yields. Table I²⁻¹² summarizes the heterocyclic phosphines and phosphine oxides prepared by nucleophilic displacement on phosphorus. It should be noted that in the reaction of indolyl Grignard reagents on phos-

(1) K. D. Berlin, T. H. Austin, M. Peterson, and M. Nagabhushanam, *Top. Phosphorus Chem.*, **1**, 17 (1964).

(2) K. Issleib and A. Brock, *Z. Anorg. Allg. Chem.*, **292**, 245 (1957).

(3) C. E. Griffin, R. P. Peller, K. R. Martin, and J. A. Peters, *J. Org. Chem.*, **30**, 97 (1965).

(4) E. Niwa, H. Aoki, H. Tanaka, and K. Munakata, *Chem. Ber.*, **99**, 712 (1966).

(5) A. Burger and N. D. Dawson, *J. Org. Chem.*, **16**, 1250 (1951).

(6) H. J. Jakobsen and J. Nielsen, *Acta Chem. Scand.*, **23**, 1070 (1969).

(7) M. D. Rausch, T. R. Criswell, and A. K. Ignatowicz, *J. Organometal. Chem.*, **13**, 419 (1968).

(8) Q. Mingoia, *Gazz. Chim. Ital.*, **60**, 144 (1930).

(9) Q. Mingoia, *ibid.*, **62**, 333 (1932).

(10) W. C. Davies and F. G. Mann, *J. Chem. Soc.*, 279 (1944).


(11) E. Plazek and R. Tyka, *Zesz. Nauk. Politech. Wroclaw. Chem.*, **79**(1957); *Chem. Abstr.*, **52**, 20156 (1958).

(12) F. G. Mann and J. Watson, *J. Org. Chem.*, **13**, 502 (1948).

Table I
Phosphines and Phosphine Oxides by Nucleophilic Displacement on Phosphorus

Heterocyclic organometallic deriv	Phosphorus reagent	Product (yield, %)	Ref
Pyrrylpotassium	PCl ₃	Tri-2-pyrrylphosphine (40)	2
Pyrrylmagnesium bromide	POCl ₃	Tri-2-pyrrylphosphine oxide (42)	3
1-Methyl-2-pyrryllithium	(EtO) ₂ POCl	Tri-2(1-methylpyrryl)-phosphine oxide (28)	3
2-Furyllithium	PBr ₃	Tri-2-furylphosphine (33)	4
2-Furyllithium	POCl ₃	Tri-2-furylphosphine oxide (68)	3, 4
2-Thienylmagnesium bromide	PCl ₃	Tri-2-thienylphosphine (70)	2
2-Thienylmagnesium bromide	(EtO) ₂ POCl	Tri-2-thienylphosphine oxide (75)	3, 5
3-Thienyllithium	PBr ₃	Tri-3-thienylphosphine (50)	6
3,4,5-Trichloro-2-thienyllithium	Ph ₂ PCl	Diphenyl(3,4,5-trichloro-2-thienyl)phosphine (34)	7
Indolylmagnesium bromide	PCl ₃	Tri-3-indolylphosphine	8
2-Methylindolylmagnesium bromide	PCl ₃	Tri-3-(2-methylindolyl)-phosphine	8
3-Methylindolylmagnesium	PCl ₃	Tri-2-(3-methylindolyl)-phosphine	8
Indolylmagnesium bromide	POCl ₃	Tri-3-indolylphosphine oxide	9
2-Methylindolylmagnesium bromide	POCl ₃	Tri-3-(2-methylindolyl)-phosphine oxide	9
2-Pyridylmagnesium bromide	PCl ₃	Tri-2-pyridylphosphine	10
2-Pyridyllithium	PCl ₃	Tri-2-pyridylphosphine (38)	11
2-Pyridylmagnesium bromide	PhPCl ₂	Phenyldi(2-pyridyl)-phosphine	12
2-Pyridylmagnesium bromide	Ph ₂ PCl	Diphenyl(2-pyridyl)-phosphine (16)	12
3-Pyridylmagnesium bromide	<i>p</i> -BrC ₆ H ₄ (Ph)PCl	Phenyl- <i>p</i> -bromophenyl-3-pyridylphosphine	10

phoryl chloride the bis(indolyl)phosphinic acids **1** are reported




as by-products along with the product phosphine oxide **2**.⁹ The origin of the phosphinic acids **1** is not clear but most probably arises from degradation of the phosphine oxide **2** upon work-up in view of the reported facile C-P bond cleavage in **3**.¹³ The alternative genesis of **1** by displacement of two chloride ions from phosphoryl chloride by the indolyl Grignard followed by hydrolysis of the remaining chlorine seems unlikely.¹ Repetition of these syntheses would be worthwhile in view of the sketchy characterization in the original work.

2. Preparation of Phosphinates, Phosphonates, and Phosphoranes

In principle, reaction of phosphoryl chloride or dialkyl phosphorochloridates with one equivalent of heterocyclic Grignard or lithium reagent should yield a phosphonic acid dihalide or diester. In practice this is not readily achieved; the product phosphonic acid derivative competes with the starting halide for reaction with the organometallic reagent, so that a complex product mixture results.

Addition of pyrrylmagnesium bromide to diethyl phosphorochloridate in refluxing ether has allowed the preparation of diethyl pyrryl-2-phosphonate (**4**) in 30% yield.¹⁴ By a similar technique 1-methyl-2-pyrryllithium reacts with diethyl phosphorochloridate with the formation of diethyl 1-methylpyrryl-2-phosphonate (**5**).¹⁴ The reaction of 2,5-dimethyl-




pyrrylmagnesium bromide (**6**) with diethyl phosphorochloridate yields 3-ethyl-2,5-dimethylpyrrole (**10**) rather than the desired phosphonate **7**. This phosphonate **7** is postulated as


(13) J. C. Powers, *Tetrahedron Lett.*, 655 (1965).

(14) C. E. Griffin, R. P. Peller, and J. A. Peters, *J. Org. Chem.*, 30, 91 (1965).

the initial product but forms anion **8** which undergoes rearrangement to **9** and C-P cleavage to **10**¹⁴ (see section III.B for further discussion).



A series of stable heterocyclic phosphorus ylides has been prepared by an extension of the well-known ylide-forming reaction in which a triarylphosphine dihalide is reacted with an activated methylene group in presence of a base. The products obtained by this procedure are shown in Table II.¹⁵



B. NUCLEOPHILIC DISPLACEMENT REACTIONS BY PHOSPHORUS NUCLEOPHILES

1. Arbuzov and Michaelis-Becker Reactions on Heterocyclic Halides

The Arbuzov and Michaelis-Becker reactions are among the most widely used procedures for forming carbon-phosphorus bonds¹⁶ and have been successfully applied in the synthesis of heterocyclic phosphorus derivatives. Nucleophilic displacements on heterocyclic halides have been well studied for alkoxides, amines, etc., and a general order of reactivity has been recognized for these nucleophiles. Thus, at one extreme 1-halo-2,4,6-triazines have a high susceptibility to displacement of halide, whereas 2-halopyridines are relatively unreactive.¹⁷ For the case of displacements by trialkyl phosphites (Arbuzov) or dialkyl alkali metal phosphonates (Michaelis-Becker), a similar order of reactivity appears to exist although no kinetic measurements have been reported. Table III¹⁸⁻²⁹

Table II

Heterocyclic Phosphoranes¹⁵

Compound		Yield, %
	R ₁ = Ph; R ₂ = Me; R ₃ = Ph	98
	R ₁ = R ₂ = R ₃ = Ph	78
	R ₁ = Ph; R ₂ = Me; R ₃ = <i>p</i> -NO ₂ C ₆ H ₄	49
	R ₁ = <i>p</i> -MeOC ₆ H ₄ ; R ₂ = Me; R ₃ = Ph	67
	R ₁ = <i>p</i> -MeOC ₆ H ₄ ; R ₂ = Me; R ₃ = <i>p</i> -NO ₂ C ₆ H ₄	52
	(R ₁) ₃ = (Ph) ₂ (<i>p</i> -Me ₂ NC ₆ H ₄); R ₂ = Me; R ₃ = Ph	57
	R = Ph	22
	R = Et	92
		73
		95

summarizes the derivatives of heterocycles prepared by these reactions. It appears that 2-chloroquinoline¹⁹ and 2-chloropyrimidine²² are the least reactive substrates successfully reacted. It has been reported that 2-bromo- or 2-chloropyridine¹⁹ and 5-halopyrimidines²² fail to react in either the Arbuzov or Michaelis-Becker reactions. It would seem that these unreactive halides could be made more susceptible to displacement by protonation or quaternization. An alternative to this approach is the use of nickel salts as catalysts which apparently enhances the nucleophilicity of the trialkyl phosphites.^{29a} In this way aryl halides can be converted into aryl phosphonates in high yield (70-90%); for example, 2-bromothiophene is converted into diethyl 2-thienylphosphonate by reaction with triethyl phosphite in the presence of nickel bromide.^{29a}

The reaction of diphenylphosphinous chloride with 2-ethoxy-1,3-dioxane (**11**) (R = H) appears to proceed by a pathway related to the Arbuzov reaction in yielding the

(19) A. Burger, J. B. Clements, N. D. Dawson, and R. B. Henderson, *J. Org. Chem.*, **20**, 1383 (1955).

(20) W. Lorenz, German Patent 930212 (1955); *Chem. Abstr.*, **52**, 14700 (1958).

(21) V. L. Narayanan, J. Bernstein, and J. Williams, *J. Pharm. Sci.*, **55**, 217 (1966).

(22) G. M. Kosolapoff and C. H. Roy, *J. Org. Chem.*, **26**, 1895 (1961).

(23) R. M. Ismail, *Justus Liebigs Ann. Chem.*, **732**, 107 (1970).

(24) W. Hewertson, R. A. Shaw, and B. C. Smith, *J. Chem. Soc.*, 1670 (1963).

(25) Y. F. Shealy, R. F. Struck, J. D. Clayton, and J. A. Montgomery, *J. Org. Chem.*, **26**, 4433 (1961).

(26) B. A. Arbuzov and B. P. Lugovkin, *Zh. Obshch. Khim.*, **22**, 1193 (1952); *Chem. Abstr.*, **47**, 4871 (1953).

(27) A. Mustafa, M. M. Sidky, and F. M. Soliman, *Justus Liebigs Ann. Chem.*, **698**, 109 (1966).

(28) H. Gross, G. Englehardt, J. Freiberg, W. Buerger, and B. Costisella, *ibid.*, **707**, 35 (1967).

(29) H. Gross, J. Freiberg, and B. Costisella, *Chem. Ber.*, **101**, 1250 (1968).

(29a) P. Tavs, *ibid.*, **103**, 2428 (1970).

(15) J. J. Pappas and E. Gancher, *J. Heterocycl. Chem.*, **6**, 265 (1969).

(16) R. G. Harvey and E. DeSombre, *Top. Phosphorus Chem.*, **1**, 57 (1964).

(17) R. G. Shepherd and J. L. Fedrick, *Advan. Heterocycl. Chem.*, **4**, 262 (1965).

(18) Y. N. Ivashchenko, L. S. Sologub, S. D. Moshchitskii, and A. V. Kirsanov, *Zh. Obshch. Khim.*, **39**, 1695 (1969); *Chem. Abstr.*, **71**, 124600 (1969).

Table III

Phosphonates by Arbuzov and Michaelis-Becker Reactions





































Halide	Nucleophile	Product	Yield, %	Ref
Pentachloropyridine	(EtO) ₃ P	Diethyl 2,3,5,6-tetrachloropyridyl-4-phosphonate		18
2-Chloroquinoline	NaPO(OEt) ₂	Diethyl quinolyl-2-phosphonate	29	19
2-Chloro-4-methylquinoline	NaPO(OEt) ₂	Diethyl 4-methylquinolyl-2-phosphonate	25	19
	(EtO) ₃ P			20
	(EtO) ₃ P			21
				
R = Me	NaPO(OEt) ₂	R' = Et	28	22
R = H	P(OPr- <i>i</i>) ₃	R' = <i>i</i> -Pr	71.5	22
	P(OPr- <i>i</i>) ₃		70	22
				
Ar = C ₆ Cl ₅	P(OEt) ₃	R = Et	88	23
Ar = 2,3,4,6-tetrachlorophenyl	P(OCH ₂ CH ₂ Cl) ₃	R = CH ₂ CH ₂ Cl	92	23
Ar = 2,4,5-trichlorophenyl	P(OEt) ₃	R = Et	93	23
Ar = 2,4,6-tribromophenyl	P(OEt) ₃	R = Et	90	23
				
X = Ph	P(OMe) ₃	R = Me	81	24
X = Ph	P(OEt) ₃	R = Et	78	24
X = C ₆ H ₁₀ N	P(OEt) ₃	R = Et	66	24
				
X = Ph	P(OEt) ₃	R = Et	72	24
X = Ph	P(OMe) ₃	R = Me	76	24
X = C ₆ H ₁₀ N	P(OEt) ₃	R = Et	67	24
				
	P(OR) ₃			
		R = Me, Et, Pr, Bu, Bz	68-94	24
	MeOP(OPh) ₂	R = Ph	70	24
	EtOP(OPh) ₂	R = Ph	57	24
	EtOPPh ₂	OR = Ph	65	24

Table III (Continued)




Halide	Nucleophile	Product	Yield, %	Ref
	P(OPr-i) ₃		64	25
	NaPO(OEt) ₂		46	26
	(RO) ₃ P			
X = H		X = H; R = Me	81	27
X = Cl		X = PO(OR) ₂ ; R = Et	83	27
X = Cl		X = PO(OR) ₂ ; R = Pr	70	27
	(EtO) ₃ P		57	28
	(EtO) ₃ P		66	28
	(EtO) ₃ P		24	28
	(EtO) ₃ P		45	28
	(EtO) ₃ P		34	
	(MeO) ₃ P			
		R = Me, CH ₂ Cl, CHCl ₂ , CCl ₃	36-61	29
	P(OEt) ₃			
		R ₁ = CO ₂ Me; R ₂ = R ₃ = R ₄ = H	71	28
		R ₁ = R ₂ = Cl; R ₃ = R ₄ = H	86	
		R ₄ = Ph; R ₁ = R ₂ = R ₃ = H	70	
		R ₄ = CO ₂ Me; R ₁ = R ₂ = R ₃ = H	50	
		R ₁ = R ₂ = R ₃ = Cl; R ₄ = H	50	

phosphine oxide **12** (R = H). The 5,5-dimethyl derivative **11** (R = Me) reacts similarly.³⁰



Although diisopropyl 2-chloropyrimidin-4-phosphonate (**14**) is readily formed from 2,4-dichloropyrimidine (**13**) and triisopropyl phosphite, displacement of the second chlorine in an Arbuzov reaction with the same phosphite to give **15a** was

not successful.²² In view of the fact that this chlorine is readily displaced by a variety of nucleophiles to produce **15b-f**,²² it would seem that less vigorous conditions or a different phosphite would yield a pyrimidin-2,4-diphosphonate (such as **15a**). This view is reinforced by the results of the reactions of cyanuric chloride (**16**) with trialkyl phosphites.²⁴ When cyanuric chloride (**16**) is allowed to react with 2 equiv of tri-

(30) W. Dietsche, *Justus Liebig's Ann. Chem.*, 712, 21 (1968).




- a, X = PO(OPr-*i*)₂ d, X = NH₂
b, X = NHMe₂ e, X = OEt
c, X = NHEt₂ f, X = H



methyl phosphite, only trisubstituted triazine **17** (64%) and unreacted chloride (32%) are isolated, which suggests that the phosphonate group may in fact enhance the reactivity of the remaining chlorine to nucleophilic displacement. The reaction of cyanuric chloride (10 mol) with triethyl phosphite (1 mol) did result in monosubstitution product **18** (77%).²⁴ The reaction of 2-chloro-4,6-bistrichloromethyltriazine (**19**) with triethyl phosphite at 145° is reported to yield the phosphonate **21** rather than the expected phosphonate **20**.³¹ It is not worthwhile to speculate on a mechanism of formation of **21**, in view of the lack of a rigorous structure proof. It is possible that **20** could be obtained by the use of less vigorous conditions as, for example, in a solvent.²⁴

9-Chloroacridine (**22**) is reported to yield diethyl acridyl-9-phosphonate (**23**) upon reaction with triethyl phosphite,³² but



(31) H. Schroeder, *J. Amer. Chem. Soc.*, **81**, 5658 (1959).
(32) G. M. Kosolapoff, *ibid.*, **69**, 1002 (1947).



recent attempts to repeat this work gave the diphosphonate **24** as the only characterizable product.³³ This diphosphonate **24** is the product of the reaction of **22** and of **23** with diethyl sodiophosphonate.³³ These results suggest that diethyl acridyl-9-phosphonate (**23**) is the initial product of Arbuzov and Michaelis-Becker reactions of 9-chloroacridine but undergoes facile nucleophilic addition reactions (see section II.C).

2. Other Nucleophilic Displacements by Phosphorus


The displacement of chloride ion from 9-chloroacridine (**22**) by the anion of dicyclohexyl- or diphenylphosphine provides a synthesis of the tertiary phosphines **25** and **26** in yields of 16 and 48%, respectively.³⁴ Triphenylphosphine displaces bromide from tetrahydrothiophene (**27**), and the resulting phosphonium salt **28** can be converted, without isolation, into the phosphorane **29**.³⁵



All the syntheses of pyridine phosphonic acid derivatives recorded to date have involved nucleophilic displacements by phosphorus. Pyridyl-3-phosphonic acid (**31**) is formed by the


(33) D. Redmore, *J. Org. Chem.*, **34**, 1420 (1969).
(34) K. Issleib and L. Bruesehaber, *Z. Naturforsch.*, **20b**, 181 (1965).
(35) H. Zimmer, F. Haupter, S. P. Kharidia, H. Pauling, R. G. Gailey, T. Pampalone, T. C. Purcell, and R. Walter, *Tetrahedron Lett.*, 5435 (1968).

treatment of 3-pyridyldiazonium fluoroborate (**30**) with phosphorus trichloride in presence of cuprous bromide followed by work-up with water.³⁶ 2-Nitropyridine *N*-oxide (**32**) when heated with triethyl phosphite yields diethyl pyridyl-2-phosphonate (**34**) presumably *via* the *N*-oxide **33**.³⁷ 4-Nitropyridine *N*-oxide fails to undergo this reaction.³⁷ A series of pyridines **35a-d** have been converted into the pyridyl-2-phosphonate esters (**36a-d**), respectively, by the reaction sequence, oxidation, O-alkylation, and treatment with an alkali metal derivative of diethyl phosphonate.³⁸ The unsymmetrical pyri-




idine (**35e**) yields a mixture of 3-methylpyridyl-2-phosphonate **36g** and the 5-methyl compound **36e** in the ratio 6:1 and **35f** yields a mixture of **36h** and **36f** in a 3:1 ratio in the same reaction sequence. The predominance of substitution adjacent to the 3-alkyl substituent is reminiscent of the reaction of phenyllithium with 3-methylpyridine.³⁹ Only in the case of *N*-methoxy-2,6-dimethylpyridinium methosulfate (**37**) was 4 substitution observed. In this case attack on the methoxy

group was a major reaction yielding, by attack at hydrogen, **38** (47%) and, by attack at carbon, **39** (6%). The 4-phosphonate **40** was obtained in 24% yield.³⁸



Reaction of 1,3-dithiacyclohexane-2-thione (**41**) with excess trimethyl phosphite leads to the formation of ylide **42**, together with trimethyl thionophosphate (**43**).⁴⁰ Although a reaction sequence has not been specified, a possible pathway is shown. The ylide **42** upon heating at 70–80° for 24 hr is isomerized into phosphonate **44**.⁴⁰ The reaction of phthalic anhydride (**45**) with triisopropyl phosphite which yields phosphonate **47** is proposed to involve carbene **46** as an intermediate.⁴¹



C. NUCLEOPHILIC ADDITION REACTIONS

Carbon-phosphorus bonds can readily be formed by the addition of an anion (phosphonate or phosphine, for example) to a multiple bond, such as carbonyl, imine, or activated carbon-carbon double bond. The addition of dialkyl phosphonates to a carbonyl function is typically carried out under anhydrous conditions in the presence of a catalytic amount of base, such as

(36) R. D. Bennett, A. Burger, and W. A. Volk, *J. Org. Chem.*, **23**, 940 (1958).

(37) J. I. G. Cadogan, D. J. Sears, and D. M. Smith, *J. Chem. Soc. C*, 1314 (1969).














(38) D. Redmore, *J. Org. Chem.*, **35**, 4114 (1970).

(39) R. A. Abramovitch and J. G. Saha, *Advan. Heterocycl. Chem.*, **6**, 280 (1966).


(40) E. J. Corey and G. Markl, *Tetrahedron Lett.*, 3201 (1967).

(41) F. Ramirez, H. Yamanaka, and O. H. Basedow, *J. Amer. Chem. Soc.*, **83**, 173 (1961).

Table IV
Nucleophilic Addition Reactions

Carbonyl compound	Nucleophile	Product	Yield, %	Ref
	HPO(OMe) ₂			44
 R = H, Me, COCH ₃	HPO(OR') ₂ R' = Me, Et, <i>i</i> -Pr		62-85	45
	HPO(OR') ₂			
		R = H or Me; R' = Me R = Me; R' = Et	75-80 32	46 47
	NH ₃ , HPX(OR') ₂			
		X = O, S; R' = Et X = O; R' = Bu	42-43 52	47 47
	HPO(OR) ₂			
		X = O; R = Me, Et, Pr X = S; R = Me, Et, Pr	48-87 53-80	48 48
	(RO) ₂ POH			
		R = Me, Et, <i>i</i> -Pr, Pr, Bu, <i>i</i> -Bu	50-65	49
	(EtO) ₂ POH			50

tertiary amine or alkoxide.⁴² In strong aqueous base the reaction is reversed readily. The reaction of a carbonyl compound



with a dialkyl phosphonate and a primary amine or ammonia yields an α -amino phosphonate most probably *via* an imine since α -hydroxy phosphonates are not readily converted into

α -amino phosphonates by treatment with amines.⁴³ Table IV⁴⁴⁻⁵⁰ summarizes the preparation of heterocyclic phos-

(43) E. K. Fields, *J. Amer. Chem. Soc.*, **74**, 1528 (1952); N. S. Kozlov, V. D. Pak, and E. S. Elin, *Zh. Obshch. Khim.*, **39**, 2407 (1969); *Chem. Abstr.*, **72**, 79156 (1970). Ammonia can apparently convert α -hydroxy phosphonates into α -amino phosphonates: M. I. Kabachnik and T. Y. Medved, *Izv. Akad. Nauk SSSR, Otd. Khim. Nauk*, 868 (1953); *Chem. Abstr.*, **49**, 840 (1955).

(44) C. Benezra and G. Ourisson, *Bull. Soc. Chim. Fr.*, 1825 (1966).

(45) A. Mustafa, M. M. Sidky, and F. M. Soliman, *Tetrahedron*, **22**, 393 (1966).


(46) I. N. Azerbaev, T. G. Sarbaev, E. U. Gafurov, A. M. Aleshin, B. D. Abiyurov, and K. B. Erzhanov, *Tr. Inst. Khim. Nauk, Akad. Nauk Kaz. SSR*, **19**, 49 (1967); *Chem. Abstr.*, **68**, 95894 (1968).

(47) T. Y. Medved and M. I. Kabachnik, *Izv. Akad. Nauk SSSR, Otd. Khim. Nauk*, 1357 (1957); *Chem. Abstr.*, **52**, 7316 (1958).

(48) I. N. Azerbaev, T. G. Sarbaev, B. D. Abiyurov, and V. S. Basalitskaya, *Izv. Akad. Nauk Kaz. SSR, Ser. Khim.*, **18**, 56 (1968); *Chem. Abstr.*, **70**, 57948 (1969).


(49) B. P. Lugovkin, *Zh. Obshch. Khim.*, **31**, 3408 (1961); *Chem. Abstr.*, **57**, 3478 (1962).

(42) K. Sasse, in Houben-Weyl, "Organophosphorus Compounds," Part I, Georg Thieme Verlag, Stuttgart, 1963, pp 475-482.




phosphonates by addition to heterocyclic carbonyl and exocyclic imine groups.



Michael-type additions of dialkyl phosphonates to activated carbon-carbon double bonds yield heterocyclic phosphonates when this moiety is part of a heterocyclic ring. The addition of dimethyl phosphonate to coumarin (**48**) yields the dihydrocoumarylphosphonate **49** (20%) without reaction at the carbonyl carbon.⁵¹ The reported addition of diethyl phosphonate




to 2-cyanofuran (**50**) to yield diethyl 2-cyano-2,3-dihydrofuryl-3-phosphonate (**51**) is surprising in view of loss of resonance energy.⁵² The addition of diethyl phosphonate to Δ^3 -sulfolene dioxide (**52**) to yield the sulfolane dioxide (**53**) is to be expected in view of the reported addition of many other nucleophiles to this double bond.⁵³ The addition of phosphorus nucleophiles to acyclic imines is a well-known re-



action⁵⁴ which has recently been applied to cyclic imines. 3,4-Dihydroisoquinoline (**54**) adds diethyl phosphonate to yield the tetrahydroisoquinolylphosphonate **55** which has been characterized as its crystalline acetyl derivative **56**.⁵⁵ Acridine (**57**) undergoes 1,4 addition of phosphorus nucleophiles; thus the lithium salts of diethyl-, diphenyl-, and dicyclo-



hexylphosphines yield the 9,10-dihydroacridines **58**, **59**, and **60** (~30%).⁵⁴ Similarly, the dihydroacridylphosphonate **63** is formed in quantitative yield by the addition of diethyl phosphonate in presence of catalytic amounts of base to acridine (**57**). This same phosphonate **63** and homolog **64** are the products of the addition of diethyl sodiophosphonate to the salts **61** and **62**.³³ The dihydroacridylphosphonate **63** can be de-



hydrogenated to **66** which itself will add diethyl phosphonate to yield the diphosphonate **68**.³³ A recent report, without experimental details, discloses the reaction of acridine, quinoline, and pyridine with a trialkyl phosphite and acetyl chloride in a Reissert-type reaction.⁵⁶ In the case of acridine, the initial product is the dihydroacridine **65** which is hydrolyzed to acridyl-9-phosphonic acid (**67**). Similar sequences are reported to give quinolyl-2-phosphonic acid and pyridyl-2-phosphonic acid.⁵⁶ The melting points of these three acids do not correspond to previously reported values^{33,19,38} and, further, other attempts to carry out this type of reaction have been unsuccessful,⁵⁷ suggesting that the above reactions have not taken place as claimed.⁵⁶

(50) G. M. Kosolapoff, K. H. Bloss, and D. K. Myers, *Zh. Obshch. Khim.*, **38**, 1517 (1968); *Chem. Abstr.*, **70**, 11860 (1969).

(51) B. A. Arbusov and V. M. Soroastrova, *Izv. Akad. Nauk SSSR*, 681 (1955); *Chem. Abstr.*, **50**, 7109 (1956).

(52) A. N. Pudovik and N. I. Plakatina, *Sb. Statei Obshch. Khim.*, **2**, 831 (1953); *Chem. Abstr.*, **49**, 62814 (1955).

(53) R. L. McConnell and N. H. Shearer, U. S. Patent 2882278 (1959); *Chem. Abstr.*, **53**, 17149 (1959).


(54) E. K. Fields, *J. Amer. Chem. Soc.*, **74**, 1528 (1952); R. W. Layer, *Chem. Rev.*, **63**, 489 (1963).

(55) D. Redmore, unpublished work.


(56) A. K. Sheinkman, G. V. Samoilenko, and S. N. Baranov, *Zh. Obshch. Khim.*, **40**, 700 (1970); *Chem. Abstr.*, **73**, 14931 (1970).

(57) D. Redmore, unpublished work.

Among the products of deoxygenation of a series of nitrobenzenes **69–72** are the 3*H*-azepinyl-7-phosphonates **73–77**^{58,59} in the yields shown. The pathway postulated for the reaction involves deoxygenation of **69** to nitrene **78** and rearrangement



to imine **79** which is attacked by phosphite with ring opening to dipolar intermediate **80** and then to azepine **73** with ethylene liberation.⁵⁸ 2-Diethylaminoazepine (**81**) results from the trapping of **79** when the deoxygenation is carried out in presence of diethylamine⁵⁹ which suggests that higher yields of azepinylphosphonates (e.g., **73**) could be obtained by deoxy-



genation in the presence of diethyl phosphonate, since diethyl phosphonate should add more readily to **79** than triethyl phosphite. Generation of nitrenes (e.g., **78**) by other means, such as from azides, in the presence of dialkyl phosphonates might provide alternative routes to azepinylphosphonates.


Two routes to the phosphonate **84** have been described; one a nucleophilic addition to **83** and the second from **82** which could involve **83** as an intermediate.^{60,61} A facile isomerization of **84** to **85** is observed upon heating in ethanol. Several other dialkyl phosphonates and phosphorous acid esters were found to undergo this addition to yield products analogous to **84** and **85**, all of which reverted to thiamine hydrochloride in aqueous HCl.^{60,61}

(58) J. I. G. Cadogan, D. J. Sears, D. M. Smith, and M. J. Todd, *J. Chem. Soc. C*, 2813 (1969).

(59) R. J. Sundberg, B. P. Das, and R. H. Smith, *J. Amer. Chem. Soc.*, **91**, 658 (1969).

(60) A. Takamizawa, K. Hirai, and Y. Hamashima, *Tetrahedron Lett.*, 5081 (1967).

(61) A. Takamizawa, K. Hirai, Y. Hamashima, Y. Matsumoto, and S. Tanaka, *Chem. Pharm. Bull.*, **16**, 1761 (1968).



D. CYCLOADDITION REACTIONS

A series of azetidone phosphonates **88** has been prepared by the cycloaddition of the imine phosphonates **86** to the ketene derived from the acid chloride **87** in the presence of triethylamine⁶² in an extension of a well-known reaction.⁶³

The 1,3-dipolar cycloaddition reaction of 1,3 dipoles with alkenes has been used in the synthesis of heterocyclic phosphorus derivatives in two different ways: the first in which the phosphorus substituent is present in the alkene (or alkyne) and the second in which it is attached to the 1,3 dipole. Diisopropyl ethynylphosphonate (**89**) upon reaction with diazomethane is converted into the pyrazolyl phosphonate **90** in 28% yield.⁶⁴ The ethynyldiphosphonate **91** is converted into the pyrazole **92** in 95% yield in a similar reaction with diazomethane.⁶⁵ At low temperatures diphenyldiazomethane adds to diethyl vinylphosphonate (**93**) to yield dihydropyrazolylphosphonate **94** (73%), although at higher temperatures the cyclopropylphosphonate **95** is the product.⁶⁶ Triphenylvinylphosphonium bromide (**96**) can also be used as a dipolarophile with diazoalkanes. Thus, with diphenyldiazomethane the product in quantitative yield is 5,5-diphenyl-2-pyrazolin-3-yltriphenylphosphonium bromide (**97**), while with diazomethane the product is reported to be 2-pyrazolin-3-yltriphenylphosphonium bromide (**98**).⁶⁷ 1,2,5-Triphenylphosphole oxide (**99**) serves as a dipolarophile for ethyl diazoacetate or diazomethane in which case the products are dihydropyrazole compounds **100** and **101**, respectively.⁶⁸ An alternative route to dihydropyrazole phosphorus compounds **104** has been provided in the addition of the nitrile imine **103** to vinylphosphorus compounds **102**.⁶⁹

(62) L. Paul and K. Zieloff, *Chem. Ber.*, **99**, 1431 (1966).

(63) R. Huisgen, R. Grashey, and J. Sauer, "The Chemistry of Alkenes," S. Patai, Ed., Interscience, London, 1964, pp 802–805.

(64) B. C. Saunders and P. Simpson, *J. Chem. Soc.*, 3351 (1963).


(65) D. Seyferth and J. D. H. Paetsch, *J. Org. Chem.*, **34**, 1483 (1969).

(66) A. N. Pudovik, R. D. Gareev, and L. I. Kuznetsova, *Zh. Obshch. Khim.*, **39**, 1536 (1969); *Chem. Abstr.*, **71**, 113049 (1969).







(67) E. E. Schweizer, C. S. Khim, and R. A. Jones, *Chem. Commun.*, 39, 1584 (1970).

(68) I. G. M. Campbell, R. C. Cookson, M. B. Hocking, and A. N. Hughes, *J. Chem. Soc.*, 2184 (1965).


(69) I. G. Kolokol'tseva, V. N. Chistoketov, B. I. Ionin, and A. A. Petrov, *Zh. Obshch. Khim.*, **38**, 1248 (1968); *Chem. Abstr.*, **69**, 96834 (1968).




R	R ₁	R ₂	Yield, %
Ph	H	Me, Et	18, 46
Ph	Me	Me, Et	29, 32
Ph	Cl	Me, Et	17, 28
Ph	Br	Me, Et	22, 27
Ph	OMe	Me, Et	36, 40
Me	H	Et	24



N-Phenylsydnone (**108**) can be used as a 1,3-dipolar compound in reaction with propynylphosphonate **107** yielding pyrazolylphosphonate **109** (53%) with extrusion of carbon dioxide.⁷⁰ The reaction between diphenylethynylphosphine



(**110a**) and diphenylnitrilimine (**103**) does not yield the expected pyrazole **112**, but rather the phosphonia diazacyclohexadiene (**111**) (98%).⁷¹ Additional acetylenes **110b** and **110c** yield **111b** and **111c** in the reaction and, further, diphenylvinylphosphine undergoes a similar reaction yielding **113**.⁷¹ This unexpected reaction of the unsaturated phosphines merits fur-



Dihydroisoxazoles **106** result when nitrile oxide **105** is substituted as the 1,3 dipole.⁶⁹


(70) A. N. Pudovik and N. G. Khusainova, *Zh. Obshch. Khim.*, **40**, 697 (1970); *Chem. Abstr.*, **73**, 14923 (1970).

(71) I. G. Kolokoltseva, V. N. Chistokletov, and A. A. Petrov, *Zh. Obshch. Khim.*, **40**, 574 (1970); *Chem. Abstr.*, **73**, 25582 (1970).

cycloaddition reaction using vinylphosphorus compounds as dipolarophiles as a method to heterocyclic phosphorus compounds seems to offer possibilities, many as yet unexamined.⁷²


The preparation of heterocyclic phosphorus compounds by the cycloadditions of phosphorus-containing dipoles to olefins has been described for a series of α -diazophosphorus compounds and activated olefins as shown in Table V.⁷³⁻⁷⁵

Table V
Addition of Phosphorus-Containing Dipoles to Activated Olefins



R	R ₁	R ₂	R ₃	Yield, %	Ref
OMe	Ph	H	COCH ₃	92	73
OMe	Ph	CO ₂ Et	CO ₂ Et	46	73
OMe	Me	H	COCH ₃ , CN, CO ₂ Et	90-98	73
Ph	Ph	H	COCH ₃	89	74
OEt	<i>p</i> -NO ₂ C ₆ H ₄	H	COCH ₃	42	75
OEt	Ph	H	COCH ₃	55	75

The addition of tosylazide to the phosphonates **114a** and **114b** almost certainly yields the triazolines **115a** and **115b** which, under the reaction conditions, yield diazophosphonate **117** and triazole **116**, respectively.^{75,76}




E. CONDENSATION REACTIONS


1. Epoxyethylphosphorus Compounds

Many of the classical condensation methods of heterocyclic synthesis have been carried out with phosphorus-bearing reactants to provide syntheses of heterocyclic phosphorus deriva-

tives. Oxirane phosphorus derivatives figure particularly prominently in this section. Interest in these structures has been stimulated by the recent discovery of the antibiotic phosphonmycin, ($-1R,2S$)-1,2-epoxypropylphosphonic acid (**118**).⁷⁷ The numerous syntheses disclosed in recent patents attest to the potential of this phosphonic acid.⁷⁸ Although little experimental detail is available on these syntheses, some of them are discussed below because of current interest.



The treatment of halohydrins with base is a well-known method for the formation of epoxides⁷⁹ and has been exploited in the synthesis of epoxy phosphonates. The halohydrin **120a** derived from the treatment of *cis*-1-propenylphosphonic acid (**119**) with sodium hypochlorite is converted into phosphonmycin (**118**) with sodium hydroxide.⁸⁰



Similarly, the dimethyl ester **120b** is converted into the phosphonmycin ester **121b** which is demethylated by reaction with trimethylchlorosilane to the acid **118**.⁸¹ Halohydrins **124** formed by heating a dialkyl phosphonate **123** with an α -chloro-carbonyl compound **122** are converted by treatment with potassium hydroxide into epoxy phosphonates **125** as summarized in Table VI.⁸²⁻⁸⁴ In the steroid field the cholestane deriva-

(77) B. G. Christensen, W. J. Leanza, T. R. Beattie, A. A. Patchett, B. H. Arison, R. E. Ormond, F. A. Kuehl, G. Albers-Schonberg, and O. Jardetzky, *Science*, **166**, 123 (1969).

(78) B. Christensen, German Offen. 1924104 (1970); *Chem. Abstr.*, **72**, 90630 (1970); J. M. Chmerda and E. J. Glamkowski, German Offen. 1924118 (1970); *Chem. Abstr.*, **72**, 132953 (1970); R. A. Firestone and E. J. Glamkowski, German Offen. 1924105 (1970); *Chem. Abstr.*, **72**, 132952 (1970); E. F. Shoenevaldt, German Offen. 1924231 (1970); *Chem. Abstr.*, **72**, 132972 (1970); M. Sletzing and S. Karady, German Offen. 1924149 (1970); *Chem. Abstr.*, **72**, 90628 (1970); R. A. Firestone, German Offen. 1924098 (1970); *Chem. Abstr.*, **72**, 90629 (1970); J. M. Chmerda and E. J. Glamkowski, German Offen. 1924173 (1970); *Chem. Abstr.*, **72**, 43871 (1970).

(79) A. Rosowsky in "Heterocyclic Compounds with Three and Four Member Rings," A. Weissberger, Ed., Interscience, New York, N. Y., 1964, pp 94-106.

(80) N. N. Girotra and N. L. Wendler, *Tetrahedron Lett.*, 4647 (1969).

(81) P. I. Pollak, B. G. Christensen, and N. L. Wendler, German Offen. 1924169 (1970); *Chem. Abstr.*, **72**, 100882 (1970).

(82) V. S. Abramov and R. N. Savitseva, *Khim. Org. Soedin. Fosfora*, **129** (1967); *Chem. Abstr.*, **69**, 67465 (1968).

(83) B. A. Arbuzov, V. S. Vinogradova, and N. A. Polezhaeva, *Dokl. Akad. Nauk SSSR*, **111**, 107 (1956); *Chem. Abstr.*, **51**, 8001 (1957).

(84) P. A. Kirpichnikov, A. S. Kapustina, and G. N. Tokareva, *Tr. Kazan. Khim.-Tekhnol. Inst.*, **33**, 188 (1964); *Chem. Abstr.*, **66**, 2619 (1967).

(72) Reference 63, pp 806-874.

(73) D. Seyferth, P. Hilbert, and R. S. Marmor, *J. Amer. Chem. Soc.*, **89**, 4811 (1967).


(74) M. Regitz and W. Anschutz, *Chem. Ber.*, **102**, 2216 (1969).

(75) M. Regitz, W. Anschutz, and A. Liedhegener, *ibid.*, **101**, 3734 (1968).


(76) M. Regitz and W. Anschutz, *Justus Liebigs Ann. Chem.*, **730**, 194 (1969).

Table VI

Structure of halohydrin 124	Yield of 125 , %	Ref
$R_1 = R_2 = H; R_3 = Me, Et, i\text{-Pr}, Bu$	25-34	82
$R_1 = H; R_2 = Me; R_3 = Et$...	83
$R_1 + R_2 = -(CH_2)_4-; R_3 = Me, Et, Pr, Bu$	30-65	84



tives **126a** or **126b** are readily converted into the epoxy **127** with sodium methoxide in high yields (85 and 75%, respectively).⁸⁵



Reaction of α -halo ketones with certain nucleophiles is a standard epoxide synthesis⁸⁶ which can be used with appropriate phosphorus nucleophiles. Alkali metal derivatives of dialkyl phosphonates attack α -halo ketones primarily at the carbonyl carbon atom forming an alkoxide anion which displaces the α -halogen. In some cases, however, attack at carbonyl oxygen can be a competitive reaction which results in formation of an enol phosphate ester.⁸⁷⁻⁸⁹ Syntheses by this route are summarized in Table VII.⁹⁰⁻⁹³

A variation of this procedure is exemplified in the conversion of the steroidal α -tosyloxy ketone **128** to epoxy phosphonate **129** in 69% yield by treatment with diethyl sodiophosphonate.⁹⁴ The reaction of diethyl ethylsodium phosphinate (**130**) on α -halo ketones is reported to take different courses

(85) C. Benezra and G. Ourisson, *Bull. Soc. Chim. Fr.*, 624 (1967).

(86) Reference 79, pp 119-147.

(87) A. Meisters and J. M. Swan, *Aust. J. Chem.*, **18**, 168 (1965).

(88) B. A. Arbutov, V. S. Vinogradova, N. A. Polezhaeva, and A. V. Shamsutdinova, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1380 (1963); *Chem. Abstr.*, **59**, 15306 (1963).

(89) T. Y. Medved and M. I. Kabachnik, *Izv. Akad. Nauk SSSR, Otd. Khim. Nauk*, 1357 (1957); *Chem. Abstr.*, **52**, 7316 (1958).

(90) G. Sturtz, *Bull. Soc. Chim. Fr.*, 2333 (1964).

(91) B. A. Arbutov, V. S. Vinogradova, and N. A. Polezhaeva, *Izv. Akad. Nauk SSSR, Otd. Khim. Nauk*, 41 (1959); *Chem. Abstr.*, **53**, 15035 (1959).


(92) M. Sprecher and D. Kost, *Tetrahedron Lett.*, 703 (1969).

(93) D. Redmore, unpublished.

(94) S. Hirai, R. G. Harvey, and E. V. Jensen, *Tetrahedron*, **22**, 1625 (1966).


(95) B. A. Arbutov, V. S. Vinogradova, and M. A. Zvereva, *Izv. Akad. Nauk SSSR, Otd. Khim. Nauk*, 1772 (1960); *Chem. Abstr.*, **55**, 16398 (1961).

Table VII


Epoxyethylphosphonates from α -Halo Ketones

α -Halo ketone	Phosphonate	Yield, %	Ref
$R_1 = Me; R_2 = R_3 = H$	$R_4 = Et$	65	83, 90, 91
$R_1 = Me; R_2 = R_3 = H$	$R_4 = Bu$	69	90
$R_1 = Et; R_2 = R_3 = H$	$R_4 = Et$	71	90
$R_1 = Ph; R_2 = R_3 = H$	$R_4 = Et^a$	55	87, 88
$R_1 = Ph; R_2 = Me; R_3 = H$	$R_4 = Et$...	88, 92
$R_1 = Ph; R_2 = R_3 = Me$	$R_4 = Et$...	^b 88
$R_1 + R_3 = -(CH_2)_4-; R_2 = H$	$R_4 = Et$	79	93
$R_1 = R_2 = Ph; R_3 = H$	$R_4 = Me$	92	
$R_1 = t\text{-Bu}; R_2 = Ph; R_3 = H$	$R_4 = Et$	92	
$R_1 = p\text{-MeOC}_6\text{H}_4; R_2 = Ph; R_3 = H$	$R_4 = Me$	92	

^a In liquid ammonia also gives $CH_2=C(Ph)OPO(OEt)_2$. ^b No epoxide, only $Me_2C=C(Ph)OPO(OEt)_2$.



depending on the ketone structure.^{95,96} This nucleophile with α -chloroacetone gives epoxyphosphinate **131**,⁹⁶ but with α -chlorocyclohexanone the product is enol phosphonate, ethyl cyclohexenyl ethylphosphonate (**132**).⁹⁶ The dependence of the reaction course upon the structure of the α -halo ketone requires more examples for clarification.




Although the reaction of α -halocarbonyl compounds with phosphorus nucleophiles has been satisfactorily applied in the synthesis of epoxyphosphonates, side reactions leading to unsaturated products can intervene as seen above. The Darzens procedure of reaction of halomethylphosphorus compounds **134** with carbonyl compounds yields epoxyphosphorus deriva-

(96) B. A. Arbutov, V. S. Vinogradova, and M. A. Zvereva, *Izv. Akad. Nauk SSSR, Otd. Khim. Nauk*, 1981 (1960); *Chem. Abstr.*, **55**, 16398 (1961).

tives **135** without side reactions. The halomethylphosphorus reactant used has been exclusively the readily available chloromethylphosphonic acid methyl or ethyl ester (**134**), while the carbonyl compound **133** is limited to ketones and arylaldehydes.⁹⁷ Table VIII summarizes the results.⁹⁸⁻¹⁰⁰

Table VIII


Epoxyphosphonates by Darzens Reaction



Carbonyl compound	Phosphonate	Yield, %	Ref
R ₁ = CH ₂ CH ₂ CHMe ₂ ; R ₂ = Me	R ₃ = Me	37	98
R ₁ = <i>c</i> -C ₆ H ₁₁ ; R ₂ = H	R ₃ = Et	43	98
R ₁ = Ph; R ₂ = Me	R ₃ = Me	36	98, 99
R ₁ = Ph; R ₂ = H	R ₃ = Me, Et	61-65	98, 99
R ₁ = <i>p</i> -MeC ₆ H ₄ ; R ₂ = H	R ₃ = Me	68	98
R ₁ = <i>p</i> -MeOC ₆ H ₄ ; R ₂ = H	R ₃ = Me	10	98
R ₁ = R ₂ = Me	R ₃ = Me	...	99
R ₁ = R ₂ = Ph	R ₃ = Me, Et	...	99
R ₁ = R ₂ = -(CH ₂) ₅ -	R ₃ = Et	31	99, 100


Direct epoxidation of α,β -unsaturated phosphonates with either acidic or basic reagents has been successful in the synthesis of epoxyphosphonates. With trifluoroperacetic acid, the initially formed epoxide may be ring opened unless mild conditions are employed.¹⁰¹ In the case of epoxidation with *tert*-butyl peroxide Michael addition of butoxide to olefin may be a significant side reaction.¹⁰² The epoxyphosphonates synthesized by epoxidation reactions are summarized in Table IX.^{103,104}

An additional procedure used in the preparation of epoxyphosphonates is the reaction of diphenyldiazomethane with dimethyl acetylphosphonate (**136a**) or dimethyl benzoylphosphonate (**136b**) which yields **137a** or **137b**, respectively.¹⁰⁵ A useful supplement to the Darzens condensation, applicable to aliphatic aldehydes, has been used to convert diethyl chloromethylphosphonate (**138**) into phosphonomycin (**118**).¹⁰⁶ Reaction of **138** with dimethyl sulfide yields sulfonium salt




a, R = Me; b, R = Ph

139 which *via* its ylide is converted into epoxyphosphonate **140** by reaction with acetaldehyde and thence to phosphonomycin (**118**).




2. Five-Membered Ring Compounds

Although the reaction of dialkyl sodiophosphonates with α -halo and γ -halo ketones yields cyclic phosphonates (epoxyethylphosphonates and tetrahydrofurylphosphonates, respectively), the reaction with β -halo ketones yields only unsaturated acyclic products such as **141**.⁹⁰ A γ -chloro ketone, 1-chloropentan-4-one (**142**), upon reaction with alkali metal phosphonates **143** and **144** gives good yields of 2-methyltetra-



hydrofurylphosphonates **145** and **146** (79 and 66%, respectively).^{90,91} In the case of 1-chlorohexanone-4 (**147**), the ex-



pected 2-ethyltetrahydrofurylphosphonate **148** is formed in only 14% yield, and phosphonate **149** is formed in significant amounts (17%).⁹⁰ In order to form the tetrahydrofuran, the phosphorus nucleophile must attack the carbonyl carbon to generate the alkoxide ion **150** which intramolecularly displaces chloride ion. It is surprising that sufficiently greater steric hindrance is introduced in **147** over **142** that direct halide displacement now competes favorably with attack at the carbonyl carbon.




(97) Cf. M. S. Newman and B. J. Magerlein, *Org. React.*, **5**, 413 (1949).(98) V. F. Martynov and V. E. Timofeev, *Zh. Obshch. Khim.*, **34**, 3890 (1964); *Chem. Abstr.*, **62**, 19457 (1965).(99) R. H. Churi and C. E. Griffin, *J. Amer. Chem. Soc.*, **88**, 1824 (1966).(100) V. F. Martynov and V. E. Timofeev, *Zh. Obshch. Khim.*, **32**, 3449 (1962); *Chem. Abstr.*, **58**, 9121 (1963).(101) K. Hunger, *Chem. Ber.*, **101**, 3530 (1968).(102) C. E. Griffin and S. K. Kundu, *J. Org. Chem.*, **34**, 1532 (1969).(103) E. J. Glamkowski, G. Gal, R. Purick, A. J. Davidson, and M. Sletzing, *ibid.*, **35**, 3510 (1970).(104) J. M. Chernerda and M. Sletzing, German Offen. 1924172 (1970); *Chem. Abstr.*, **72**, 90631 (1970).(105) A. N. Pudovik, R. D. Gareev, and L. A. Stabrovskaya, *Zh. Obshch. Khim.*, **40**, 698 (1970); *Chem. Abstr.*, **73**, 14933 (1970).(106) B. G. Christensen and R. A. Firestone, German Offen. 1924135 (1969); *Chem. Abstr.*, **72**, 43870 (1970).

Table IX
Epoxyethylphosphonates by Oxidation of Olefins




Reagent	Epoxyde	Yield, %	Ref
CF ₃ CO ₂ H	R ₁ = H; R ₂ + R ₃ = -(CH ₂) ₄ -; R ₄ = Me	73	101
CF ₃ CO ₂ H	R ₁ = R ₂ = R ₃ = H; R ₄ = Et	67	101
MeOH-H ₂ O ₂ , pH 9.5-10	R ₁ = R ₂ = R ₃ = H; R ₄ = Et	10	102
<i>t</i> -BuOOH-Triton B	R ₁ = R ₂ = R ₃ = H; R ₄ = Et	62	102
<i>t</i> -BuOOH-Triton B	R ₁ = R ₂ = R ₃ = H; R ₄ = Me	18	102
CF ₃ CO ₂ H	R ₁ = R ₂ = Me; R ₃ = H; R ₄ = Et	76	101
CF ₃ CO ₂ H	R ₁ = R ₂ = Me; R ₃ = H; R ₄ = Et	65	101
<i>m</i> -ClC ₆ H ₄ CO ₂ H	R ₁ = R ₃ = Ph; R ₂ = H; R ₄ = Me	..	92
<i>m</i> -ClC ₆ H ₄ CO ₂ H	R ₁ = Me; R ₂ + R ₃ = -(CH ₂) ₄ -; R ₄ = Me	..	92
H ₂ O ₂ -NaWO ₄ , pH 5.0	R ₁ = Me; R ₂ = R ₃ = H; R ₄ = H	64	77, 103
CF ₃ CO ₂ H	R ₁ = Me; R ₂ = R ₃ = Cl; R ₄ = PhCH ₂	..	104

Levulinic acid (**151**) reacts with diethyl phosphonate to form a rather unstable adduct **152** which can be cyclized with *p*-toluenesulfonic acid to furanone **153**.¹⁰⁷




Generation of the anion from allylphosphonate **154** in a protic solvent in presence of carbonyl compounds yields mixtures of 2,3-dihydrofuryl-3-phosphonates **155** and epoxides **156**.¹⁰⁸ In aprotic solvent the Wadsworth-Emmons reaction is the major pathway leading to butadienes.¹⁰⁸



2,5-Diphenylfuryl-3-phosphonic acid (**158**) is conveniently obtained by reaction of phosphorus trichloride with dibenzoyl-ethylene (**157**) in the presence of acetic anhydride followed by careful hydrolytic work-up.¹⁰⁹ A less efficient alternative is radical-induced¹⁰⁹ or base-catalyzed addition¹¹⁰ of diethyl

phosphonate to **157** yielding phosphonate **159** which with strong acid is cyclized to the acid **158**. The furylphosphonic



acid **158** undergoes a Diels-Alder addition with maleic anhydride to yield phosphonic acid **160**.¹⁰⁹ Subjecting phosphonate **161a** or phosphinate **161b** to reaction with phenylhydrazine under Fisher indole synthesis conditions affords the indolyl-2-phosphonate **162a** or -2-phosphinate **162b**, respectively, in low yield.^{111,112} One would have anticipated the corresponding 3-substituted indole as the product;¹¹³ thus the structure assignment may be incorrect (see section III.B).

Acetals of formyl phosphonates or phosphinates **164** have been utilized in condensations with amino, hydroxy, or mercapto 1,2-disubstituted benzene derivatives **163** to yield a

(107) J. A. Cade, *J. Org. Chem.*, **23**, 1372 (1958).

(108) G. Lavielle and H. Normant, *C. R. Acad. Sci., Ser. B*, **270**, 86 (1970).


(109) C. E. Griffin and J. T. Brown, *J. Org. Chem.*, **26**, 853 (1961).

(110) N. Kreutzkamp and W. Mengel, *Arch. Pharm. (Weinheim)*, **300**, 389 (1967).

(111) A. I. Razumov and P. A. Gurevich, *Zh. Obshch. Khim.*, **37**, 1615 (1967); *Chem. Abstr.*, **68**, 39730 (1968).


(112) A. I. Razumov and P. A. Gurevich, *Tr. Kazan. Khim. Tekhnol. Inst.*, **480** (1967); *Chem. Abstr.*, **70**, 20160 (1969).

(113) B. Robinson, *Chem. Rev.*, **63**, 373 (1963); **69**, 227 (1969).





a, R = OEt; b, R = Ph

series of benzoheterocyclic phosphorus compounds **165**.¹¹⁴⁻¹¹⁸ The yields of benzoxazole and benzothiazole derivatives are acceptable, but very poor yields of benzimidazoles are obtained. The source of oxidant in these reactions is not apparent, and in fact the reactions are carried out under a nitrogen blanket. Catechol and naphthalene-2,3-diol upon reaction




Phosphorus reactant (164)		Yield, %	Ref
X = O	R ₁ = R ₂ = Et	68	114, 115
X = S	R ₁ = R ₂ = Et	30	116
X = NH	R ₁ = R ₂ = Me	7.5	117
X = NH	R ₁ = Et; R ₂ = <i>p</i> -Cl-C ₆ H ₄	6.5	117

with formyl phosphonate **166** are converted into dioxolane phosphonates **167** and **168**, respectively, in a similar reac-





tion.¹¹⁸ Phenacyldiphenylphosphine oxide (**169**) condenses with imine **170** to form ketoimine **171** which upon reaction with phenyl- or *p*-nitrophenylhydrazine yields the diphenylpyrazolylphosphine oxides **172a** and **172b**.¹¹⁹

The Arbuzov reaction of triethyl phosphite and malonic acid derivatives **173** yields phosphonate **174** which upon condensation with phenylhydrazine is converted into pyrazolone phosphonate **175** in low yield.¹²⁰



a, Ar = Ph (56%); b, Ar = *p*-NO₂C₆H₄ (55%)



Diethyl acetylphosphonate (**176**), readily obtained from chloroacetone and triethyl phosphite by an Arbuzov reaction, after bromination, condenses with thiourea or thioformamide to thiazole phosphonates **177a** and **177b**. Both are rather unstable unless converted into salts.¹²¹

Conventional condensations of 3,4-diaminophenylphosphonic acid (**178**) with formic or acetic acid provides syntheses of benzimidazolyl-5-phosphonic acids **179a** and **179b**.¹²²

Upon reaction with tosyl azide, acetamides **180a** and **180b** are converted into triazolyl phosphonate **181a**⁷⁵ and triazolyl-diphenylphosphine oxide **181b**,⁷⁴ respectively, in a reaction involving diazo intermediates. The pyridyltriazolylphosphine oxide **183** is formed from **182** by a similar sequence.⁷⁴

3. Six-Membered Ring Compounds

A few phosphonic acid derivatives of six-membered rings have been prepared, mostly by straightforward condensation procedures. Diethyl 4-phenyltetrahydropyran-4-phosphonate (**184**) is the product from the reaction of diethyl benzylphosphonate and bis(β -chloroethyl) ether in the presence of sodamide.¹²³ Magnesium ethoxide brings about cyclization of the

(114) A. I. Razumov, B. G. Liorber, and P. A. Gurevich, *Zh. Obshch. Khim.*, **37**, 2782 (1967); *Chem. Abstr.*, **69**, 43977 (1968).

(115) A. I. Razumov, P. A. Gurevich, B. G. Liorber, and T. B. Borisova, *Zh. Obshch. Khim.*, **39**, 392 (1969); *Chem. Abstr.*, **71**, 115230 (1969).

(116) A. I. Razumov, B. G. Liorber, and P. A. Gurevich, *Zh. Obshch. Khim.*, **38**, 199 (1968); *Chem. Abstr.*, **69**, 52216 (1968).

(117) A. I. Razumov and P. A. Gurevich, *Zh. Obshch. Khim.*, **37**, 1620 (1967); *Chem. Abstr.*, **68**, 39731 (1968).



(118) A. I. Razumov and P. A. Gurevich, *Zh. Obshch. Khim.*, **38**, 944 (1968); *Chem. Abstr.*, **69**, 67467 (1968).

(119) H. G. Henning, G. Petzold, and G. Busse, *Z. Chem.*, **8**, 302 (1968).
(120) M. H. Maguire, R. K. Ralph, and G. Shaw, *J. Chem. Soc.*, 2299 (1958).


(121) N. D. Dawson and A. Burger, *J. Amer. Chem. Soc.*, **74**, 5312 (1952).

(122) R. W. Bost and L. D. Quin, *J. Org. Chem.*, **18**, 358 (1953).



(123) P. Malatesta and A. Ciaramella, *Ann. Chim. (Rome)*, **51**, 230 (1961).





a, R = OEt; b, R = Ph




ester **185** into the tetrahydrothiopyranylphosphonate **186** in 56% yield.¹²⁴



The Skraup quinoline synthesis has successfully been applied to quinolyl-6-phosphonic acid (**188**) from *p*-amino-phenylphosphonic acid (**187**).¹²⁵ Two rather low yield preparations of pyrimidylphosphonic acid derivatives have been reported: uracil-6-phosphonic acid (**189**) is obtained by condensing **174** with urea¹²⁰ while pentane-2,4-dione 3-phosphonate (**190**) with urea yields the pyrimidyl-5-phosphonate (**191**).¹²⁶ Tetraethyl 1,4-dioxane-2,6-phosphonate (**194**) re-




portedly is formed by condensing diphosphonate **192** and chloro ether **193**.¹²⁷



F. RADICAL REACTIONS^{127a}

Syntheses of phosphorus derivatives of heterocyclic systems by reactions involving radical intermediates are very few and of little importance. However, dibenzofuran (**195a**) and carbazole (**195b**) undergo phosphonation in good yield when heated with diethyl phosphonate in the presence of *tert*-butyl peroxide. After work-up by hydrolysis the products in each case are a mixture of isomeric phosphonic acids **196a** and **196b** not readily separable.¹²⁸ The formation of diethyl 2-



phenylindolyl-3-phosphonate (**198**) by deoxygenation of 1-hydroxy-2-phenylindole (**197**) with triethyl phosphite is postulated to proceed by radical intermediates.¹²⁹ The generation of aryl radicals by photolysis of aryl iodides in the presence of trimethyl or triethyl phosphite gives good yields of aryl phosphonates.¹³⁰ In the case of 2-iodofuran (**199a**) and 2-iodothiophene (**199b**), however, rather low yields of the corresponding

(127) A. N. Pudovik, E. A. Ishmaeva, and I. V. Shergina, USSR Patent 253804 (1969); *Chem. Abstr.*, **72**, 90618 (1970).

(127a) For a review of radical reactions of organophosphorus compounds see C. Walling and M. S. Pearson, *Top. Phosphorus Chem.*, **3**, 1 (1966).

(128) E. F. Jason and E. K. Fields, *J. Org. Chem.*, **27**, 1402 (1962).


(129) R. J. Sundberg, *ibid.*, **30**, 3604 (1965).

(130) R. Obrycki and C. E. Griffin, *ibid.*, **33**, 632 (1968).

(124) F. Korte and F. F. Wiese, *Chem. Ber.*, **97**, 1963 (1964).

(125) G. M. Kosolapoff, *J. Org. Chem.*, **21**, 1046 (1956).

(126) A. N. Pudovik and T. M. Moshkina, *Zh. Obshch. Khim.*, **27**, 1611 (1957); *Chem. Abstr.*, **52**, 3713 (1958).




a, X = O; b, X = S

phosphonates **200a** and **200b**, respectively, are obtained owing to difficulty in obtaining pure iodides, and also owing to instability of the product phosphonates **200** under the photolysis conditions.¹³⁰ A superior method of preparation of these phosphonates is the nickel salt catalyzed Arbuzov reaction^{29a} (section II.B).

G. ELECTROPHILIC SUBSTITUTION BY PHOSPHORUS


Although many heterocyclic rings, *e.g.*, pyridine, are rather inert to electrophilic attack, the more electron-rich systems, *e.g.*, thiophene, readily undergo electrophilic substitution. For example, the Friedel-Crafts reaction of phosphorus trichloride and thiophene in the presence of stannic chloride yields 2-thienylphosphonous dichloride (**201**) in 50% yield.¹³¹ This procedure is a modification of that described by Sachs in 1892.¹³² The phosphonous dichloride **201** can be converted by standard transformations into a series of derivatives (Scheme I).



2-Methylindolyl-3-phosphonic dichloride (**202**) has been prepared by reaction of phosphorus pentachloride with 2-methylindole followed by mild hydrolytic work-up.^{132a}

The following preparations are discussed in this section, although no strong evidence is available to define the mechanism. The reaction of phosphorus pentasulfide with excess thiophene is reported, without supporting physical data, to

yield 2-thienylthiophosphonic anhydride (**203**).¹³³ A similar type of substitution reaction results from the treatment of pyrazole **204** with phosphoryl chloride in which the product after esterification is the ester **205**.¹³⁴




III. Chemistry

A. CARBON-PHOSPHORUS BOND CLEAVAGE

1. Wittig and Wadsworth-Emmons Reactions

Appropriate heterocyclic phosphorus compounds undergo the title reactions in a manner similar to the corresponding acyclic and alicyclic compounds.¹³⁵ Thus, pyrazoline **206** with base



a, R = Ph; b, R = Me

(131) M. Ventov, L. David, and E. D. Bergmann, *J. Chem. Soc.*, 4750 (1964).

(132) H. Sachs, *Chem. Ber.*, **25**, 1514 (1892).


(132a) J. C. Powers, *J. Org. Chem.*, **31**, 2627 (1966).

(133) H. Hirai and H. Yoshiska, German Offen. 1896105 (1969); *Chem. Abstr.*, **71**, 50213 (1969).


(134) I. I. Grandberg and A. N. Kost, *Zh. Obshch. Khim.*, **31**, 129 (1961); *Chem. Abstr.*, **55**, 22292 (1961).

(135) A. W. Johnson, "Ylid Chemistry," Academic Press, New York, N. Y., 1966, pp 132-215.

and treatment of the resulting ylide with benzaldehyde gives benzylidene compound **207** in 65% yield,⁶⁷ and ylide **208** with salicylaldehyde is converted to **209**.⁶⁵ Similarly, the phosphonium salt **210** upon treatment with base yields an ylide which with benzaldehyde or acetaldehyde is converted into pyran derivatives **211a** and **211b** in 40 and 42% yields, respectively.¹³⁶ The anion generated from dihydroacridinylphosphonate **64** is converted into a series of benzylidene dihydroacridines **212a-c** by treatment with the appropriate aldehyde in yields of 40-




70%.³³ A useful synthesis of pyrrolinones is presumed to involve a heterocyclic phosphonate intermediate, derived from **213**, which undergoes elimination of diethyl phosphate to give good yields of **214**.¹³⁷




2. Other C-P Bond Cleavages


Carbon-phosphorus bonds in heterocyclic systems seem to be much more susceptible to cleavage upon treatment with base than C-P bonds in aliphatic or most aromatic derivatives. Thus, triphenylphosphine oxide can withstand vigorous base treatment without much C-P cleavage, although certain aminophenylphosphonic acids undergo C-P bond cleavage with nucleophiles.¹³⁸ The α -hydroxyphosphonic acid esters are particularly susceptible to C-P bond cleavage upon treatment with aqueous base. Thus, the addition reaction of phosphorus anions to carbonyl groups (see section II.C) can be easily reversed. It is thought that the base removes a proton and the resulting anion expels a phosphonate anion (eq 1). It should be noted that the α -hydroxy acids are quite stable to base. Tri-2-thienylphosphine oxide (**215**) undergoes significant C-P bond cleavage upon heating with aqueous sodium hydroxide to a mixture of di-2-thienylphosphinic acid (**216**) and thiophene together with sodium meta-



phosphate.¹³⁹ The greater degree of cleavage in **215** compared with triphenylphosphine oxide is thought to be a reflection of the greater stability of the 2-thienyl anion in comparison with the phenyl anion.¹³⁹ Anions of the aryl group are thought to be involved in these cleavages. Tri-2-pyrrolylphosphine oxide (**217**)



is similarly cleaved in high yield (87%) by aqueous sodium hydroxide to yield pyrrole (**218a**).¹⁴ Diethyl pyrrolyl-2-phosphonate (**219a**) suffers C-P bond cleavage under these conditions to pyrrole (**218a**) (70%) and 2-ethylpyrrole (**218b**) (28%).¹⁴ Other bases, such as sodium hydride, ethylmagnesium bromide, or pyrrolylmagnesium bromide also serve to



a, R = H; b, R = Et; c, R = Me

convert the phosphonate **219a** into the cleavage products **218a** and **218b**.^{14,140} An attractive reaction pathway is as shown in which the base forms anion **220** which may or may not be intramolecularly alkylated before cleavage of the C-P bond. Nmr suggests the presence of **221**¹⁴⁰ (see section III.B). Supporting evidence for this pathway is provided by the observation that diethyl 1-methylpyrrolyl-2-phosphonate (**219c**) undergoes normal hydrolysis to its monoethyl ester with aqueous base.¹⁴ In common with several other 3-substituted indoles, 2-methyl-3-indolylphosphonic dichloride (**222**) is cleaved of its 3 substituent with aqueous base to yield, in this case, 2-methylindole (**224**). It is thought that indolenine intermediates, such as **223**, are involved.¹³

Triazine derivative **225** is particularly susceptible to hydrolytic cleavage so that attempted recrystallization from aqueous ethanol in air results in the formation of cyanuric acid (**226**) (93%) and diphenylphosphinic acid (**227**) (80%).²⁴ It is not


(136) S. V. Krivun, *Dokl. Akad. Nauk SSSR*, **182**, 347 (1968); *Chem. Abstr.*, **70**, 29009 (1969).

(137) G. Stork and R. Matthews, *Chem. Commun.*, 445 (1970).


(138) L. D. Freedman and G. O. Doak, *Chem. Rev.*, **57**, 479 (1957).

(139) K. R. Martin and C. E. Griffin, *J. Heterocycl. Chem.*, **3**, 92 (1966).


(140) C. E. Griffin, private communication.



clear whether air is essential for this cleavage. There are no reports regarding the hydrolytic stability of other phosphorus-substituted triazines.^{140a} 3,3-Diphenyl-5-triphenylphosphonia- Δ^1 -pyrazoline bromide (**228**) yields 5,5-diphenyl-3*H*-pyrazo-




line (**229**) in 87% as expected when treated with 10% sodium hydroxide.⁶⁷ This same phosphonium salt **228** undergoes ring





opening with extrusion of nitrogen upon heating to yield **230**, whereas the related pyrazoline **231** yields pyrazolium hydro-

bromide (**232**) quantitatively upon heating.⁶⁷ Thermal C-P bond cleavage in phosphonic acids has been observed in some cases,¹³⁸ so that the thermal conversion of 2,5-diphenylfuryl-3-phosphonic acid (**233**) to 2,5-diphenylfuran (**234**) is not exceptional.¹⁰⁹




B. REARRANGEMENTS

The rearrangement of epoxides to carbonyl compounds, usually induced by Lewis acids, is a well-established reaction.¹⁴¹ 1,2-Epoxyalkylphosphonates have been found to readily undergo rearrangement, with migration of the phosphonyl group, to β -carbonyl-containing phosphonates.^{92,99,102,142} Although in some cases an efficient thermal rearrangement has been observed, the use of boron trifluoride at room temperature is much more generally useful.^{92,99} In fact, the rearrangement products themselves can be thermally labile undergoing dephosphonation.⁹⁹ From the results of the rearrangement studies summarized in Table X, the following order of migratory aptitudes has been established: Ph > PO(OR)₂ > H > alkyl. It has been found that there are limitations in this rearrangement; for example, the less substituted members can undergo ring opening without rearrangement. Thus, diethyl 1,2-epoxyethylphosphonate (**235a**) upon treatment with boron trifluoride etherate yields hydroxy ether **236**,¹⁰² and phosphonoyl dimethyl ester (**235b**) gave no characterizable product with this reagent.⁸⁰ It is interesting to note, however, that mild base treatment of bromohydrin **237** induces a rearrangement with apparent phosphorus migration yielding alde-



hyde **238** (90%).⁸⁰ In the synthesis of indolyl-2-phosphonates **243a** and **243b**, it is postulated that the initially formed prod-




(140a) Recently it has been reported [J. P. Moreau and L. H. Chance, *J. Chem. Eng. Data*, **15**, 581 (1970)] that ammonia readily displaces one or two phosphonate groups from trisubstituted triazines such as 17.

(141) Reference 79, pp 230-261.

(142) B. A. Arbusov, N. A. Polezhaeva, and V. S. Vinogradova, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1146 (1967); *Chem. Abstr.*, **68**, 13092 (1968).

Table X
Rearrangement of 1,2-Epoxyethylphosphonates




Epoxide 239	Conditions	Product	Yield, %	Ref
R ₁ = Et; R ₂ = Ph; R ₃ = Me; R ₄ = H	Δ, 170°	240	86	99
R ₁ = Me; R ₂ = R ₃ = Me; R ₄ = H	Δ, 170°	240	...	99
R ₁ = Me or Et; R ₂ = R ₃ = Ph; R ₄ = H	Δ, 170°	240	...	99
R ₁ = Me or Et; R ₂ + R ₃ = -(CH ₂) ₅ -; R ₄ = G	BF ₃ , rt ^b	240	100	92, 99
R ₁ = Et; R ₂ = Ph; R ₃ = R ₄ = H	BF ₃ , rt	240	100	99
R ₁ = Et; R ₂ = H; R ₃ = Ph; R ₄ = Me	BF ₃ , rt	240	100	92
R ₁ = Et; R ₂ = H; R ₃ = Ph; R ₄ = <i>t</i> -Bu	BF ₃ , rt	240	33	92
R ₁ = Me; R ₂ = Me; R ₃ + R ₄ = -(CH ₂) ₄ -	BF ₃ , rt	240	100	92
R ₁ = Me; R ₂ = H; R ₃ = Ph; R ₄ = <i>p</i> -MeC ₆ H ₄	BF ₃ , rt	241	100	92
R ₁ = Me; R ₂ = H; R ₃ = R ₄ = Ph ^a	BF ₃ , rt	240, 11% 241	66	92
R ₁ = Et; R ₂ = R ₃ = H; R ₄ = Me	ZnCl ₂ , 160-170°	240	...	142
R ₁ = Et; R ₂ = H; R ₃ = R ₄ = Me	ZnCl ₂ , 160-170°	240	...	142
R ₁ = Et; R ₂ = H; R ₃ + R ₄ = -(CH ₂) ₃ -	ZnCl ₂ , 160-170°	240	...	142

^a This result is for (*E*)-stilbene oxide. The *Z* isomer gives 249 (34%) and HCOC(Ph)₂PO(OMe)₂ (66%) under similar conditions. The mode of formation of this latter product remains to be clarified. ^b rt = room temperature.

ucts are the 3-phosphonates 242a and 242b, which rearrange under the reaction conditions.¹¹¹ Although this rearrangement appears to be established for the indolylmethylphosphonate 242b, conclusive characterization is lacking in the case of the lower homolog where the product of the synthesis could be 242a and not 243a as claimed.¹¹¹

Diethyl 1-methyl-9,10-dihydroacridinyl-9-phosphonate (64) undergoes an isomerization reaction upon treatment with methyl- (or phenyl-) magnesium chloride yielding the 9-ethyl compound 245.³³ The Grignard reagent functions as a base to form anion 244 which undergoes intramolecular alkylation to yield 245. These results lend strong support for the postulated mechanism of the dephosphonation of diethyl pyrrolyl-2-phosphonate (219a) with base resulting in the formation of 2-ethylpyrrole^{14,140} (see section III.A.2).



C. OTHER REACTIONS

Much of the chemistry of heterocyclic phosphorus compounds is unexceptional and follows the behavior of the parent heterocycle or for manipulations on the phosphorus group, the behavior of that group in simple systems. Thus, for example, diethyl 1,2-epoxyethylphosphonate undergoes ring opening of

the epoxide upon treatment with nucleophiles,¹⁰² and the phosphonate ester, diethyl pyridyl-2-phosphonate, is readily hydrolyzed to the corresponding acid with aqueous acid.³⁸ Although this acid hydrolysis could not be applied to epoxyphosphonates without affecting the epoxide function, gentle methods are available for the conversion of phosphonate esters to phosphonic acids such as reaction with chlorotrimethylsilane followed by water.¹⁴³ This procedure has not been widely exploited, but its potential is demonstrated in the conversion of diethyl *cis*-1,2-epoxypropylphosphonate to the acid, phosphonmycin, leaving the epoxide ring intact.⁸⁰

IV. Properties

A. SPECTRA

Spectral measurements, particularly ultraviolet and nuclear magnetic resonance spectra, provide a means for determining the extent of π bonding involving the overlap of a phosphorus d orbital with a p orbital of an adjacent carbon of an aryl or vinyl group. Since certain aromatic heterocyclic systems are more effective electron donors than unsubstituted phenyl rings, heteroaryl phosphorus derivatives are particularly important systems in the study of $d\pi-p\pi$ bonding in organophosphorus compounds. This bonding is expressed in terms of canonical structures, such as B, and is reflected in bathochromic shifts in




Table XI
Ultraviolet Spectra of Heterocyclic Phosphorus Derivatives

Compound	λ_{max}, nm	$\epsilon_{max} \times 10^3$	$\Delta\lambda, nm$	Ref
Pyrrole	208	7.3	...	3
Tri-2-pyrrolylphosphine oxide	237.5	11.6	39.5	3
Diethyl pyrrolyl-2-phosphonate	211	6.5	3	14
1-Methylpyrrole	213	6.7	...	3
Tri-2-(1-methylpyrrolyl)phosphine oxide	243	12.9	30	3
Diethyl 1-methylpyrrolyl-2-phosphonate	218	6.4	5	14
Furan	205	6.4	...	3
Tri-2-furylphosphine oxide	238	33.6	33	3
	245	38.9	40	4
Tri-2-furylphosphine	243	21.8	38	4
Tri-2-furylphosphine sulfide	241	30.2	36	4
Tri-2-furylphosphine selenide	239	24.0	34	4
2,5-Diphenylfuran	324	29.2	...	109
2,5-Diphenylfuryl-3-phosphonic acid	318	36.4	-6	109
Thiophene	231	7.1	...	3
Tri-2-thienylphosphine oxide	238	33.0	7	3
Pyridine	251, 270	2.0, 0.45	...	144
Diethyl pyridyl-2-phosphonate	259, 267	2.79, 1.97	8, -4	38
2,6-Dimethylpyridine	266	4.5	...	145
Diethyl 2,6-dimethylpyridyl-4-phosphonate	279	3.2	13	
Acridine	359	10	...	146
Diethyl acridyl-9-phosphonate	369	11.48	10	33
2-Phenylindole	309	24.6	...	147
Diethyl 2-phenylindolyl-3-phosphonate	293	17.8	-16	129

the ultraviolet spectrum and, in the nmr spectra, by deshielding of certain ring protons in the heterocyclic ring. These measurements do not necessarily give identical results but are, in fact, complementary, since uv spectra are excited-state measurements and nmr spectra are ground-state measurements.


1. Ultraviolet Adsorption Spectra

Table XI summarizes the uv spectral data and shows the shift from the parent heterocyclic system ($\Delta\lambda$).¹⁴⁴⁻¹⁴⁷ Several trends are apparent from the data: those heterocyclic rings which are strongest electron donors show largest interaction and phosphine oxides conjugate more effectively than phosphonates. The 2-pyrrolyl- and 2-furylphosphine oxides show appreciable shifts from the parent heterocycle and a significant amount of $d\pi-p\pi$ bonding, but this is smaller than the shift observed for the formyl derivatives. In the case of tri-2-furylphosphine oxide there is a discrepancy in position of the maximum adsorption reported by two groups of workers.^{3,4} Thiophene and pyridine phosphorus derivatives show virtually no $d\pi-p\pi$ interaction from the measurements of the uv spectra. It should be noted, however, there is a much larger effect from a phosphonate in the 4 position than in the 2 position of pyridine. Thus, relative to the appropriate pyridine, diethyl pyridyl-2-phosphonate has $\Delta\lambda = 4$ nm and diethyl 2,6-dimethylpyridyl-4-phosphonate has $\Delta\lambda = 13$ nm.³⁸ The uv spectrum of the unknown tri-4-pyridylphosphine oxide *vs.* that of the 2-pyridyl isomer would be informative in determining the relative extent of $d\pi-p\pi$ bonding in 2- and 4-pyridyl systems. It should be

noted that 4-trimethylsilylpyridine is less basic than its 2 isomer, suggesting that $d\pi-p\pi$ interaction is stronger for 4-substituted pyridines.¹⁴⁸

2. Nuclear Magnetic Resonance Spectra

Detailed analysis has been made of the nmr spectra of a number of heteroarylphosphines and phosphine oxides and sulfides. In addition to chemical shift data, $^1H-^{31}P$ spin coupling data have been obtained for coupling through three, four, and five bonds. The chemical shift data, which for selected compounds are summarized in Table XII, have been used to determine the presence of $d\pi-p\pi$ bonding in the heteroaryl phosphorus compounds. For the five-membered heterocycles substantial deshielding of H(5) may be a reflection of $d\pi-p\pi$ interactions represented by the contribution of canonical form C. The interaction is much greater in tetracoordinate phosphorus



derivatives as can be seen by a comparison of the chemical shifts of tri-2-furylphosphine and its oxide (Table XII). The magnitude of the $^1H-^{31}P$ coupling strongly supports this conclusion in both five- and six-membered heteroaryl phosphorus

(144) A. I. Scott, "Interpretation of Ultraviolet Spectra of Natural Products," Pergamon Press, Oxford, 1964, p 179.

(145) H. C. Brown and X. R. Mihn, *J. Amer. Chem. Soc.*, **77**, 1723 (1955).

(146) A. Albert, "The Acridines," 2nd ed, Edward Arnold, London, 1966, p 176.

(147) M. J. Kamlet and J. C. Dacons, *J. Org. Chem.*, **26**, 220 (1961).

(148) D. G. Anderson, J. R. Chipperfield, and D. E. Webster, *J. Organometal. Chem.*, **12**, 323 (1968).

Table XII
Nmr Spectra for Heteroaryl Phosphine Derivatives and Model Compounds

	Solvent	Chemical shift, δ			Ref	
		H(3)	H(4)	H(5)		
Furan	CDCl ₃	6.37	6.37	7.42	149	
2-Formylfuran	CDCl ₃	7.28	6.63	7.72	149	
Tri-2-furylphosphine	CDCl ₃	6.63	6.26	7.45	151	
Tri-2-furylphosphine oxide	CDCl ₃	7.14	6.53	7.71	149, 151	
Tri-2-furylphosphine sulfide	CDCl ₃	7.15	6.50	7.69	151	
Tri-2-furylphosphine selenide	CDCl ₃	7.19	6.50	7.71	151	
Diethyl 2-furylphosphonate	(CH ₃) ₂ CO	7.10	6.62	7.88	149	
2-Formylpyrrole	CDCl ₃	7.17	6.30	6.98	149	
Tri-2-pyrrolylphosphine oxide	CDCl ₃	6.33	6.33	6.82	149	
Thiophene	CDCl ₃	7.20	7.20	7.30	149	
2-Formylthiophene	CDCl ₃	7.78	7.22	7.78	149	
Tri-2-thienylphosphine oxide	CDCl ₃	7.58	7.17	7.74	149, 151	
Diethyl 2-thienylphosphonate	(CH ₃) ₂ CO	7.64	7.27	7.94	149	
		H(3)	H(4)	H(5)	H(6)	
2-Formylpyridine	(CH ₃) ₂ SO	8.36	8.17	7.88	9.03	150
Tri-2-pyridylphosphine oxide	CDCl ₃	8.19	7.78	7.35	8.75	150, 152
Diethyl 2-pyridylphosphonate	CCl ₄	8.10	8.02	7.65	8.88	38

compounds.¹⁴⁹⁻¹⁵² For example, in tri-2-thienylphosphine oxide the coupling between phosphorus and hydrogens at C₃, C₄, and C₅ is 7.98, 2.00, and 4.48 Hz, while in tri-2-thienylphosphine these values are 6.17, 1.33, and 0.19 Hz, respectively.¹⁵¹ The increased deshielding of H(5) with increasing solvent polarity, Table XIII, argues in favor of an important

constants range from 1.18 to 3.10 Hz [for ³¹P with H(5)] for tri-2-pyridylphosphine to tri-2-pyridylphosphine selenide.

The nmr spectra of a series of azepine phosphonates **246** have been studied and, in fact, formed part of the structure proof.¹⁵⁴ This analysis also disclosed a five-bond ¹H-³¹P spin coupling ($J = 2$ Hz).


Table XIII

Solvent Effect on Chemical Shift of Tri-2-thienylphosphine¹⁵¹

Solvent	Chemical shift, δ		
	H(3)	H(4)	H(5)
Carbon tetrachloride	7.26	6.96	7.41
Deuteriochloroform	7.32	7.02	7.50
Deuterioacetone	7.35	7.09	7.70
Deuterioacetonitrile	7.33	7.06	7.62
Deuteriodimethyl sulfoxide	7.35	7.10	7.79

contribution from resonance hybrid C and hence to the existence of $d\pi-p\pi$ interaction. The general conclusions from Table XII are that in furan and thiophene phosphorus derivatives $d\pi-p\pi$ interaction is present, whereas in pyrrole and pyridine derivatives there is little or no evidence for interaction.

Studies on other derivatives not summarized in Table XII include tri-2-thienylphosphine sulfide and selenide,¹⁵¹ tri-2-(5-methylfuryl)phosphine and its sulfide,¹⁵³ tri-3-thienylphosphine and its oxide, sulfide, and selenide,¹⁵¹ and tri-2-pyridylphosphine and its sulfide and selenide.^{150,152} Analysis of the spectrum of the pyridylphosphines has given the first values of ¹H-³¹P coupling through five bonds.^{150,152} These coupling



- 246a**, R = Et; R₁ = Me; R₂ = H
b, R = Et; R₁ = Et; R₂ = H
c, R = Me; R₁ = H; R₂ = Me
d, R = Me; R₁ = Me; R₂ = H
e, R = Et; R₁ = R₂ = H

B. BIOLOGICAL ACTIVITY

Although the stated purpose of many of the syntheses of heterocyclic phosphorus compounds, particularly phosphonic acids, was to determine biological activity, very few positive results are reported. From this class of compounds, phosphonmycin appears to stand alone in its biological activity.⁷⁷ It is reported to be a bactericidal antibiotic effective against a wide range of organisms comparable to tetracycline or chloramphenicol in its activity. This antibiotic, which has low toxicity, functions by inhibiting bacterial cell-wall synthesis.⁷⁷ Among compounds reported to have weak or no biological activity are quinolinephosphonic acids,¹⁹ 3-pyridylphosphonic acid,⁸⁶ thiazolephosphonate ethyl ester,¹²¹ and pyrimidinephosphonate esters.²² It would seem that many of the phosphonates tested only as esters should be reexamined as the corresponding acids which may well be more readily assimilated by organisms.

(149) R. H. Kemp, W. A. Thomas, M. Gordon, and C. E. Griffin, *J. Chem. Soc. B*, 527 (1969).

(150) G. E. Griffin and W. A. Thomas, *ibid.*, 477 (1970).

(151) H. J. Jakobsen and J. A. Nielsen, *J. Mol. Spectrosc.*, **33**, 474 (1970).

(152) H. J. Jakobsen, *ibid.*, **34**, 245 (1970).

(153) H. J. Jakobsen and M. Begrup, *ibid.*, **35**, 158 (1970).

(154) J. I. G. Cadogan and R. K. Mackie, *J. Chem. Soc. C*, 2819 (1969).