

SUBJECT INDEX FOR VOLUME 72, 1972

Biosynthesis, of monoterpenes, 115
 Cancer, metals, and ligands, 203
 Carbenes, complexes with transition metals, 545
 Catalysis, general acid–base, of complex reactions in water, 705
 (CH)_n compounds, rearrangements and interconversions, 181
 Coordination chemistry, of pyrazole-derived ligands, 497
 Critical constants of inorganic substances, 71
 Cycloaddition reactions, 157
 Cyclohexanone, vibrational analysis of, 439
 Dielectric relaxation, use of in the dipole correlation function in, 55
 Dihydropyridines, 1
 Dioxolan-2-ylum cations, 1,3-, 357
 Dipole correlation function, in dielectric relaxation, use of, 55
 Electron-transfer reactions, aqueous V(IV) as reductant *vs.* Fe(II), 215
 General acid–base catalysis, of complex reactions in water, 705
 Group frequencies, in vibrational analysis of cyclohexanone, 439
 Heterocyclic cations, related to 1,3-dioxolan-2-ylum cations, 357
 Hybridization, by the maximum overlap method, 43
 Hydride complexes, of transition metals, 231
 Hydrogen atoms, reactions of in aqueous solutions, 533
 Hydrogen bond, theory of, 283
 Interconversions, of compounds of formula (CH)_n, 181
 Iron(II), as reductant *vs.* V(IV), in aqueous electron-transfer reactions, 215
 Isocyanate chemistry, 457
 Isocyanides, polymerization of, 101
 Isotope effects, kinetic, as measure of structure–reactivity relationships, 511

Chemical Reviews Indexes.—Each annual volume has its own author and subject indexes in the December issue. Cumulative indexes for Volumes 1–60 and 61–70 were published in the December issues of 1960 and 1970, respectively. The next cumulative

Kinetic isotope effects, as measure of structure–reactivity relationship, 511
 Lead dioxide electrode, 679
 Ligands, metals, and cancer, 203
 Maximum overlap method, hybridization by, 43
 Mercury, surface properties of, 575
 Metals, ligands, and cancer, 203
 Monoterpenes, biosynthesis of, 115
 Neighboring group interactions, in ortho-substituted nitrobenzenes, 627
 Nitrobenzene derivatives, ortho-substituted, synthetic methods involving neighboring group interaction, 627
 Normal modes, in vibrational analysis of cyclohexanone, 439
 Phosphazene chemistry, 315
 Phosphonitrilic compounds, 315
 Polymerization, of isocyanides, 101
 Pyrazole-derived ligands, coordination chemistry of, 497
 Pyridines, dihydro-, 1
 Rearrangement, of compounds of formula (CH)_n, 181
 Structure–reactivity relationships, kinetic isotope effects as measure of, 511
 Surface properties, of mercury, 575
 Terpenes, mono-, biosynthesis of, 115
 Thujane derivatives, 305
 Transition metals, complexes with hydrides, 231; complexes with carbenes, 545
 Vanadium(IV), as reductant *vs.* Fe(II), in aqueous electron-transfer reactions, 215
 Vibrational analysis, of cyclohexanone, 439

index will appear in the December 1980 issue. Cumulative indexes may be purchased from the Special Issues Sales Department, American Chemical Society, 1155 Sixteenth St., N.W., Washington, D. C. 20036.

ERRATA

Volume 71, 1971

Sites and Thermodynamic Quantities Associated with Proton and Metal Ion Interaction with Ribonucleic Acid, Deoxyribonucleic Acid, and Their Constituent Bases, Nucleosides, and Nucleotides. By R. M. Izatt, J. J. Christensen, and J. H. Rytting

Page 449: The work in references 130 and 131 *disproves* rather than *supports* binding of one copper(II) ion simultaneously to the N(3) and N(9) positions of a purine ring. Later work by these authors supports this conclusion: *Chem. Commun.*, 1573 (1970); *Nature*, **229**, 191 (1971); *Biochem. Biophys. Res. Commun.*, **42**, 63 (1971). The copper ions are held in pairs by a purine bridge. Within each Cu–adenine–Cu unit one copper binds to N(3) and the other to N(9).

Page 463: Table VI. Reaction 5'-AMP (L), Mⁿ⁺ + H₂L⁻ + MH₂Lⁿ⁻¹ should read 5'-AMP (L), Mⁿ⁺ + HL²⁻ = MHLⁿ⁻² (these complexes are formed with the protonated phosphate group).

Page 463: And first entry on p 464. The log *K* values from ref 201 under the heading 5'-AMP–M²⁺ were measured in the presence of the metal ion concentrations listed in ref 201 on p 1712. Therefore the given log *K* values are only “apparent” constants.

The charges on several of the complexes are known. Changes should be made on the indicated pages.

Page 473: GTP–Cu²⁺ (L), H⁺ + L²⁻ = HL^{+?} (ionic charge of GTP–Cu²⁺ not given) should read Cu(GTP)³⁺ + H⁺ = Cu(HGTP)²⁻.

Page 474: ITP–Cu²⁺ (L), H⁺ + L²⁻ = HL^{+?} (Base) (ionic charge of ITP–Cu²⁺ not given) should read Cu(ITP)³⁻ + H⁺ = Cu(HITP)²⁻.

Page 475: TTP–Cu²⁺ (L), H⁺ + L²⁻ = HL^{+?} (ionic charge of TTP–Cu²⁺ not given) should read Cu(TTP)³⁻ + H⁺ = Cu(HTTP)²⁻.

Page 477: UTP–Cu²⁺ (L), H⁺ + L²⁻ = HL^{+?} (ionic charge of UTP–Cu²⁺ not given) should read Cu(UTP)³⁻ + H⁺ = Cu(HUTP)²⁻.