

Chemical Reviews

Volume 76, Number 3 June 1976

Chromium, Molybdenum, and Tungsten: Thermodynamic Properties, Chemical Equilibria, and Standard Potentials

INGEMAR DELLIEN,¹ F. M. HALL,² and LOREN G. HEPLER*

*Department of Chemistry, University of Lethbridge,
Lethbridge, Alberta, Canada*

Received January 27, 1975 (Revised Manuscript Received June 3, 1975)

Contents

I. Introduction	283
II. Chromium	283
A. Descriptive Chemistry	283
B. Element and Oxides	284
C. Cr(VI): Compounds and Aqueous Ions	285
D. Cr(II) and Cr(III): Compounds and Aqueous Ions	287
E. Cr(IV) and Cr(V): Compounds and Aqueous Ions	292
F. Other Inorganic Compounds of Chromium	293
G. Organic Compounds and Complexes	294
III. Molybdenum and Tungsten	294
A. Descriptive Chemistry	294
B. Molybdenum, Molybdenum Oxides, and Molybdic Acid	295
C. $\text{MoO}_4^{2-}(\text{aq})$ and Molybdates	295
D. Halides and Oxyhalides of Molybdenum	298
E. Other Molybdenum Compounds	300
F. Tungsten, Tungsten Oxides, and Tungstic Acid	301
G. $\text{WO}_4^{2-}(\text{aq})$ and Tungstates	303
H. Halides and Oxyhalides of Tungsten	304
I. Other Tungsten Compounds	305
IV. Appendix	306
V. References and Notes	306

I. Introduction

In this review we are primarily concerned with the thermodynamic properties of compounds and aqueous ions of chromium, molybdenum, and tungsten. Thermodynamic data for all of these have long been known to be useful in electrochemistry, analytical chemistry, metallurgy, and many applications of chemical principles. Because many of these thermodynamic quantities are most useful when presented in the form of equilibrium constants or standard potentials, we have listed a large number of these quantities.

We have been critical in our tabulations of data and have recalculated many of the published results cited here. When data from several sources are not in good agreement, we have attempted to justify our choices. In general, we have been explicit about the sources of data and also our treatment of data so that interested readers can check the steps leading to our tabulated values of thermodynamic properties (also related equilibrium constants and standard potentials) and form their own opinions about reliability and accuracy.

In several cases we have combined experimental data from a cited source with an estimate of some thermodynamic

property. Numerical values of estimates are given in parentheses. We have used auxiliary thermodynamic data [such as ΔH_f° of $\text{Cl}^-(\text{aq})$] from the National Bureau of Standards Technical Notes 270-3,³ 270-4,⁴ and 270-6,⁵ which are the most comprehensive sources of such data available to us.

In spite of considerable discussion in recent years, general agreement is still lacking on "sign conventions" for potentials. Much of the confusion arises because "sign" can be either electrical or algebraic, as previously discussed.⁶ All potentials listed in this review are reduction potentials with algebraic signs and may also be taken to be electrode potentials with electrical signs relative to the potential of the standard hydrogen electrode.

The symbols E° and K are used for potentials and equilibrium constants that have been determined in very dilute solutions or in such ways that activity coefficients could be considered in treating the experimental data. Similarly, we use ΔG° , ΔH° , and ΔS° to indicate data that refer to the usual standard states.^{3-5,7} Other useful investigations have been carried out in solutions maintained at some constant ionic strength by means of some "inert" supporting electrolyte. In some cases it has been possible to make reasonable estimates of adjustments to the usual infinite dilution standard states. In other cases we adopt properties based on "uncorrected" data and call attention to the fact that the cited properties refer to solutions having some stated ionic strength.

Except where explicitly stated otherwise, all thermodynamic properties cited in this review refer to 298.15 K (25 °C).

II. Chromium

A. Descriptive Chemistry

In addition to the familiar +2, +3, and +6 oxidation states, compounds and ions containing chromium in the -2, -1, 0, +4, and +5 oxidation states have been reported. In this section we present brief qualitative descriptions of some compounds and ions characteristic of the various oxidation states and then turn to detailed discussion of thermodynamic properties, equilibrium constants, and standard potentials in following sections.

The best known compound containing chromium in the zero oxidation state is the hexacarbonyl, $\text{Cr}(\text{CO})_6$. Reaction of $\text{Cr}(\text{CO})_6$ with alkaline solutions yields compounds containing chromium in the -1 and -2 oxidation states. Represent-

tative compounds of this sort are $\text{Na}_2[\text{Cr}(\text{CO})_5]$ and $\text{Na}_2[\text{Cr}_2(\text{CO})_{10}]$. Although it is well established that these and similar compounds are easily oxidized, we have no relevant quantitative data to cite later in this review.

Reduction of $\text{K}_3\text{Cr}(\text{CN})_6$ in liquid NH_3 yields $\text{K}_6\text{Cr}(\text{CN})_6$, which may be described as containing chromium in the zero oxidation state.⁸

Nash, Campbell, and Block⁹ have studied the preparation of chromium by thermal decomposition of bis(benzene)chromium.

Electron spin resonance and optical measurements have shown¹⁰ that it is reasonable to describe $\text{K}_3[\text{Cr}(\text{CN})_5\text{NO}]\cdot\text{H}_2\text{O}$ as containing "effective" Cr^+ and NO^+ ions. Petrakis¹¹ has made lattice energy calculations that lead to estimates of enthalpies of formation of the hypothetical Cr^+ halides. These calculated ΔH_f° values show that the hypothetical solid CrX compounds are quite unstable with respect to disproportionation.

Reasonably common compounds containing chromium in the +2 oxidation state are the halides, the sulfate, and the acetate. Such hydrated salts as $\text{CrCl}_2\cdot 4\text{H}_2\text{O}$, $\text{CrSO}_4\cdot 7\text{H}_2\text{O}$, and $[\text{Cr}(\text{Ac})_2]_2\cdot 2\text{H}_2\text{O}$ have been obtained from aqueous solutions. Because of its low solubility in aqueous solution, dimeric chromous acetate is one of the most easily prepared compounds of $\text{Cr}(\text{II})$. This interesting red compound has an unusually short $\text{Cr}-\text{Cr}$ distance and is diamagnetic at room temperature, presumably because of considerable interaction between the metal ions. Deyrup¹² has described preparation of some air-stable $\text{Cr}(\text{II})$ compounds: Na_2CrF_4 and $\text{CrXOH}\cdot\text{HBO}_3$ ($\text{X} = \text{Cl}, \text{Br}, \text{or I}$). Beauchamp and Sullivan¹³ have developed procedures for large-scale preparation of CrCl_2 .

Under most conditions the +3 oxidation state is the most stable state for chromium. The oxide, halides, sulfate, and nitrate are all common compounds. Many octahedral complexes of $\text{Cr}(\text{III})$ are known in solution and in the solid state. One indication of the number and variety of such complexes is afforded by the work of Kyuno, Kamada, and Tanaka,¹⁴ who have applied a systematic synthetic procedure to preparation of a series of 25 ammine complexes of $\text{Cr}(\text{III})$. Because of the slow substitution reactions of many complexes of $\text{Cr}(\text{III})$, these species may persist for long times under conditions where they are thermodynamically unstable. In this review our attention is largely limited to "inorganic" complexes for which fairly complete thermodynamic data are available.

Swaddle et al.¹⁵ have recently discussed the aqueous chemistry of $\text{Cr}(\text{III})$ above 100 °C, with particular emphasis on various spinels, which have also been prepared by direct reaction of Cr_2O_3 with various other oxides.

A few solid and gaseous compounds of $\text{Cr}(\text{IV})$ and $\text{Cr}(\text{V})$ have been prepared.¹⁶ Some of these are the following: Ba_2CrO_4 , Cr_2O_5 , CrO_2 , CrF_4 , and oxyfluorides of $\text{Cr}(\text{V})$. Soluble species containing $\text{Cr}(\text{IV})$ and $\text{Cr}(\text{V})$ are unstable with respect to disproportionation or reaction with solvent under most conditions. Several aqueous species of $\text{Cr}(\text{IV})$ and $\text{Cr}(\text{V})$ have been postulated as intermediates in oxidation-reduction reactions, and it is possible to make reasonable estimates of standard potentials for half-reactions involving these species. Evidence for moderately stable $\text{Cr}(\text{V})$ species in sulfuric acid solutions¹⁷ and in strong base¹⁸ has been reported.

Peroxo complexes of $\text{Cr}(\text{III}), \text{IV}, \text{V}, \text{and VI}$ have been reviewed by Connor and Ebsworth¹⁹ and more recently by Cotton and Wilkinson.²⁰ We also call attention to a recent equilibrium investigation of peroxo species of $\text{Cr}(\text{V})$ and VI by Bartlett and Quane.²¹ Among the peroxo compounds and ions that have been identified or postulated are the following: K_3CrO_8 , $\text{CrO}(\text{O}_2)_2$ and related salts, $\text{CrO}_5\cdot\text{A}$ ($\text{A} = \text{ether}, \text{pyridine}, \text{etc.}$), $\text{Cr}_2(\text{O}_2)^{4+}$, and $\text{Cr}_3(\text{O}_2)_2^{5+}$.

Many solid compounds and aqueous ions containing $\text{Cr}(\text{VI})$ are known and have been investigated enough that we can

later quote reliable data. Several of these compounds and ions are useful oxidizing agents in acidic solution. The most common $\text{Cr}(\text{VI})$ compounds and aqueous ions are CrO_3 , $\text{K}_2\text{Cr}_2\text{O}_7$, K_2CrO_4 , $\text{Cr}_2\text{O}_7^{2-}(\text{aq})$, $\text{HCrO}_4^-(\text{aq})$, and $\text{CrO}_4^{2-}(\text{aq})$. There are several slightly soluble chromates that are of some importance, including Ag_2CrO_4 , PbCrO_4 , BaCrO_4 , etc.

B. Element and Oxides

The best general sources of thermodynamic data for chromium and its compounds are the National Bureau of Standards Technical Notes 270-4⁴ and 270-6,⁵ to which we hereafter refer as NBS 270-4 and NBS 270-6.

We follow NBS 270-4 in taking $\Delta H_f^\circ = 0$, $\Delta G_f^\circ = 0$, and $S^\circ = 5.68 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{Cr}(\text{c})$ at 298 K. Kelley and King²² have reviewed earlier C_p measurements that lead to this $S^\circ = 5.68 \text{ cal K}^{-1} \text{ mol}^{-1}$. Two more recent investigations^{23,24} have led to $S^\circ = 5.64$ and $5.72 \text{ cal K}^{-1} \text{ mol}^{-1}$. We also note that Bonilla and Garland²⁵ have made C_p measurements on $\text{Cr}(\text{c})$ at various pressures near the Néel transition.

We accept $S^\circ = 41.68 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{Cr}(\text{g})$ as listed in NBS 270-4, based on statistical thermodynamic calculations. Stull and Sinke²⁶ and more recently Storms²⁷ have reviewed results of high-temperature vapor pressure studies and have calculated $\Delta H_f^\circ = 95.0$ and $94.89 \pm 0.31 \text{ kcal mol}^{-1}$ for $\text{Cr}(\text{g})$. Dickson, Myers, and Saxer²⁸ have made measurements (not considered above^{26,27}) that lead to $\Delta H_f^\circ = 95.3 \text{ kcal mol}^{-1}$ for $\text{Cr}(\text{g})$ at 298 K. On the basis of all these results, we adopt $\Delta H_f^\circ = 95.0 \text{ kcal mol}^{-1}$ for $\text{Cr}(\text{g})$ and combine with the S° to obtain $\Delta G_f^\circ = 84.3 \text{ kcal mol}^{-1}$.

For the $\text{Cr}^{n+}(\text{g})$ ions we adopt the ΔH_f° values from NBS 270-4 after adjustment for the change in ΔH_f° for $\text{Cr}(\text{g})$ from the NBS 270-4 value ($94.8 \text{ kcal mol}^{-1}$) to our value ($95.0 \text{ kcal mol}^{-1}$).

For $\text{CrO}(\text{g})$ we adopt $\Delta H_f^\circ \simeq 53 \text{ kcal mol}^{-1}$ from Grimley, Burns, and Inghram²⁹ and $S^\circ = 53.0 \text{ cal K}^{-1} \text{ mol}^{-1}$ from Kelley and King.²² These values lead to $\Delta G_f^\circ = 46 \text{ kcal mol}^{-1}$ for $\text{CrO}(\text{g})$.

The NBS 270-4 lists ΔH_f° (at 0 K) = $-14 \text{ kcal mol}^{-1}$ for $\text{CrO}_2(\text{g})$. This value is consistent with the results of Grimley et al.²⁹ More recent investigations^{30,31} have led to less exothermic values. On the basis of all three investigations,²⁹⁻³¹ we adopt $\Delta H_f^\circ \simeq -3 \text{ kcal mol}^{-1}$ and combine with the $S^\circ = 62.0 \text{ cal K}^{-1} \text{ mol}^{-1}$ from Nagarajan³² to obtain ΔG_f° for this compound.

The NBS 270-4 lists $\Delta H_f^\circ = -143 \text{ kcal mol}^{-1}$ for $\text{CrO}_2(\text{c})$. Although the only calorimetric investigation we know of for $\text{CrO}_2(\text{c})$ (by Ariya et al.³³) has led to a less exothermic value, we tentatively accept the value from NBS 270-4 and combine with an estimated S° to obtain its ΔG_f° .

For $\text{Cr}_2\text{O}_3(\text{c})$ we have a well-established $S^\circ = 19.4 \text{ cal K}^{-1} \text{ mol}^{-1}$, based on work reviewed by Kelley and King,²² and as adopted in NBS 270-4. Unfortunately, the situation is not so clear for ΔH_f° and ΔG_f° for this important compound.

Combustion calorimetry by Mah^{34,35} gives $\Delta H_f^\circ = -272.6 \pm 0.6 \text{ kcal mol}^{-1}$ for $\text{Cr}_2\text{O}_3(\text{c})$. High-temperature equilibrium measurements of the reduction of $\text{Cr}_2\text{O}_3(\text{c})$ by $\text{H}_2(\text{g})$ have led Novokhatskii and Lenev³⁶ to report ΔH_f° (at 298 K) = $-272.8 \pm 0.8 \text{ kcal mol}^{-1}$ for $\text{Cr}_2\text{O}_3(\text{c})$. Our own second law calculation with their results leads to a slightly more negative value, while our third law calculation with the same results leads to a slightly less negative value, with both calculations making use of high-temperature data from Kelley.³⁷ All of these values are in agreement with $\Delta H_f^\circ = -272.4 \text{ kcal mol}^{-1}$ for $\text{Cr}_2\text{O}_3(\text{c})$ as listed in NBS 270-4.

But there is also evidence in favor of a significantly less negative ΔH_f° for $\text{Cr}_2\text{O}_3(\text{c})$. For example, high-temperature equilibrium results have led Ramsey, Caplan, and Burr³⁸ to report $\Delta H_f^\circ = -268.5 \text{ kcal mol}^{-1}$. High-temperature emf

measurements by Tretjakow and Schmalzried³⁹ have led to results that permit us to calculate (second law) $\Delta H_f^\circ = -261$ kcal mol⁻¹ and (third law) $\Delta H_f^\circ = -268$ kcal mol⁻¹ for Cr₂O₃(c). More recent high-temperature emf measurements by Mazandarany and Pehike⁴⁰ and also high-temperature equilibrium measurements by Jeannin, Mannerskantz, and Richardson⁴¹ have yielded results that we have used for calculation of ΔH_f° (at 298 K) = -270.0 kcal mol⁻¹ for Cr₂O₃(c), again making use of auxiliary data from Kelley.³⁷

It is obvious from the discussion in the two paragraphs above that there are substantial and unexplained discrepancies between the results of high-quality investigations. Kulkarni and Worrell⁴² have selected $\Delta H_f^\circ = -270.0$ kcal mol⁻¹ for Cr₂O₃(c), partly on the basis of the work of Richardson et al.⁴¹ and partly to achieve consistency with some results for chromium carbides that we discuss later. But it should also be pointed out that consistency with some other results for chromium carbides is best achieved by way of $\Delta H_f^\circ = -272.6$ kcal mol⁻¹ for Cr₂O₃(c) as found by Mah^{34,35} or $\Delta H_f^\circ = -272.4$ kcal mol⁻¹ as listed in NBS 270-4.

In the absence of further definitive measurements involving Cr₂O₃(c), we must make a somewhat arbitrary choice of a best ΔH_f° and then make as many other values as possible be consistent with this chosen value. Partly for convenience in relating our Tables to NBS 270-4 and 270-6 and partly because of our respect for the work of Mah^{34,35} and prejudice in favor of calorimetry, we adopt $\Delta H_f^\circ = -272.4$ kcal mol⁻¹ for Cr₂O₃(c) as listed in NBS 270-4. Therefore we also adopt $\Delta G_f^\circ = -252.9$ kcal mol⁻¹ as listed in NBS 270-4.

Combination of enthalpies of solution⁴³ of CrO₃(c) and (NH₄)₂Cr₂O₇(c) with the enthalpy of decomposition⁴⁴ of (NH₄)₂Cr₂O₇(c) and the ΔH_f° of Cr₂O₃(c) adopted above leads to $\Delta H_f^\circ = -140.9$ kcal mol⁻¹ for CrO₃(c), which is the value listed in NBS 270-4 and is also the value we adopt for this compound. We note, however, that there are other paths to this ΔH_f° . Bichowsky and Rossini⁴⁵ have quoted (from Roth and Becker) a ΔH for decomposition of CrO₃(c) to Cr₂O₃(c) and O₂(g) that leads us to $\Delta H_f^\circ = -139.7$ kcal mol⁻¹ for CrO₃(c), based on our adopted ΔH_f° for Cr₂O₃(c). More recent measurements by Richelmi and Laffitte⁴⁶ of the enthalpy of decomposition of CrO₃(c) lead to $\Delta H_f^\circ = -137.0$ kcal mol⁻¹ for CrO₃(c). Because of the disagreements between these values, it must be concluded that at least one of the three kinds of measurements [enthalpies of solution, enthalpy of decomposition of CrO₃, enthalpy of decomposition of (NH₄)₂Cr₂O₇] is in error. We combine an estimated S° with our adopted ΔH_f° to obtain the ΔG_f° of CrO₃(c).

For CrO₃(g) we adopt $\Delta H_f^\circ \approx -65$ kcal mol⁻¹ from the results of Grimley et al.²⁹ and the more recent results of Farber and Srivastava.³¹ We combine this value, which is considerably less negative than the value listed in NBS 270-4, with $S^\circ = 63.6$ cal K⁻¹ mol⁻¹ from Nagarajan³² to obtain $\Delta G_f^\circ \approx -60$ kcal mol⁻¹ for this compound.

McDonald and Margrave⁴⁷ have investigated vaporization of CrO₃ to form various (CrO₃)_n(g) species.

The investigation by Richelmi and Laffitte⁴⁶ of the decomposition of CrO₃(c) to Cr₂O₃(c) indicates that the reaction proceeds via Cr₅O₁₂(c) and Cr₈O₂₁(c). Their results lead to ΔH_f° values for these compounds, which we list as $\Delta H_f^\circ = -140.3$ kcal mol⁻¹ for Cr₅O₁₂(c) and $\Delta H_f^\circ = -139.1$ kcal mol⁻¹ for Cr₈O₂₁(c). Because the ΔH_f° of CrO₃(c) from these workers⁴⁶ differs from the value we have adopted, the ΔH_f° values quoted above for the intermediate oxides may not bear the right relationship to our ΔH_f° values for CrO₃(c) and Cr₂O₃(c).


C. Cr(VI): Compounds and Aqueous Ions

Bomb calorimetric measurements by Neugebauer and

Margrave⁴⁴ have led to ΔH° of decomposition of (NH₄)₂Cr₂O₇(c) to Cr₂O₃(c). Combination of this result with our selected ΔH_f° of Cr₂O₃(c) leads to $\Delta H_f^\circ = -431.8$ kcal mol⁻¹ for (NH₄)₂Cr₂O₇(c), which is the same as the value listed in NBS 270-4 for this compound. As noted in the preceding section, we have used this result with enthalpies of solution⁴³ in establishing the ΔH_f° of CrO₃(c).

The most direct route to the ΔH_f° of CrO₄²⁻(aq) is by way of the calorimetrically measured⁴³ enthalpy of solution of CrO₃(c) in excess OH⁻(aq). Combination of the resulting ΔH° of solution with our adopted ΔH_f° of CrO₃(c) leads to $\Delta H_f^\circ = -210.60$ kcal mol⁻¹ for CrO₄²⁻(aq), which is the value we adopt and is also the value listed in NBS 270-4.


The entropy of CrO₄²⁻(aq) can be obtained by way of Ag₂CrO₄(c) as follows. Smith, Pitzer, and Latimer⁴⁸ have reported $S^\circ = 51.82$ cal K⁻¹ mol⁻¹ for Ag₂CrO₄(c), based on their C_p measurements. Subsequent recalculation²² has led to $S^\circ = 52.0$ cal K⁻¹ mol⁻¹, which is the value listed in NBS 270-4 and adopted here. Smith, Pitzer, and Latimer⁴⁸ have also made calorimetric measurements leading to $\Delta H^\circ = 14.45$ kcal mol⁻¹ for



Subsequent emf measurements and the derived $d \ln K_{sp}/dT$ from Pan⁴⁹ lead to $\Delta H^\circ = 14.49$ kcal mol⁻¹ for this reaction, in good agreement with the calorimetric result.⁴⁸ Smith, Pitzer, and Latimer⁴⁸ chose $K_{sp} = 1.1 \times 10^{-12}$ from the work of Cann and Mueller.⁵⁰ We combine the ΔG° of solution from this K_{sp} with their⁴⁸ enthalpy of solution and entropies of Ag⁺(aq) and Ag₂CrO₄(c) to obtain $S^\circ = 11.1$ cal K⁻¹ mol⁻¹ for CrO₄²⁻(aq). There are, however, two other K_{sp} values worth consideration in connection with this calculation. Pan's measurements⁴⁹ led to $K_{sp} = 1.3 \times 10^{-12}$ at 25 °C, while conductance measurements by Howard and Nancollas⁵¹ have led to $K_{sp} = 2.5 \times 10^{-12}$ for Ag₂CrO₄(c). Use of this latter K_{sp} in the calculation described above leads to $S^\circ = 12.6$ cal K⁻¹ mol⁻¹ for CrO₄²⁻(aq), while Pan's results lead to a value intermediate between 11.1 and 12.6 cal K⁻¹ mol⁻¹ for this ion. It is also possible to calculate S° for CrO₄²⁻(aq) from the solubility, enthalpy of solution, activity coefficient in saturated solution, and the third law entropy of K₂CrO₄(c), but the result is not accurate enough to help us choose between the entropies cited above. In the absence of another reliable path or new data for silver chromate, the best we can do is adopt $S^\circ = 12$ cal K⁻¹ mol⁻¹ for CrO₄²⁻(aq). The NBS 270-4 lists $S^\circ = 12.00$ cal K⁻¹ mol⁻¹ for this ion, but we are unable to justify the last two figures in this value.

Combination of our selected S° and ΔH_f° values leads to $\Delta G_f^\circ = -173.9_6$ kcal mol⁻¹ for CrO₄²⁻(aq), in agreement with the NBS 270-4.


The properties of HCrO₄⁻(aq) can now be derived from the thermodynamics of the reaction



The equilibrium results of several sets of investigators⁵²⁻⁵⁷ are in agreement with $K = 3.3 \times 10^{-7}$ for this reaction at 298 K. We therefore adopt the corresponding $\Delta G_f^\circ = -182.8$ kcal mol⁻¹ for HCrO₄⁻(aq), which is also the value listed in NBS 270-4. We also adopt $\Delta H_f^\circ = -209.9$ kcal mol⁻¹ for this ion as derived from a calorimetric⁴³ ΔH° for reaction 2 and as listed in NBS 270-4. This ΔH_f° is in good agreement with the $d \ln K/dT$ results of Linge and Jones.⁵² Combination of the ΔG_f° and ΔH_f° values above leads to $S^\circ = 44$ cal K⁻¹ mol⁻¹ for HCrO₄⁻(aq).

We also note that there have been several investigations⁵⁶⁻⁵⁸ of reaction 2 in 3 M ClO₄⁻ medium, with the best results probably being those of Arnek and Johansson.⁵⁸

For the reaction




we choose $K = 33.9$ from the results of several investigators⁵⁹⁻⁶¹ and the corresponding $\Delta G_f^\circ = -311.0 \text{ kcal mol}^{-1}$ for $\text{Cr}_2\text{O}_7^{2-}(\text{aq})$, with this same ΔG_f° being listed in NBS 270-4. This K is in only fair agreement with the results of other investigators.^{53,62} We also adopt $\Delta H_f^\circ = -356.2 \text{ kcal mol}^{-1}$ for $\text{Cr}_2\text{O}_7^{2-}(\text{aq})$ from the results of calorimetric investigation⁴³ of reaction 3 and as listed in NBS 270-4. This ΔH_f° is in good agreement with the $d \ln K/dT$ results of Linge and Jones⁵⁹ and Davies and Prue⁶⁰ but in poorer agreement with the results of other investigators.⁶² Combination of ΔG_f° and ΔH_f° values leads to $S^\circ = 62.6 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{Cr}_2\text{O}_7^{2-}(\text{aq})$, as in NBS 270-4.

Several investigations^{56-58,61,63,64} have yielded results for reaction 3 in various constant ionic strength media.

Results of two investigations^{65,66} are in reasonable agreement with the following:




We use this K with the ΔG_f° of $\text{HCrO}_4^-(\text{aq})$ to calculate $\Delta G_f^\circ = -181.8 \text{ kcal mol}^{-1}$ for $\text{H}_2\text{CrO}_4(\text{aq})$. The $d \ln K/dT$ results of Tong and Johnson⁶⁵ permit calculation of $\Delta H_f^\circ = -201 \text{ kcal mol}^{-1}$ and $S^\circ = 70 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{H}_2\text{CrO}_4(\text{aq})$. Their investigations⁶⁵ have also led to the following:




$$K(1 \text{ M}) = 11$$

This K is in agreement with results of an earlier investigation.⁶⁶ We use the above⁶⁵ K and also $d \ln K/dT$ to calculate $\Delta G_f^\circ = -158.9 \text{ kcal mol}^{-1}$, $\Delta H_f^\circ = -180.4 \text{ kcal mol}^{-1}$, and $S^\circ = 49 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{CrO}_3\text{Cl}^-(\text{aq})$.


We also have the following equilibrium constants^{66,67} for other reactions of $\text{HCrO}_4^-(\text{aq})$:



$$K(3 \text{ M}) = 4$$




$$K(0.25 \text{ M}) = 3$$



$$K(0.25 \text{ M}) = 9$$

We combine the second and third of these equilibrium constants with ΔG_f° values for $\text{H}_2\text{PO}_4^-(\text{aq})$ and $\text{H}_3\text{PO}_4(\text{aq})$ from NBS 270-3 to obtain our tabulated ΔG_f° values for the Cr-P species. We do not do this same calculation for $\text{CrSO}_7^{2-}(\text{aq})$ because of the high ionic strength of the medium to which the K applies.

We also have



$$K(1 \text{ M}) = 0.43$$

from the work of Espenson and Helzer.⁶⁸

Two high-temperature investigations^{31,69} are in good agreement with $\Delta H_f^\circ = -176 \text{ kcal mol}^{-1}$ for $\text{H}_2\text{CrO}_4(\text{g})$, as compared with $-174 \text{ kcal mol}^{-1}$ listed in NBS 270-4.

We now consider the properties of various chromates and dichromates, beginning with $\text{Ag}_2\text{CrO}_4(\text{c})$. As previously stated, we have^{22,48} $S^\circ = 52.0 \text{ cal K}^{-1} \text{ mol}^{-1}$ from C_p measurements and the third law. We combined the calorimetric enthalpy of solution from Smith, Pitzer, and Latimer⁴⁸ with ΔH_f° values for $\text{Ag}^+(\text{aq})$ and $\text{CrO}_4^{2-}(\text{aq})$ to obtain $\Delta H_f^\circ = -174.58 \text{ kcal mol}^{-1}$ for $\text{Ag}_2\text{CrO}_4(\text{c})$. Further combination of ΔH_f° and S° leads to $\Delta G_f^\circ = -153.11 \text{ kcal mol}^{-1}$ for this compound and thence to a calculated $K_{sp} = 1.8 \times 10^{-12}$ for $\text{Ag}_2\text{CrO}_4(\text{c})$.

Ferrante, Stuve, and Krug⁷⁰ have made C_p measurements leading to $S^\circ = 42.212 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{Na}_2\text{CrO}_4(\text{c})$ at 298 K. They⁷⁰ have also made high-temperature measurements leading to values of $(H_T - H_{298})$ and related thermodynamic quantities. The enthalpy of solution⁷¹ leads to $\Delta H_f^\circ = -320.8 \text{ kcal mol}^{-1}$, which we combine with the entropy to obtain $\Delta G_f^\circ = -295.2 \text{ kcal mol}^{-1}$ for $\text{Na}_2\text{CrO}_4(\text{c})$.

The enthalpy of reaction⁷¹ of $\text{Na}_2\text{Cr}_2\text{O}_7(\text{c})$ with excess $\text{OH}^-(\text{aq})$ leads to its $\Delta H_f^\circ = -473.0 \text{ kcal mol}^{-1}$, which we combine with our estimated $S^\circ = (64) \text{ cal K}^{-1} \text{ mol}^{-1}$ to obtain $\Delta G_f^\circ = -430.3 \text{ kcal mol}^{-1}$ for $\text{Na}_2\text{Cr}_2\text{O}_7(\text{c})$.

We use the calorimetric enthalpy⁴³ of solution of $\text{K}_2\text{CrO}_4(\text{c})$ to obtain its $\Delta H_f^\circ = -335.4 \text{ kcal mol}^{-1}$ and combine with the entropy²² ($S^\circ = 47.8 \text{ cal K}^{-1} \text{ mol}^{-1}$) to obtain $\Delta G_f^\circ = -309.6 \text{ kcal mol}^{-1}$ for $\text{K}_2\text{CrO}_4(\text{c})$.

Measurements by Farber and Srivastava³¹ lead to $\Delta H_f^\circ = -249 \text{ kcal mol}^{-1}$ for $\text{K}_2\text{CrO}_4(\text{g})$ and to $\Delta H_f^\circ = -217 \text{ kcal mol}^{-1}$ for $\text{KHCrO}_4(\text{g})$.

The enthalpy of reaction⁷¹ of $\text{K}_2\text{Cr}_2\text{O}_7(\text{c})$ with excess $\text{OH}^-(\text{aq})$ leads to its $\Delta H_f^\circ = -492.9 \text{ kcal mol}^{-1}$. We combine this quantity with the $S^\circ = 69.6 \text{ cal K}^{-1} \text{ mol}^{-1}$ from Kelley and King²² to obtain $\Delta G_f^\circ = -450.0 \text{ kcal mol}^{-1}$ for $\text{K}_2\text{Cr}_2\text{O}_7(\text{c})$. We also note that Shmagin and Shidlovskii⁷² have measured enthalpies of dilution of supersaturated solutions of $\text{K}_2\text{Cr}_2\text{O}_7(\text{aq})$.

For $(\text{NH}_4)_2\text{CrO}_4(\text{c})$ we adopt $\Delta H_f^\circ = -279.0 \text{ kcal mol}^{-1}$ as listed in NBS 270-4, consistent with an old enthalpy of solution.⁴⁵

We also adopt $\Delta H_f^\circ = -321.1 \text{ kcal mol}^{-1}$ for $\text{MgCrO}_4(\text{c})$ as listed in NBS 270-6, based on results of older investigations that are cited in NBS Circular 500.⁷³


Shidlovskii, Voskresenskii, and Balakireva⁷⁴ have measured the enthalpy of solution of $\text{SrCrO}_4(\text{c})$ in acid with results that lead us to $\Delta H_f^\circ = -341.6 \text{ kcal mol}^{-1}$ for this compound. Sillén⁷⁵ has listed $K_{sp} = 2 \times 10^{-5}$ for $\text{SrCrO}_4(\text{c})$, which leads us to its $\Delta G_f^\circ = -314.0 \text{ kcal mol}^{-1}$. Combination of this ΔG_f° with the ΔH_f° gives $S^\circ = 23.6 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{SrCrO}_4(\text{c})$. This calculated entropy is smaller than seems reasonable. Combination of the ΔH_f° above with a "more reasonable" larger entropy leads to a calculated K_{sp} that is smaller than the listed⁷⁵ value. Further measurements are needed.

Shidlovskii et al.⁷⁴ have also measured the enthalpy of precipitation of $\text{BaCrO}_4(\text{c})$, which leads us to $\Delta H_f^\circ = -345.3 \text{ kcal mol}^{-1}$ for this compound. Taking $K_{sp} = 1.2 \times 10^{-10}$ from Beyer and Rieman,⁷⁶ we calculate $\Delta G_f^\circ = -321.53 \text{ kcal mol}^{-1}$ for $\text{BaCrO}_4(\text{c})$, which is the value listed in NBS 270-6. Combination of ΔH_f° and ΔG_f° values leads to $S^\circ = 38.9 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{BaCrO}_4(\text{c})$. Our values for the ΔH_f° and S° of $\text{BaCrO}_4(\text{c})$ differ slightly from those listed in NBS 270-6, possibly because we have used an enthalpy of precipitation that appeared too late for consideration there. A third law entropy would be a useful check on the other results.

Old (somewhat uncertain) calorimetric results⁴⁵ lead us to $\Delta H_f^\circ = -11 \text{ kcal mol}^{-1}$ for the enthalpy of precipitation of $\text{PbCrO}_4(\text{c})$ and thence to $\Delta H_f^\circ = -222 \text{ kcal mol}^{-1}$ for this compound. Taking $K_{sp} = 2.5 \times 10^{-13}$ from Kolthoff, Perlich, and Weiblen,⁷⁷ we calculate $\Delta G_f^\circ = -197.0 \text{ kcal mol}^{-1}$ for $\text{PbCrO}_4(\text{c})$ and combine this value with the ΔH_f° above to obtain $S^\circ = 35 \text{ cal K}^{-1} \text{ mol}^{-1}$ for this compound. On the basis of apparently reliably known entropies for other compounds of lead and barium, it seems safe to predict that the entropy (at 298 K) of $\text{PbCrO}_4(\text{c})$ should be larger than that of $\text{BaCrO}_4(\text{c})$. Because the S° we have calculated here for $\text{PbCrO}_4(\text{c})$ is smaller than that calculated in the preceding paragraph for $\text{BaCrO}_4(\text{c})$, we suggest that there is at least one error in the various properties listed above for these compounds. Further measurements are needed.

Sillén⁷⁵ has listed $K_{sp} = 9.77 \times 10^{-13}$ for $\text{Ti}_2\text{CrO}_4(\text{c})$, with

reference to Suzuki.⁷⁸ We use this K_{sp} to calculate $\Delta G_f^\circ = -205.82 \text{ kcal mol}^{-1}$ for $\text{Ti}_2\text{CrO}_4(\text{c})$, in close agreement with the value listed in NBS 270-4. The abstract⁷⁸ of Suzuki's paper quotes $\Delta H_f^\circ = -50.74 \text{ kcal mol}^{-1}$ [Neither ΔG° nor ΔS° is given in the abstract.] for the reaction



from which we calculate $\Delta H_f^\circ = -225.3 \text{ kcal mol}^{-1}$ for $\text{Ti}_2\text{CrO}_4(\text{c})$. About half of the difference between this value and the value ($-225.8 \text{ kcal mol}^{-1}$) listed in NBS 270-4 is due to the difference between our ΔH_f° for $\text{Ag}_2\text{CrO}_4(\text{c})$ and that listed in NBS 270-4. Combination of our ΔH_f° and ΔG_f° values leads to a calculated $S^\circ = 69.2 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{Ti}_2\text{CrO}_4(\text{c})$, as compared to $S^\circ = 67.5 \text{ cal K}^{-1} \text{ mol}^{-1}$ listed in NBS 270-4. Because entropies of various other compounds of silver and thallium lead us to estimate $S^\circ = (61) \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{Ti}_2\text{CrO}_4(\text{c})$, we suspect some error in the reported⁷⁸ data. We adopt ΔH_f° and S° values that are averages of the values we have quoted above and the values listed in NBS 270.4.

Gaultier and Pannetier⁷⁹ have investigated the thermodynamics of solid-state transitions and melting of Ti_2CrO_4 , while Natarajan and Secco⁷⁹ have investigated enthalpies of phase transitions of $\text{Ti}_2\text{CrO}_4(\text{c})$ and $\text{Ag}_2\text{CrO}_4(\text{c})$.

For the very reactive chromyl fluoride (CrO_2F_2) we have only vapor pressure data⁸⁰ from which we calculate $\Delta S^\circ = 18 \text{ cal K}^{-1} \text{ mol}^{-1}$ for melting and $\Delta S^\circ = 27 \text{ cal K}^{-1} \text{ mol}^{-1}$ for vaporization at the normal boiling point. Edwards, Falconer, and Sunder⁸¹ have reported a new synthesis of CrOF_4 . Their vapor pressure measurements show that there is a solid-state transition at 24.1°C ($\Delta H_f^\circ = 3.7 \text{ kcal mol}^{-1}$ and $\Delta S^\circ = 12 \text{ cal K}^{-1} \text{ mol}^{-1}$) and have also led to $\Delta S^\circ = 8 \text{ cal K}^{-1} \text{ mol}^{-1}$ for melting (55°C) and to $\Delta S^\circ = 27 \text{ cal K}^{-1} \text{ mol}^{-1}$ for vaporization at the normal boiling point (95°C). It is interesting to note that the sum of the entropies of transition and melting for CrOF_4 is close to the entropy of melting of CrO_2F_2 and that entropies of vaporization of both compounds suggest that both liquids are associated.

The properties listed in NBS 270-4 for $\text{CrO}_2\text{Cl}_2(\text{liq and g})$ are consistent with enthalpies of reaction,^{73,82,83} vapor pressures,^{73,82} and an entropy of $\text{CrO}_2\text{Cl}_2(\text{g})$ calculated from estimated molecular constants.

D. Cr(II) and Cr(III): Compounds and Aqueous Ions

The NBS 270-4 lists $S^\circ = 22.44 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{CrF}_3(\text{c})$, which is the value obtained by Hansen and Griffel⁸⁴ from their C_p measurements. Old results cited by Bichowsky and Rossini,⁴⁵ NBS Circular 500,⁷³ Brewer et al.,⁸⁵ and more recent high-temperature emf results from Tanaka, Yamaguchi, and Moriyama⁸⁶ lead to various values of the thermodynamic properties of $\text{CrF}_2(\text{c})$ and $\text{CrF}_3(\text{c})$. Weighting the recent results⁸⁶ most heavily leads us to values for the thermodynamic properties of these compounds that are consistent with those listed in NBS 270-4, which we adopt here. High-temperature measurements by Kent and Margrave⁸⁷ lead to $\Delta H_f^\circ = -99 \text{ kcal mol}^{-1}$ for $\text{CrF}_2(\text{g})$, as also listed in NBS 270-4, and to $\Delta H_f^\circ = 5 \text{ kcal mol}^{-1}$ for $\text{CrF}(\text{g})$. The enthalpy of sublimation of $\text{CrF}_3(\text{c})$ from Zmbov and Margrave⁸⁸ leads to $\Delta H_f^\circ = -217 \text{ kcal mol}^{-1}$ for $\text{CrF}_3(\text{g})$.

For $\text{CrCl}_2(\text{c})$ we adopt $S^\circ = 27.56 \text{ cal K}^{-1} \text{ mol}^{-1}$ from the C_p measurements of Stout and Chisholm,⁸⁹ whose review of earlier high-temperature equilibrium investigations has led to $\Delta H_f^\circ = -94.52 \text{ kcal mol}^{-1}$ and $\Delta G_f^\circ = -85.1 \text{ kcal mol}^{-1}$ for this compound. All of these values are the same as those listed in NBS 270-4. It might be noted that these values are not consistent with the recent high-temperature equilibrium results of Papapietro and Monnier.⁹⁰

The $\Delta H_f^\circ = -30.7 \text{ kcal mol}^{-1}$ for $\text{CrCl}_2(\text{g})$ that is listed in NBS 270-4 is consistent with the work of Maier,⁹¹ who has also calculated $S^\circ = 74.2 \text{ cal K}^{-1} \text{ mol}^{-1}$ for this compound. A more recent calculation⁹² leads to $S^\circ = 74.8 \text{ cal K}^{-1} \text{ mol}^{-1}$, which we adopt and use with the ΔH_f° above to obtain $\Delta G_f^\circ = -35.4 \text{ kcal mol}^{-1}$ for $\text{CrCl}_2(\text{g})$.

The NBS 270-4 lists $\Delta H_f^\circ = -111 \text{ kcal mol}^{-1}$ for $(\text{CrCl}_2)_2(\text{g})$, consistent with the high-temperature mass spectroscopic results of Schoonmaker, Friedman, and Porter.⁹³ Results from these investigators might also be used to calculate approximate ΔG_f° and S° values for $(\text{CrCl}_2)_2(\text{g})$.

The NBS 270-4 lists $S^\circ = 29.38 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{CrCl}_3(\text{c})$, which is the value derived by Hansen and Griffel⁸⁴ from their C_p measurements. A weighted average of results cited in NBS Circular 500⁷³ and by Sano⁹⁴ is consistent with $\Delta H_f^\circ = -133.0 \text{ kcal mol}^{-1}$ for $\text{CrCl}_3(\text{c})$ as listed in NBS 270-4 and as adopted here. These values lead to $\Delta G_f^\circ = -116.2 \text{ kcal mol}^{-1}$ for $\text{CrCl}_3(\text{c})$.

There are also two other routes to the ΔH_f° of $\text{CrCl}_3(\text{c})$. Vasil'kova et al.⁹⁵ have made calorimetric measurements with results that we combine with our tabulated ΔH_f° of $\text{CrO}_3(\text{c})$ to obtain $\Delta H_f^\circ = -124 \text{ kcal mol}^{-1}$ for $\text{CrCl}_3(\text{c})$. Or we may combine some of the results of Vasil'kova et al.⁹⁵ with the enthalpy of solution of $\text{CrCl}_3(\text{c})$ reported by Gregory and Burton⁹⁶ to obtain $\Delta H_f^\circ = -130 \text{ kcal mol}^{-1}$ for $\text{CrCl}_3(\text{c})$. Because it is not certain that the same Cr(III) species were formed in all of these experiments,^{95,96} this last calculation that involves combination of results obtained under different conditions may be unjustified.

Maier's⁹¹ work leads to $\Delta H_f^\circ = -71 \text{ kcal mol}^{-1}$, $S^\circ = 84 \text{ cal K}^{-1} \text{ mol}^{-1}$, and $\Delta G_f^\circ = -71 \text{ kcal mol}^{-1}$ for $\text{CrCl}_3(\text{g})$.

The NBS 270-4 lists $\Delta H_f^\circ = -72.2 \text{ kcal mol}^{-1}$ for $\text{CrBr}_2(\text{c})$, a value that is consistent with our calculations (± 1 or 2 kcal mol^{-1}) based on the high-temperature equilibrium results of Shchukarev et al.⁹⁷ We estimate $S^\circ = (32) \text{ cal K}^{-1} \text{ mol}^{-1}$ and combine with the ΔH_f° above to obtain $\Delta G_f^\circ = -69 \text{ kcal mol}^{-1}$ for $\text{CrBr}_2(\text{c})$. The high-temperature vaporization measurements of Sime and Gregory⁹⁸ and the mass spectral investigations of Schoonmaker et al.⁹³ lead to $\Delta H_f^\circ = -17 \text{ kcal mol}^{-1}$ for $\text{CrBr}_2(\text{g})$ and $\Delta H_f^\circ = -84 \text{ kcal mol}^{-1}$ for $(\text{CrBr}_2)_2(\text{g})$ as listed in NBS 270-4. We also have $S^\circ = 80.62 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{CrBr}_2(\text{g})$ from Shaw and Porter⁹² and thence $\Delta G_f^\circ = -28.5 \text{ kcal mol}^{-1}$ for $\text{CrBr}_2(\text{g})$.

It is possible to use quantities reported by Rupcheva et al.⁹⁹ to calculate $\Delta H_f^\circ = -97.4$ and $-76.6 \text{ kcal mol}^{-1}$ for $\text{CrBr}_3(\text{c})$, with no clear indication as to the reliability of either value. We therefore omit $\text{CrBr}_3(\text{c})$ from our Table I. It might also be noted that the properties listed by Rupcheva et al.⁹⁹ for $\text{CrBr}_2(\text{c})$ are much different from those cited above.

Combination of the calorimetric results of Gregory and Burton⁹⁶ with our tabulated ΔH_f° of $\text{CrCl}_2(\text{c})$ leads to $\Delta H_f^\circ = -37.5 \text{ kcal mol}^{-1}$ for $\text{CrI}_2(\text{c})$, which is also the value listed in NBS 270-4 for this compound. We estimate $S^\circ = (37) \text{ cal K}^{-1} \text{ mol}^{-1}$ and combine with the ΔH_f° to obtain $\Delta G_f^\circ = -39 \text{ kcal mol}^{-1}$ for $\text{CrI}_2(\text{c})$.

Allen¹⁰⁰ has made vapor pressure measurements on $\text{CrI}_2(\text{c})$ and has used a second law method to calculate ΔH_f° of vaporization at 298 K , which we combine with the ΔH_f° of $\text{CrI}_2(\text{c})$ to obtain $\Delta H_f^\circ = 34 \text{ kcal mol}^{-1}$ for $\text{CrI}_2(\text{g})$, as compared with $\Delta H_f^\circ = 24 \text{ kcal mol}^{-1}$ listed in NBS 270-4. Our third law treatment of Allen's¹⁰⁰ results leads to somewhat smaller values for ΔH_f° of vaporization and ΔH_f° . We adopt $\Delta H_f^\circ = 32 \text{ kcal mol}^{-1}$ for $\text{CrI}_2(\text{g})$ and combine with $S^\circ = 84.15 \text{ cal K}^{-1}$ from Shaw and Porter⁹² (whose results were also used in our third law calculations) to obtain $\Delta G_f^\circ = 17 \text{ kcal mol}^{-1}$ for $\text{CrI}_2(\text{g})$.

Combination of the calorimetric results of Gregory and Burton⁹⁶ with our tabulated ΔH_f° of $\text{CrCl}_3(\text{c})$ leads to $\Delta H_f^\circ = -45.8 \text{ kcal mol}^{-1}$ for $\text{CrI}_3(\text{c})$. We also use the results of de-

TABLE I.^a Thermodynamic Properties of Chromium Compounds at 298 K

Substance	ΔH_f° kcal mol ⁻¹	ΔG_f° kcal mol ⁻¹	S° cal K ⁻¹ mol ⁻¹	Substance	ΔH_f° kcal mol ⁻¹	ΔG_f° kcal mol ⁻¹	S° cal K ⁻¹ mol ⁻¹
Cr(c)	0	0	5.68 ²²⁻²⁴	Cr ²⁺ (aq)	-34.3 ^{45,96}	-35	(-24)
Cr(g)	95.0 ²⁶⁻²⁸	84.3	41.68	Cr(H ₂ O) ₆ ³⁺ (aq)	~-467	~-386.6	~24
Cr ⁺ (g)	252.5 ^b			Cr ³⁺ (aq)	~-57	~-46.5	~-76
Cr ²⁺ (g)	634.4 ^b			Cr(H ₂ O) ₅ (OH) ²⁺ (aq)	-457 ¹¹⁴⁻¹¹⁶	-381.4 ^{75,114,115}	~42
Cr ³⁺ (g)	1350			Cr(OH) ²⁺ (aq, i.s. ^c)	-115 ¹¹⁴⁻¹¹⁶	-98.0 ^{75,114,115}	~-42
Cr ⁴⁺ (g)	2495			Cr(H ₂ O) ₄ (OH) ₂ ⁺ (aq)	-449 ^{73,75,114}	-372 ^{75,114}	~35
Cr ⁵⁺ (g)	4182			Cr(OH) ₂ ⁺ (aq, i.s.)	-176 ^{73,75,114}	-145 ^{75,114}	~-32
Cr ⁶⁺ (g)	6273			Cr(OH) ₃ (ppt)		-200.2 ^{75,114}	
Cr ⁷⁺ (g)	9991			Cr(OH) ₄ ⁻ (aq)	-298 ¹¹⁷	~-236 ⁷⁵	-26 ¹¹⁷
CrO(g)	~53 ²⁹	~46	53.0 ²²	Cr(H ₂ O) ₅ Cl ²⁺ (aq)	-434 ¹²⁰⁻¹²²	-360.3 ¹¹⁹⁻¹²³	35
CrO ₂ (c)	-143	-131	(13)	CrCl ²⁺ (aq, i.s.)	-92 ¹²⁰⁻¹²²	-76.9 ¹¹⁹⁻¹²³	-49
CrO ₂ (g)	~-3 ²⁹⁻³¹	~-5	62.0 ³²	Cr(H ₂ O) ₆ Cl ²⁺ (aq, o.s. ^d)		-417.9 ¹²³	
Cr ₂ O ₃ (c)	-272.4	-252.9	19.4 ²²	CrCl ²⁺ (aq, o.s.)		-77.8 ¹²³	
CrO _{2.40} (c)	-140.3 ⁴⁶			Cr(H ₂ O) ₄ Cl ₂ ⁺ (aq)	-401 ^{117,122}	-336.3 ^{75,112}	~52
CrO _{2.62} (c)	-139.1 ⁴⁶			CrCl ₂ ⁺ (aq, i.s.)	-127 ^{117,122}	-109.6 ^{75,112}	~-15
CrO ₃ (c)	-140.9 ^{43,44}	-122	(16)	Cr(H ₂ O) ₃ (OH)Cl ₂ (aq)		-328.5 ¹²⁴	
CrO ₃ (g)	~-65 ^{29,31}	~-60	63.6 ³²	Cr(OH)Cl ₂ (aq, i.s.)		-158.4 ¹²⁴	
CrO ₄ ²⁻ (aq)	-210.60 ⁴³	-173.9 ₆	12 ⁴⁸⁻⁵¹	Cr(H ₂ O) ₅ Br ²⁺ (aq)	-422.7 ¹²⁶	-351.1 ¹²⁶	41.5 ¹²⁶
HCrO ₄ ⁻ (aq)	-209.9 ⁴³	-182.8 ⁶²⁻⁵⁷	44.0	CrBr ²⁺ (aq, i.s.)	-81.1 ¹²⁶	-67.7 ¹²⁶	-42.0 ¹²⁶
H ₂ CrO ₄ (aq)	-201 ⁶⁵	-181.8 ^{65,66}	70 ⁶⁵	Cr(H ₂ O) ₄ Br ₂ ⁺ (aq)	-377.1 ⁴⁵		
H ₂ CrO ₄ (g)	-176 ^{31,69}			CrBr ₂ ⁺ (aq, i.s.)	-103.8 ⁴⁵		
Cr ₂ O ₇ ²⁻ (aq)	-356.2 ⁴³	-311.0 ⁶⁹⁻⁶¹	62.6	Cr(H ₂ O) ₅ SCN ²⁺ (aq)	-382.5 ¹¹⁶	-312.0 ¹¹⁶	~49 ¹¹⁶
CrO ₄ ³⁻ (aq)		-176.3 ¹⁸		CrSCN ²⁺ (aq, i.s.)	-40.9 ¹¹⁶	-28.6 ¹¹⁶	~-35 ¹¹⁶
CrO ₃ Cl ⁻ (aq)	-180.4 ⁶⁵	-158.9 ^{65,66}	49 ⁶⁵	FeCr ₂ O ₄ (c)	-345.3 ^{39,145-147}	-321.2	34.9 ²²
HCrPO ₇ ²⁻ (aq)		-396.9 ⁶⁷		MgCr ₂ O ₄ (c)	-426.9 ¹⁴⁸	-399.5	25.3 ⁴²²
H ₂ CrPO ₇ ⁻ (aq)		-400.5 ⁶⁷		NiCr ₂ O ₄ (c)	-330.6 ¹⁴⁸	-305.3	(31.5)
Ag ₂ CrO ₄ (c)	-174.58 ⁴⁸	-153.11	52.0 ^{22,48}	ZnCr ₂ O ₄ (c)	-370.9 ¹⁴⁸	-344.4	(30.5)
Na ₂ CrO ₄ (c)	-320.8 ⁷¹	-295.2	42.2 ¹²⁷⁰	CdCr ₂ O ₄ (c)	-344.3 ¹⁴⁸	-318.1	(34)
Na ₂ Cr ₂ O ₇ (c)	-473.0 ⁷¹	-430.3	(64)	CuCr ₂ O ₄ (c)	-307.3 ¹⁴⁸	-282.2	(33)
K ₂ CrO ₄ (c)	-335.4 ⁴³	-309.6	47.8 ²²	NaCrO ₂ (c)	-210 ¹¹⁰		
K ₂ CrO ₄ (g)	-249 ³¹			NH ₄ Cr(SO ₄) ₂ ·12H ₂ O(c)			170.9 ²²
KHCrO ₄ (g)	-217 ³¹			CrH _{0.84} (c)			8.98 ¹⁵⁸
K ₂ Cr ₂ O ₇ (c)	-492.9 ⁷¹	-450.0	69.6 ²²	CrH _{0.91} (c)			8.06 ¹⁵⁸
(NH ₄) ₂ CrO ₄ (c)	-279.0 ⁴⁵			CrH _{0.94} (c)			8.20 ¹⁵⁸
(NH ₄) ₂ Cr ₂ O ₇ (c)	-431.8 ⁴⁴			Cr ₇ H ₂ (c)	-3.8 ¹⁵⁹		
MgCrO ₄ (c)	-321.1 ⁷³			Cr ₃ C ₂ (c)	-19.3	-19.5	20.4 ²²
SrCrO ₄ (c)	-341.6 ⁷⁴	-314.0 ⁷⁵	23.6 (?)	Cr ₇ C ₃ (c)	-38.7	-39.9	48.0 ²²
BaCrO ₄ (c)	-345.3 ⁷⁴	-321.5 ²⁷⁶	38.9 (?)	Cr ₂₃ C ₆ (c)	-87.2	-89.3	145.8 ²²
PbCrO ₄ (c)	-222 ⁴⁵	-197.0 ⁷⁷	35 (?)	Cr ₂ (g)	~184 ¹⁶³		
Tl ₂ CrO ₄ (c)	-225.6 ⁷⁸	-205.8 ₂	68 (?)	CrN(c)	-29 ^{45,73,164,165}	-22	(6)
CrO ₂ Cl ₂ (liq)	-138.5 ^{73,82,83}	-122.1 ^{73,82}	53.0	Cr ₂ N(c)	-30 ^{45,73,164,165}	-18	(12)
CrO ₂ Cl ₂ (g)	-128.6 ^{73,82}	-119.9 ^{73,82}	78.8	CrSi(c)	-12.7 ¹⁶⁶	-12.9	10.5 ¹⁶⁶
CrF(g)	5 ⁸⁷			CrSi ₂ (c)	-19.1 ¹⁶⁶	-18.7	13.3 ^{24,166,167}
CrF ₂ (c)	-186	-176	(20)	Cr ₃ Si(c)	-22.0 ¹⁶⁶	-21.8	20.6 ¹⁶⁶
CrF ₂ (g)	-99 ⁸⁷			Cr ₅ Si ₃ (c)	-50.4 ¹⁶⁶	-50.9	43.5 ¹⁶⁶
CrF ₃ (c)	-277 ^{45,73,85,86}	-260	22.44 ⁸⁴	CrGe(c)			14.6 ¹⁶⁸
CrF ₃ (g)	-217 ⁸⁸			Cr ₅ Ge ₃ (c)			56.6 ¹⁶⁸
CrF ₄ (c)	-298 ¹⁵³	-277	(33)	Cr _{1,1} Ge ₁₉ (c)			227.3 ¹⁶⁸
CrCl ₂ (c)	-94.5 ₂ ⁸⁹	-85.1 ⁸⁹	27.56 ⁸⁹	CrS(c)	-32 ¹⁶⁹	-33 ¹⁶⁹	17
CrCl ₂ (g)	-30.7 ⁹¹	-35.4	74.8 ⁹²	CrS(g)	~83 ¹⁷⁰	~69	(60)
(CrCl ₂) ₂ (g)	-111 ⁹³			Cr ₂ Te ₃ (c)			49.86 ¹⁷¹
CrCl ₃ (c)	-133.0 ^{73,94}	-116.2	29.38 ⁸⁴	Cr ₃ Te ₄ (c)			70.05 ¹⁷¹
CrCl ₃ (g)	-71 ⁹¹	-71 ⁹¹	84 ⁹¹	Cr ₅ Te ₆ (c)			112.49 ¹⁷¹
CrCl ₄ (g)	-102 ^{91,154,155}	-97.5	87.5 ⁹¹	Cr(CO) ₆ (c)	-234(?) ¹⁷⁵	-207	72 ^{173,176}
CrBr ₂ (c)	-72.2 ⁹⁷	-69	(32)	Cr(CO) ₆ (g)	-217 ^{173,176}	-202	111.6 ¹⁷⁸
CrBr ₂ (g)	-179 ^{93,98}	-28.5	80.62 ⁹²	Cr(PF ₃) ₆ (c)	-1446.4 ¹⁷⁹		
(CrBr ₂) ₂ (g)	-84 ^{93,98}						
CrI ₂ (c)	-37.5 ⁹⁶	-39	(37)				
CrI ₂ (g)	32 ¹⁰⁰	17	84.15 ⁹²				
CrI ₃ (c)	-49 ^{96,101,102}	-49	47 ^{101,102}				
CrI ₄ (g)	3 ¹⁰²	-12	112 ¹⁰²				
CrICl ₂ (c)	-100 ¹⁰³						
CrIBr ₂ (c)	-79 ¹⁰³						

^a Italicized values agree with values in NBS 270-4 or NBS 270-6. Values in parentheses are estimates. ^b From NBS 270-4, adjusted for different ΔH_f° of Cr(g). ^c i.s. = inner sphere. ^d o.s. = outer sphere.

composition studies on CrI₃(c) by Handy and Gregory¹⁰¹ and by Shieh and Gregory¹⁰² with our tabulated ΔH_f° for CrI₂(c) to obtain $\Delta H_f^\circ = -50$ kcal mol⁻¹ for CrI₃(c). Their results^{101,102} are also consistent with $S^\circ = 47$ cal K⁻¹ mol⁻¹

and thence $\Delta G_f^\circ = -50$ kcal mol⁻¹ for CrI₃(c). We adopt $\Delta H_f^\circ = -49$ kcal mol⁻¹, $S^\circ = 47$ cal K⁻¹ mol⁻¹, and $\Delta G_f^\circ = -49$ kcal mol⁻¹ for CrI₃(c).

Handy and Gregory¹⁰³ have made high-temperature equi-

librium measurements on decomposition of $\text{CrCl}_2(\text{c})$ and $\text{CrBr}_2(\text{c})$, with results that we combine with our tabulated ΔH_f° values for $\text{CrCl}_2(\text{c})$ and $\text{CrBr}_2(\text{c})$ to obtain $\Delta H_f^\circ = -100$ kcal mol⁻¹ for $\text{CrCl}_2(\text{c})$ and $\Delta H_f^\circ = -79$ kcal mol⁻¹ for $\text{CrBr}_2(\text{c})$. These same values are listed in NBS 270-4. We use the thermodynamic properties to calculate $\Delta H^\circ = -16$ kcal mol⁻¹ of $\text{CrI}_3(\text{c})$ for




This indication that $\text{CrCl}_2(\text{c})$ is unstable with respect to the CrX_3 compounds suggests that further work is needed.

Lascelles, Shelton, and Schäfer¹⁰⁴ have carried out high-temperature measurements on the $\text{CrCl}_2\text{-AlCl}_3$ system. We are unable to use either their reported K values or their equation for $K(T)$, which is inconsistent with the reported K values, for calculation of thermodynamic properties to list in Table I for the postulated $\text{CrCl}_2\cdot 2\text{AlCl}_3(\text{g})$.

Vasil'kova et al.¹⁰⁵ have reported results of their calorimetric measurements on various "chlorochromates (III)" and summarized results of some related earlier work. Although it is possible to calculate values of ΔH_f° (from the elements) for various $\text{M}_3\text{CrCl}_6(\text{c})$ and $\text{M}_3\text{Cr}_2\text{Cl}_9(\text{c})$ compounds ($M =$ alkali metal) from their results, it may be more useful to have available the enthalpies of formation from the constituent halides as in Table II. We also call attention to related high-temperature vaporization measurements by Galitskii et al.,¹⁰⁶ whose results do not permit us to calculate any thermodynamic properties for our tables. It should also be noted that Cook¹⁰⁷ has concluded from phase equilibrium data that $\text{Na}_3\text{Cr}_2\text{Cl}_9(\text{c})$ is apparently nonexistent.

Now we turn to consideration of the thermodynamic properties of aqueous chromic [Cr(III)] and chromous [Cr(II)] ions, beginning with the latter and then continuing with a large number of Cr(III) species.

Gregory and Burton⁹⁶ have measured the enthalpy of solution of $\text{CrCl}_2(\text{c})$ in 1.0 M $\text{Cl}^-(\text{aq})$, with a result in fair agreement with an earlier⁴⁵ enthalpy of solution. Combination of these experimental results with our estimated enthalpies of dilution and the ΔH_f° of $\text{CrCl}_2(\text{c})$ is consistent with $\Delta H_f^\circ = -34.3$ kcal mol⁻¹ for $\text{Cr}^{2+}(\text{aq})$ as listed in NBS 270-4. We estimate $S^\circ = (-24)$ cal K⁻¹ mol⁻¹ and combine with the ΔH_f° to obtain $\Delta G_f^\circ = -35$ kcal mol⁻¹ for $\text{Cr}^{2+}(\text{aq})$. This ΔG_f° leads to the potential



Jenkins, Mamantov, and Manning¹⁰⁸ have reported E° for the $\text{Cr(II)}|\text{Cr(c)}$ couple in molten fluoride medium.

The NBS 270-4 lists $\Delta H_f^\circ = -477.8$ kcal mol⁻¹ for $\text{Cr}(\text{H}_2\text{O})_6^{3+}(\text{aq})$, which corresponds to $\Delta H_f^\circ = -477.8 - 6$ (-68.315) = -67.9 kcal mol⁻¹ for $\text{Cr}^{3+}(\text{aq})$ in which we do not explicitly include any water. Our assessment of various experimental results leads us to a considerably less negative ΔH_f° as described below.

The calorimetrically determined¹⁰⁹ enthalpy of reduction of Cr(VI) in acidic solution to Cr(III) in combination with our tabulated properties of $\text{Cr}_2\text{O}_7^{2-}(\text{aq})$ and $\text{HCrO}_4^-(\text{aq})$ leads to $\Delta H_f^\circ = -58$ kcal mol⁻¹ for $\text{Cr}^{3+}(\text{aq})$.

We have other calorimetric results (cited below) that we can use for evaluation of ΔH_f° of $\text{Cr}^{3+}(\text{aq})$. For this purpose we must make use of ΔH° values for dissociation of $\text{CrCl}_2^{2+}(\text{aq})$ and $\text{CrCl}_2^+(\text{aq})$ as discussed later in this section.

Neuman, Kroger, and Kunz (cited by Bichowsky and Rossini⁴⁵) have determined the enthalpy of solution of Cr(c) in $\text{HCl}(\text{aq})$. Assuming that $\text{CrCl}_2^+(\text{aq})$ was the principal Cr(III) species in their solutions, we obtain $\Delta H_f^\circ = -61$ kcal mol⁻¹ for $\text{Cr}^{3+}(\text{aq})$. Assuming that $\text{CrCl}_2^{2+}(\text{aq})$ was the principal species leads to $\Delta H_f^\circ = -56$ kcal mol⁻¹ for $\text{Cr}^{3+}(\text{aq})$.

Gregory and Burton⁹⁶ have measured the enthalpy of solution of $\text{CrCl}_3(\text{c})$ in 1.0 M $\text{Cl}^-(\text{aq})$. Assuming that $\text{CrCl}_2^+(\text{aq})$

TABLE II. Enthalpies of Reaction¹⁰⁵

$$m\text{MCl}(\text{c}) + n\text{CrCl}_3(\text{c}) = \text{M}_n\text{Cr}_m\text{Cl}_{3m+n}(\text{c})$$

$\text{M}_n\text{Cr}_m\text{Cl}_{3m+n}$	$\Delta H^\circ/\text{kcal mol}^{-1}$	$\text{M}_n\text{Cr}_m\text{Cl}_{3m+n}$	$\Delta H^\circ/\text{kcal mol}^{-1}$
Na_3CrCl_6	0.7	$\text{NaRb}_2\text{CrCl}_6$	-14.9
K_3CrCl_6	-11.0	$\text{NaCs}_2\text{CrCl}_6$	-20.6
Rb_3CrCl_6	-16.0	$\text{K}_3\text{Cr}_2\text{Cl}_9$	-12.7
Cs_3CrCl_6	-18.4	$\text{Rb}_3\text{Cr}_2\text{Cl}_9$	-25.8
$\text{NaK}_2\text{CrCl}_6$	-8.0	$\text{Cs}_3\text{Cr}_2\text{Cl}_9$	-34.7



was the principal species in solution, we calculate $\Delta H_f^\circ = -56$ kcal mol⁻¹ for $\text{Cr}^{3+}(\text{aq})$, while the choice of $\text{CrCl}_2^{2+}(\text{aq})$ as principal species in solution leads to $\Delta H_f^\circ = -51$ kcal mol⁻¹ for $\text{Cr}^{3+}(\text{aq})$.

Shchukarev et al.⁹⁵ have also measured the enthalpy of solution of $\text{CrCl}_3(\text{c})$, using 0.12 M $\text{Cl}^-(\text{aq})$ as solvent. Assuming that $\text{CrCl}_2^{2+}(\text{aq})$ was the principal species in solution leads to $\Delta H_f^\circ = -57$ kcal mol⁻¹ for $\text{Cr}^{3+}(\text{aq})$, while choice of $\text{CrCl}_2^+(\text{aq})$ as principal solute species leads to $\Delta H_f^\circ = -62$ kcal mol⁻¹ for $\text{Cr}^{3+}(\text{aq})$.


On the basis of all of the results above [$\Delta H_f^\circ = -58, -61, -56, -56, -51, -57,$ and -62 kcal mol⁻¹ for $\text{Cr}^{3+}(\text{aq})$], we choose $\Delta H_f^\circ \approx -57$ kcal mol⁻¹ for $\text{Cr}^{3+}(\text{aq})$, which corresponds to $\Delta H_f^\circ \approx -467$ kcal mol⁻¹ for $\text{Cr}(\text{H}_2\text{O})_6^{3+}(\text{aq})$. These values are 11 kcal mol⁻¹ less negative than corresponding values from NBS 270-4. Further measurements are required to establish satisfactorily the thermodynamic properties of this important ion.

We note that Gross, Wilson, and Gutteridge¹¹⁰ have measured the enthalpy of solution of Cr(c) in aqueous HCl-FeCl_3 at 110 °C with results that are roughly consistent with our choice above.

Now we have two unsatisfactory routes to the ΔG_f° and S° of $\text{Cr}^{3+}(\text{aq})$. Grube and Schlecht¹¹¹ have measured potentials for $\text{Cr(III)}|\text{Cr(II)}$ couples in several solutions, using several different "inert" electrodes. We select $E^\circ \approx -0.4_3$ V (uncertainty $\pm 0.1_3$ V?) for the $\text{Cr}^{3+}(\text{aq})|\text{Cr}^{2+}(\text{aq})$ couple and combine with the ΔG_f° of $\text{Cr}^{2+}(\text{aq})$ cited above to obtain $\Delta G_f^\circ \approx -45$ kcal mol⁻¹ for $\text{Cr}^{3+}(\text{aq})$. Combination of this ΔG_f° with our selected $\Delta H_f^\circ = -57$ kcal mol⁻¹ for $\text{Cr}^{3+}(\text{aq})$ leads to $S^\circ \approx -81$ cal K⁻¹ mol⁻¹ for this ion. Because this entropy seems to be too negative in comparison with entropies of some other 3+ aqueous ions, we also consider an alternative approach that begins with an estimated (more reasonable?) $S^\circ \approx (-72)$ cal K⁻¹ mol⁻¹ for $\text{Cr}^{3+}(\text{aq})$. Now combination of this entropy with our selected ΔH_f° leads to $\Delta G_f^\circ \approx -48$ kcal mol⁻¹ for $\text{Cr}^{3+}(\text{aq})$ and thence to a calculated $E^\circ \approx -0.5_6$ V for the $\text{Cr}^{3+}(\text{aq})|\text{Cr}^{2+}(\text{aq})$ couple. Because of substantial uncertainties throughout the calculations summarized here, we arbitrarily select mean values as follows: $\Delta G_f^\circ \approx -46.5$ kcal mol⁻¹ and $S^\circ \approx -76$ cal K⁻¹ mol⁻¹ for $\text{Cr}^{3+}(\text{aq})$. These values correspond to $\Delta G_f^\circ \approx -386.6$ kcal mol⁻¹ and $S^\circ \approx 24$ cal K⁻¹ mol⁻¹ for $\text{Cr}(\text{H}_2\text{O})_6^{3+}(\text{aq})$ in which specifically indicated waters of solvation are included in the thermodynamic properties. Our selected ΔG_f° values correspond to the following potentials:



$$E^\circ = 1.29 \text{ V}$$




$$E^\circ = 1.31 \text{ V}$$

Jenkins, Mamantov, and Manning¹⁰⁸ have measured the $\text{Cr(III)}|\text{Cr(II)}$ potential in molten fluoride medium.

In the early years of this century N. Bjerrum found it neces-

sary to interpret his emf determinations of $H^+(aq)$ in Cr(III) solutions in terms of hydrolytic dimers and polymers as well as such monomeric species as $Cr(OH)^{2+}(aq)$. Since that time a number of investigators have confirmed the importance of species containing more than one Cr(III), and by taking advantage of kinetic inertness have isolated a few such species. Results^{75,112,113} are summarized briefly as follows. Solutions containing appreciable concentrations of dimers and larger species can be obtained by refluxing chromium(III) perchlorate solutions, by making the pH of many solutions containing Cr(III) greater than about 3, and by oxidation of Cr(II) or reduction of Cr(VI) solutions. Dimeric and trimeric species bearing charges of +4 and +5 have been identified and equilibrium quotients for their formation have been reported. Dimers and larger species must certainly be considered in connection with many solutions in equilibrium with precipitated chromic hydroxide.

We begin our consideration of hydrolytic species of Cr(III) with the first hydroxy complex, which can be represented by $Cr(H_2O)_5(OH)^{2+}(aq)$ or by $Cr(OH)^{2+}(aq)$, inner sphere). Meyenburg, Siroky, and Schwarzenbach¹¹⁴ have recently investigated the equilibrium that we represent by




at 5 and 20 °C in 0.1 M solution. Swaddle and Kong¹¹⁵ have investigated the same reaction at 25° and other temperatures in 0.5 M solution. Earlier investigations have been cited by Sillén.⁷⁵ On the basis of all these results, we choose K ($\mu = 0$, 298 K) = 1.6×10^{-4} for the equilibrium represented by (4). We use this value with our ΔG_f° of $Cr(H_2O)_6^{3+}(aq)$ to calculate $\Delta G_f^\circ = -381.4$ kcal mol⁻¹ for $Cr(H_2O)_5(OH)^{2+}(aq)$. We also obtain $\Delta G_f^\circ = -98.0$ kcal mol⁻¹ for the equivalent $Cr(OH)^{2+}(aq)$, inner sphere).

The NBS 270-4 ΔH_f° values correspond to $\Delta H^\circ = 14.2$ kcal mol⁻¹ for (4), while values tabulated in the older NBS Circular 500⁷³ lead to $\Delta H^\circ = 14.3$ kcal mol⁻¹ for this reaction. Results in the papers cited there⁷³ suggest to us that a smaller ΔH° might be better for this reaction. Three more recent investigations¹¹⁴⁻¹¹⁶ of equilibrium constants at different temperatures are all consistent with $\Delta H^\circ = 10$ kcal mol⁻¹ for the reaction represented by (4), which is the value we adopt and use in calculating $\Delta H_f^\circ = -457$ kcal mol⁻¹ for $Cr(H_2O)_5(OH)^{2+}(aq)$. Combination of ΔH_f° and ΔG_f° values leads to $S^\circ = 42$ cal K⁻¹ mol⁻¹ for $Cr(H_2O)_5(OH)^{2+}(aq)$. We also obtain $\Delta H_f^\circ = -115$ kcal mol⁻¹ and $S^\circ = -42$ cal K⁻¹ mol⁻¹ for the equivalent $Cr(OH)^{2+}(aq)$, inner sphere). This latter entropy is more negative than is usual for 2+ ions, which suggests that there may be an error in the results for reaction 4 or that there is an error in the entropy adopted for $Cr^{3+}(aq)$.

It is important to emphasize here that the ΔH_f° , ΔG_f° , and S° values in the paragraph above are based on our choices for $Cr(H_2O)_6^{3+}(aq)$ and $Cr^{3+}(aq)$, with our ΔH_f° values for these ions 11 kcal mol⁻¹ less negative than previously selected in NBS 270-4. Further, our ΔH_f° values for $Cr(H_2O)_5(OH)^{2+}(aq)$ and $Cr(OH)^{2+}(aq)$, inner sphere) are different from those in NBS 270-4 because we have selected a different ΔH° for reaction 4 than was selected by the authors of NBS 270-4.


For the second hydrolysis of $Cr^{3+}(aq)$ that can be represented by



we choose $K = 10^{-7}$ from the results of several investigations.^{75,114} The ΔH_f° values listed in NBS 270-4 correspond to $\Delta H^\circ = 7.8$ kcal mol⁻¹ for this reaction, in agreement with results of Bjerrum cited previously^{73,75} and with the more recent results of Meyenburg, Siroky, and Schwarzenbach.¹¹⁴ All of these results, in combination with our tabulated properties of $Cr(H_2O)_5(OH)^{2+}(aq)$, lead to $\Delta G_f^\circ = -372$ kcal mol⁻¹,



$\Delta H_f^\circ = -449$ kcal mol⁻¹, and $S^\circ \approx 35$ cal K⁻¹ mol⁻¹ for $Cr(H_2O)_4(OH)_2^+(aq)$. We also obtain $\Delta G_f^\circ = -145$ kcal mol⁻¹, $\Delta H_f^\circ = -176$ kcal mol⁻¹, and $S^\circ \approx -32$ cal K⁻¹ mol⁻¹ for the equivalent $Cr(OH)_2^+(aq)$, inner sphere). We note that this latter entropy is considerably more negative than is usual for 1+ ions.

On the basis of old⁷⁵ and recent¹¹⁴ investigations, we choose the following solubility product for precipitated chromic hydroxide:



Combination of this K_{sp} with our ΔG_f° for $Cr^{3+}(aq)$ leads to $\Delta G_f^\circ = -200.2$ kcal mol⁻¹ for $Cr(OH)_3(ppt)$. The ΔH° of the reaction above that is derived from the NBS 270-4 ΔH_f° values in combination with our other thermodynamic properties leads to a negative S° for $Cr(OH)_3(ppt)$, which is clearly unacceptable. We may also obtain an approximate ΔH° for this reaction from the work of Meyenburg, Siroky, and Schwarzenbach,¹¹⁴ which leads to $S^\circ \approx 8$ cal K⁻¹ mol⁻¹ for $Cr(OH)_3(ppt)$. This S° is only about one-third of a "reasonable" estimate for this quantity. In view of these calculations, we omit ΔH_f° and S° values for $Cr(OH)_3(ppt)$ from our Table I.

It is well established that $Cr(OH)_3(ppt)$ and various other Cr(III) compounds dissolve in excess $OH^-(aq)$ to yield anions containing Cr(III). The work of Schug and King¹¹⁷ has shown that these "chromite" solutions cannot be realistically described in any simple way. Nevertheless, we use results summarized by Sillén⁷⁵ to calculate an approximate $\Delta G_f^\circ \approx -236$ kcal mol⁻¹ for "chromite" ion that we represent in over-simplified fashion by $Cr(OH)_4^-(aq)$. This ΔG_f° is consistent with the following:




$$E^\circ \approx -0.21 \text{ V}$$

We also use calorimetric results from Schug and King¹¹⁷ to obtain $\Delta H_f^\circ = -298$ kcal mol⁻¹ and $S^\circ \approx -26$ cal K⁻¹ mol⁻¹ for $Cr(OH)_4^-(aq)$. In connection with these thermodynamic properties it must be emphasized that Schug and King¹¹⁷ have stated clearly that "chromite" solutions do *not* have the properties that might be expected for solutions of " $Cr(OH)_4^-(aq)$."


The stability of $CrF^{2+}(aq)$ has been investigated by Swaddle and King,¹¹⁸ who have also cited earlier work. The $CrF^{2+}(aq)$ ion is less stable with respect to dissociation than the corresponding ions of iron, scandium, and aluminum, but is more stable than similar ions of indium, yttrium, and lanthanum.

Results of several investigators¹¹⁹⁻¹²³ are in reasonable agreement with $K = 0.2$ for the reaction



This K value leads to $\Delta G_f^\circ = -360.3$ kcal mol⁻¹ for $Cr(H_2O)_5Cl^{2+}(aq)$ and also to $\Delta G_f^\circ = -76.9$ kcal mol⁻¹ for the equivalent $CrCl^{2+}(aq)$, inner sphere). Results of some of these investigations¹²⁰⁻¹²² are also consistent with $\Delta H^\circ = 5$ kcal mol⁻¹ for this reaction. We calculate $\Delta H_f^\circ = -434$ kcal mol⁻¹ and $S^\circ = 35$ cal K⁻¹ mol⁻¹ for $Cr(H_2O)_5Cl^{2+}(aq)$, and also obtain $\Delta H_f^\circ = -92$ kcal mol⁻¹ and $S^\circ = -49$ cal K⁻¹ mol⁻¹ for the equivalent $CrCl^{2+}(aq)$, inner sphere).


The work of Morris and Hammond¹²³ leads to K ($\mu = 1.0$ M) = 0.9 for formation of the outer-sphere complex represented by



A somewhat smaller equilibrium quotient was found by Gates


and King¹²² for solutions having higher ionic strength. We use the former result¹²³ to obtain $\Delta G_f^\circ = -417.9$ kcal mol⁻¹ for Cr(H₂O)₆Cl²⁺(aq) and also $\Delta G_f^\circ = -77.8$ kcal mol⁻¹ for the equivalent CrCl²⁺(aq, outer sphere).

From investigations cited by Sillén^{75,112} we adopt $K = 0.1$ for



This K leads to $\Delta G_f^\circ \approx -336.3$ kcal mol⁻¹ for Cr(H₂O)₄Cl₂⁺(aq) and also to $\Delta G_f^\circ \approx -109.6$ kcal mol⁻¹ for the equivalent CrCl₂⁺(aq, inner sphere). Measurements by King et al.^{117,122} lead to $\Delta H^\circ = 5$ kcal mol⁻¹ for this reaction and thence to $\Delta H_f^\circ \approx -401$ kcal mol⁻¹ for Cr(H₂O)₄Cl₂⁺(aq) and to $\Delta H_f^\circ \approx -127$ kcal mol⁻¹ for the equivalent CrCl₂⁺(aq, inner sphere). Combination of enthalpies and free energies leads to $S^\circ \approx 52$ cal K⁻¹ mol⁻¹ for Cr(H₂O)₄Cl₂⁺(aq) and to $S^\circ \approx -15$ cal K⁻¹ mol⁻¹ for CrCl₂⁺(aq, inner sphere).


Results of several investigators¹²⁴ are in agreement with $K = 2 \times 10^{-6}$ for



We calculate $\Delta G_f^\circ \approx -328.5$ kcal mol⁻¹ for Cr(H₂O)₃(OH)Cl₂(aq) and $\Delta G_f^\circ \approx -158.4$ kcal mol⁻¹ for the equivalent Cr(OH)Cl₂(aq, inner sphere).


The Cr(H₂O)₄Cl₂⁺(aq) species can exist in cis and trans forms. Two investigations¹²⁵ are consistent with $K = 0.5$ for cis = trans and the conclusion that the cis form is 0.4 kcal mol⁻¹ more stable than the trans form.

Equilibrium investigations by Espenson and King¹²⁶ are consistent with $K = 2 \times 10^{-3}$ and $\Delta H^\circ = 5$ kcal mol⁻¹ for




We calculate $\Delta G_f^\circ \approx -351.1$ kcal mol⁻¹, $\Delta H_f^\circ \approx -422.7$ kcal mol⁻¹, and $S^\circ \approx 41.5$ cal K⁻¹ mol⁻¹ for Cr(H₂O)₅Br²⁺(aq), and also obtain $\Delta G_f^\circ \approx -67.7$ kcal mol⁻¹, $\Delta H_f^\circ \approx -81.1$ kcal mol⁻¹, and $S^\circ \approx -42$ cal K⁻¹ mol⁻¹ for the equivalent CrBr²⁺(aq, inner sphere). Old calorimetric results from Recoura⁴⁵ lead to $\Delta H_f^\circ = -377.1$ kcal mol⁻¹ for Cr(H₂O)₄Br₂⁺(aq) and to $\Delta H_f^\circ \approx -103.8$ kcal mol⁻¹ for the equivalent CrBr₂⁺(aq, inner sphere).

Results of Spreer and King¹²⁷ lead to $K(\mu = 4.1) = 0.2$ for



From Swaddle and Guastalla¹²⁸ we have equilibrium quotients at several temperatures for replacement of water in Cr(H₂O)₆³⁺(aq) by iodide ion; at 25 °C they give $K(\mu = 4.2) = 7 \times 10^{-5}$.

Espenson and Binau¹²⁹ have reported $K(\mu = 1.0)$ values from 45–65 °C for




We use their derived enthalpy and entropy to calculate $K(\mu = 1.0) = 17$ at 25 °C.


Two investigations¹³⁰ have provided enthalpy data for cis-trans transformations for [CrCl₂(NH₃)₄]Cl·H₂O(c) and [CrBr(H₂O)(NH₃)₄]Br₂(c).

Jones and Bjerrum¹³¹ have made measurements on concentrated solutions with results that lead to estimates of the stability of the inner-sphere complex Cr(H₂O)₅ClO₄²⁺(aq).

Results of Mercer and Hormuth¹³² lead to $K(\mu = 0.5) = 1.3 \times 10^2$ for




Ardon and Sutin¹³³ have reported $K(\mu = 1.0) = 1 \times 10^{-2}$ for



Their K values at several temperatures also lead to enthalpies and entropies.


Results of two investigations¹³⁴ of aqueous chromic sulfate complexes are in poor agreement with each other, although it does appear to be well established that there is a kinetically inert inner-sphere complex and a labile outer-sphere complex.

Measurements by Hume and Kolthoff¹³⁵ have led to the following:



Guzzetta and Hadley¹³⁶ have reported calorimetric data for formation of Cr(CN)₆⁴⁻(aq), but their results do not justify calculation of a ΔH_f° value.

Poulsen, Bjerrum, and Poulsen¹³⁷ and Postmus and King¹¹⁶ have investigated the chromic thiocyanate complex at several temperatures and ionic strengths. From the latter investigation¹¹⁶ we take the following:




$$K = 1.2 \times 10^3 \quad \Delta H^\circ = -2.13 \text{ kcal mol}^{-1}$$


$$\Delta S^\circ = 6.9 \text{ cal K}^{-1} \text{ mol}^{-1}$$

We use these values in calculating $\Delta G_f^\circ \approx -312.0$ kcal mol⁻¹, $\Delta H_f^\circ \approx -382.5$ kcal mol⁻¹, and $S^\circ \approx 49$ cal K⁻¹ mol⁻¹ for Cr(H₂O)₅SCN²⁺(aq), and also $\Delta G_f^\circ \approx -28.6$ kcal mol⁻¹, $\Delta H_f^\circ \approx -40.9$ kcal mol⁻¹, and $S^\circ \approx -35$ cal K⁻¹ mol⁻¹ for the equivalent Cr(SCN)₂⁺(aq, inner sphere).

Templeton and King¹³⁸ have investigated the protonation of the chromic azide complex and report the following (30 °C):



Equilibrium¹³⁹ and calorimetric¹⁴⁰ measurements have led to the following:



$$K = 1 \times 10^{-5} \quad \Delta H^\circ = 8.3 \text{ kcal mol}^{-1}$$

$$\Delta S^\circ = 5.5 \text{ cal K}^{-1} \text{ mol}^{-1}$$

Duffy and Earley¹⁴¹ have made extensive measurements on the replacement of H₂O by thiocyanate and chloride in Cr(NH₃)₅(H₂O)³⁺(aq) and have also investigated outer-sphere complex or ion-pair species involving this cation with thiocyanate and chloride. They have reported several K , ΔH° , and ΔS° values for these reactions. Guastalla and Swaddle¹⁴² have carried out both equilibrium and rate measurements on the aquation of chromic ammine halides and have obtained ΔV° and ΔV^\ddagger values for these reactions.

Equilibrium constants for formation of such species as Cr(NCS)Hg⁴⁺(aq), Cr(NC)Hg⁴⁺(aq), and {[Cr(NH₃)₅(NCS)]-Ag}³⁺(aq) have been reported.¹⁴³

DTA measurements leading¹⁴⁴ to ΔH values for deaquation of such compounds as [Cr(NH₃)₅(H₂O)]X₃(c) and K₃[Cr(C₂O₄)₃]·3H₂O(c) are generally in poor agreement with other measurements on the same compounds. Some reported ΔH values for reactions where the products are H₂O(liq) and H₂O(g), respectively, are inconsistent with the well-known enthalpy of vaporization of water.

For the spinel FeCr₂O₄(c) the NBS 270-4 lists $S^\circ = 34.9$ cal K⁻¹ mol⁻¹, which is the same as the value derived from results previously cited by Kelley and King.²² The $\Delta H_f^\circ = -345.3$ kcal mol⁻¹ and $\Delta G_f^\circ = -321.2$ kcal mol⁻¹ given in NBS 270-4 for this compound are consistent with the high-temperature emf and equilibrium results obtained by Tretjakow and Schmalzried,³⁹ Boericke and Bangert,¹⁴⁵ Chen and Chipman,¹⁴⁶ and Katsura and Muan.¹⁴⁷ We therefore adopt

the ΔH_f° and ΔG_f° values listed in NBS 270-4, rather than the properties derived from the high-temperature results of Novokhatskii and Lenev.³⁶

The NBS 270-6 lists $S^\circ = 25.34 \text{ cal K}^{-1} \text{ mol}^{-1}$ for the spinel $\text{MgCr}_2\text{O}_4(\text{c})$, in agreement with the value derived from results cited by Kelley and King.²² The $\Delta H_f^\circ = -426.3 \text{ kcal mol}^{-1}$ that is listed in NBS 270-6 for this compound is in good agreement with the value we calculate from the high-temperature emf results of Tretjakow and Schmalzried³⁹ [with $(H_T - H_{298})$ values from Kelley³⁷]. A slightly different and probably better $\Delta H_f^\circ = -426.9 \text{ kcal mol}^{-1}$ can be similarly calculated from the recent high-temperature (900 °C) calorimetric results of Müller and Kleppa.¹⁴⁸ We combine this latter ΔH_f° with the S° above to obtain $\Delta G_f^\circ = -399.5 \text{ kcal mol}^{-1}$ for $\text{MgCr}_2\text{O}_4(\text{c})$.

Calorimetric measurements by Müller and Kleppa¹⁴⁸ lead to ΔH° of formation of $\text{NiCr}_2\text{O}_4(\text{c})$ from the constituent oxides at 900 °C. Combination of this value with our estimated $(H_T - H_{298})$ leads to $\Delta H_f^\circ = -330.6 \text{ kcal mol}^{-1}$ for $\text{NiCr}_2\text{O}_4(\text{c})$ at 298 K. This value appears to be preferable to that derived from the temperature derivative of the emf results of Tretjakow and Schmalzried.³⁹ Although the ΔH° based on these latter³⁹ results appears to have a large uncertainty, the uncertainty associated with ΔG° at any temperature is less large and can be used with estimated $(H_T - H_{298})$ and $(S_T - S_{298})$ values and the calorimetric ΔH° cited above to obtain ΔG_f° and S° at 298 K. Or we can directly estimate S° at 298 K and combine with the above ΔH_f° to obtain ΔG_f° at 298 K. On the basis of these two approaches, we adopt $S^\circ = (31.5) \text{ cal K}^{-1} \text{ mol}^{-1}$ and $\Delta G_f^\circ = -305.3 \text{ kcal mol}^{-1}$ for $\text{NiCr}_2\text{O}_4(\text{c})$.

Calorimetric measurements by Müller and Kleppa¹⁴⁸ have led to ΔH° values for formation of $\text{ZnCr}_2\text{O}_4(\text{c})$ and $\text{CdCr}_2\text{O}_4(\text{c})$ from their constituent oxides at 900 °C. Combination of these values with estimated $(H_T - H_{298})$ values leads to $\Delta H_f^\circ = -370.9 \text{ kcal mol}^{-1}$ for $\text{ZnCr}_2\text{O}_4(\text{c})$ and $\Delta H_f^\circ = -344.3 \text{ kcal mol}^{-1}$ for $\text{CdCr}_2\text{O}_4(\text{c})$, both at 298 K. We estimate $S^\circ = (30.5) \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{ZnCr}_2\text{O}_4(\text{c})$ and calculate its $\Delta G_f^\circ = -344.4 \text{ kcal mol}^{-1}$. We also estimate $S^\circ = (34) \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{CdCr}_2\text{O}_4(\text{c})$ and calculate its $\Delta G_f^\circ = -318.1 \text{ kcal mol}^{-1}$.

Calorimetric results from Müller and Kleppa¹⁴⁸ lead to ΔH° for formation of $\text{CuCr}_2\text{O}_4(\text{c})$ from the oxides at 900 °C. We combine this result with our estimated $(H_T - H_{298})$ to obtain $\Delta H_f^\circ = -307.3 \text{ kcal mol}^{-1}$ at 298 K. We adopt this value rather than that derived from the temperature derivative of the high-temperature results of Schmahl and Minzl.¹⁴⁹ We proceed as described above for $\text{NiCr}_2\text{O}_4(\text{c})$ to obtain $S^\circ = (33) \text{ cal K}^{-1} \text{ mol}^{-1}$ and $\Delta G_f^\circ = -282.2 \text{ kcal mol}^{-1}$ for $\text{CuCr}_2\text{O}_4(\text{c})$.

Tretjakow and Schmalzried³⁹ and also Aukrust and Muan¹⁵⁰ have made measurements leading to apparently reliable ΔG° values (in satisfactory agreement with each other) for formation of $\text{CoCr}_2\text{O}_4(\text{c})$ at high temperatures. Because there are substantial uncertainties in the derived ΔH° values and our calculation of thermodynamic properties at 298 K also requires estimated C_p or related properties, we omit this compound from our Table I.

Gross, Wilson, and Gutteridge¹¹⁰ have made calorimetric measurements leading to ΔH° of formation of $\text{NaCrO}_2(\text{c})$ from $\text{Na}_2\text{O}(\text{c})$ and $\text{Cr}_2\text{O}_3(\text{c})$ by way of somewhat uncertain reactions. We use their ΔH° value for this reaction with $\Delta H_f^\circ = -99.9 \text{ kcal mol}^{-1}$ for $\text{Na}_2\text{O}(\text{c})$ from the JANAF tables¹⁵¹ with ΔH_f° for $\text{Cr}_2\text{O}_3(\text{c})$ from Table I to obtain $\Delta H_f^\circ = -210 \text{ kcal mol}^{-1}$ for $\text{NaCrO}_2(\text{c})$.

Zalazinskii et al.¹⁵² have carried out high-temperature equilibrium measurements on the reduction of "cuprous chromite" of formula $\text{CuCrO}_2(\text{c})$. Our calculations with their results lead to $S^\circ \approx 6 \text{ cal K}^{-1} \text{ mol}^{-1}$ for this compound. Be-

cause this unreasonably small entropy value indicates either error or very large uncertainty, we do not list $\text{CuCrO}_2(\text{c})$ in our Table I.

E. Cr(IV) and Cr(V): Compounds and Aqueous Ions




The NBS 270-4 lists $\Delta H_f^\circ = -298 \text{ kcal mol}^{-1}$ for $\text{CrF}_4(\text{c})$, which is in agreement with the results of Wartenberg.¹⁵³ We estimate $S^\circ = (33) \text{ cal K}^{-1} \text{ mol}^{-1}$ and calculate $\Delta G_f^\circ = -277 \text{ kcal mol}^{-1}$ for $\text{CrF}_4(\text{c})$.

The results of several investigators^{91,154,155} are in reasonable agreement with the NBS 270-4 $\Delta H_f^\circ = -102 \text{ kcal mol}^{-1}$ (maybe a slightly more negative value would be better) for $\text{CrCl}_4(\text{g})$. We combine this ΔH_f° with $S^\circ = 87.5 \text{ cal K}^{-1} \text{ mol}^{-1}$ from Maier⁹¹ to obtain $\Delta G_f^\circ = -97.5 \text{ kcal mol}^{-1}$ for $\text{CrCl}_4(\text{g})$.






We can use the somewhat uncertain results of Sime and Gregory¹⁵⁶ for reaction of $\text{CrBr}_3(\text{c})$ with $\text{Br}_2(\text{g})$ with the more uncertain ΔH_f° of $\text{CrBr}_3(\text{c})$ discussed in the preceding section to obtain a ΔH_f° value for $\text{CrBr}_4(\text{g})$ that seems far too negative in comparison to ΔH_f° values for $\text{CrCl}_4(\text{g})$ and $\text{CrI}_4(\text{g})$. Shieh and Gregory¹⁰² have estimated $\Delta H_f^\circ = (-42.6) \text{ kcal mol}^{-1}$ for $\text{CrBr}_4(\text{g})$, a value which seems more reasonable to us than the value we have calculated as described above. The S° has been estimated as (100.1) and (106.9) $\text{cal K}^{-1} \text{ mol}^{-1}$. Because of all the uncertainties, we omit $\text{CrBr}_4(\text{g})$ from our Table I.

The results of Shieh and Gregory¹⁰² in combination with our ΔH_f° for $\text{CrI}_3(\text{c})$ leads to $\Delta H_f^\circ = 3 \text{ kcal mol}^{-1}$ for $\text{CrI}_4(\text{g})$. Combination of this ΔH_f° with $S^\circ = 112 \text{ cal K}^{-1} \text{ mol}^{-1}$ from Shieh and Gregory¹⁰² gives $\Delta G_f^\circ = -12 \text{ kcal mol}^{-1}$ for $\text{CrI}_4(\text{g})$.

We now consider approximate potentials for various couples involving aqueous Cr(IV) and Cr(V), beginning with alkaline solutions. Bailey and Symons¹⁸ have deduced an approximate ΔG_f° for the Cr(V) species they represent by $\text{CrO}_4^{3-}(\text{aq})$. Revision of their value to make it consistent with our ΔG_f° for $\text{CrO}_4^{2-}(\text{aq})$ leads to $\Delta G_f^\circ = -176.3 \text{ kcal mol}^{-1}$ for $\text{CrO}_4^{3-}(\text{aq})$ and thence the following potentials:



Kinetic evidence¹⁵⁷ leads to generally consistent (approximate) potentials for various half reactions involving Cr(IV) and Cr(V) in acidic solutions. Another approach is to assume that the Cr(V) species in acidic solution is $\text{H}_3\text{CrO}_4(\text{aq})$ and then calculate its free energy from the ΔG_f° above for $\text{CrO}_4^{3-}(\text{aq})$ and a ΔG° of protonation estimated by way of the known ΔG° of protonation of $\text{PO}_4^{3-}(\text{aq})$. The following potentials (for acidic solutions) are consistent with all of this.




These potentials (acidic solution) are conveniently summarized in the following potential diagram.

TABLE III. ΔH_f° (298 K) Values for Chromium Carbides

Carbide	$\Delta H_f^\circ/\text{kcal mol}^{-1}$							
	NBS 270-4	Mah ^a	Mah ^b	Kulkarni ^c	Kulkarni ^d	Mabuchi ^e	Kleykamp ^f	Storms ^g
Cr ₃ C ₂ (c)	-19.3	-22.5	-22.2			-16.75	-13.5	-13.6 to -29.4
Cr ₇ C ₃ (c)	-38.7	-43.3	-42.6	-36.7	-34.3		-26.8	
Cr ₂₃ C ₆ (c)	-87.2	-94.7	-92.4	-81.9	-77.1			-69, -81.9

^a Reference 35 (combustion calorimetry). Published values quoted here are based on $\Delta H_f^\circ = -272.6 \text{ kcal mol}^{-1}$ for Cr₂O₃(c). ^b Results of ref 35 recalculated on the basis of our adopted $\Delta H_f^\circ = -272.4 \text{ kcal mol}^{-1}$ for Cr₂O₃(c). Similar recalculation on the basis of $\Delta H_f^\circ = -270.0 \text{ kcal mol}^{-1}$ for Cr₂O₃(c) as suggested by Kulkarni and Worrell⁴² leads to ΔH_f° values that are 1.2 kcal mol⁻¹ of Cr less negative than values given here. ^c Reference 42 (high-temperature equilibrium). Published values quoted here are based on their selected $\Delta H_f^\circ = -270.0 \text{ kcal mol}^{-1}$ for Cr₂O₃(c). ^d Results of ref 42 recalculated to conform with our tabulated $\Delta H_f^\circ = -272.4 \text{ kcal mol}^{-1}$ for Cr₂O₃(c). ^e Reference 161 (high-temperature emf). This value does not depend on ΔH_f° of Cr₂O₃(c). ^f Reference 162 (high-temperature emf). These values do not depend on ΔH_f° of Cr₂O₃(c). ^g Reference 27 (review of earlier works from 1926 to 1965). Some of these values are independent of properties assigned to Cr₂O₃(c).



F. Other Inorganic Compounds of Chromium

Wolf¹⁵⁸ has measured heat capacities of CrH_x(c) ($x = 0.84, 0.91, \text{ and } 0.94$) and has evaluated entropies at 298 K as listed in Table I. For the compound or mixture of phases with composition described by Cr₇H₂(c) we adopt $\Delta H_f^\circ = -3.8 \text{ kcal mol}^{-1}$ as listed in NBS 270-4 and as consistent with the published results of Sieverts and Gotta.¹⁵⁹ We also note that Stock and Hardcastle¹⁶⁰ have carried out phase and composition analysis of the chromium hydride system.

The chromium carbides are of considerable importance in metallurgy and have been investigated several times^{27,35,42,161,162} by calorimetric, emf, and equilibrium methods. Results in the form of ΔH_f° values at 298 K are summarized in Table III. The wide range in reported ΔH_f° values cannot be resolved by our data analysis; still more measurements are required. The ΔH_f° values listed in the NBS 270-4 appear to be reasonable choices and are adopted for our Table I. Entropies listed in NBS 270-4 are all consistent with results summarized by Kelley and King.²² We also have $\Delta H_f^\circ \approx 184 \text{ kcal mol}^{-1}$ for CrC₂(g) from the work of Kohl and Stearns.¹⁶³

The NBS 270-4 lists $\Delta H_f^\circ = -30.5 \text{ kcal mol}^{-1}$ for Cr₂N(c) and $\Delta H_f^\circ = -29.8 \text{ kcal mol}^{-1}$ for CrN(c). This value for CrN(c) is consistent with results cited previously by Bichowsky and Rossini⁴⁵ and in NBS Circular 500.⁷³ The value above for Cr₂N(c) is identical with the value derived from Mah's¹⁶⁴ combustion calorimetry and the $\Delta H_f^\circ = -270.4 \text{ kcal mol}^{-1}$ for Cr₂O₃(c) that has been adopted here and in NBS 270-4. Had we adopted $\Delta H_f^\circ = -270.0 \text{ kcal mol}^{-1}$ for Cr₂O₃(c) as suggested by other considerations,⁴² we would obtain $\Delta H_f^\circ = -28.1 \text{ kcal mol}^{-1}$ for Cr₂N(c) from the combustion results. Results of various other calorimetric and high-temperature equilibrium studies^{45,73,165} lead to ΔH_f° values for CrN(c) and Cr₂N(c) that are equal to or less negative than the values listed in NBS 270-4. On the basis of all of these results, we select $\Delta H_f^\circ = -29 \text{ kcal mol}^{-1}$ for CrN(c) and $\Delta H_f^\circ = -30 \text{ kcal mol}^{-1}$ for Cr₂N(c). Combination of these values with $S^\circ = (6) \text{ cal K}^{-1} \text{ mol}^{-1}$ for CrN(c) and $S^\circ = (12) \text{ cal K}^{-1} \text{ mol}^{-1}$ for Cr₂N(c) leads to the tabulated ΔG_f° values. Although several investigations¹⁶⁵ have shown that the nitrides are nitrogen deficient with stoichiometries (under particular conditions) that lead to formulas such as Cr₂N_{0.977}(c) and CrN_{0.982}(c), we have followed earlier practice in using the simple formulas Cr₂N(c) and CrN(c).

Chart¹⁶⁶ has recently carried out a thorough review of the

thermodynamic properties of silicides of chromium (and other elements) and has selected "best" values for these properties. Our analysis of recent results^{24,167} cited by Chart¹⁶⁶ and earlier results cited in these recent papers leads to our selected (in Table I) ΔH_f° values that are identical with those selected by Chart¹⁶⁶ for all of the chromium silicides. All of these ΔH_f° values are less negative than values listed for the same compounds in NBS 270-4. Our selection of entropies is identical with that of Chart¹⁶⁶ except for a small difference for CrSi₂(c). Free energies (Table I) are consistent with the tabulated ΔH_f° and S° values.

Low-temperature C_p measurements by Kalishevich et al.¹⁶⁸ lead to the entropies listed in Table I for CrGe(c), Cr₅Ge₃(c), and Cr₁₁Ge₁₉(c).

The high-temperature equilibrium results of Hager and Elliott¹⁶⁹ lead to $\Delta H_f^\circ = -32 \text{ kcal mol}^{-1}$, $\Delta G_f^\circ = -33 \text{ kcal mol}^{-1}$, and $S^\circ = 17 \text{ cal K}^{-1} \text{ mol}^{-1}$ for CrS(c). For CrS(g) we have $\Delta H_f^\circ \approx 83 \text{ kcal mol}^{-1}$ from Drowart et al.¹⁷⁰ We estimate $S^\circ = (60) \text{ cal K}^{-1} \text{ mol}^{-1}$ and then obtain $\Delta G_f^\circ \approx 69 \text{ kcal}$ for CrS(g).

The entropies listed in NBS 270-4 and in our Table I for Cr₂Te₃(c), Cr₃Te₄(c), and Cr₅Te₆(c) are consistent with the C_p results of Grønwald and Westrum.¹⁷¹ We also note that Grønwald¹⁷² has reported high-temperature C_p and related properties for Cr₃Te₄(c).

The NBS 270-4 has listed $\Delta H_f^\circ = -257.4 \text{ kcal mol}^{-1}$ for Cr(CO)₆(c), which is exactly the same as the value we calculate from the combustion results of Cotton, Fischer, and Wilkinson¹⁷³ and $\Delta H_f^\circ = -272.4 \text{ kcal mol}^{-1}$ for Cr₂O₃(c) from NBS 270-4 and our Table I. More recently, Skinner and Virmani¹⁷⁴ have made calorimetric measurements on decomposition of Cr(CO)₆(c) to Cr(c) and CO(g) and have obtained $\Delta H_f^\circ = -223.5 \text{ kcal mol}^{-1}$ for Cr(CO)₆(c). Similar measurements by Connor, Skinner, and Virmani¹⁷⁵ have led to $\Delta H_f^\circ = -222.9 \text{ kcal mol}^{-1}$ for Cr(CO)₆(c). These latter workers have also made calorimetric measurements on the iodination of Cr(CO)₆(c). Combination of their results with ΔH_f° values already adopted here for CrI₂(c) and CrI₃(c) leads to $\Delta H_f^\circ = -234.3 \text{ kcal mol}^{-1}$ for Cr(CO)₆(c). In the absence of any clear indication as to which of the results cited here is best, we hesitantly adopt $\Delta H_f^\circ = -234 \text{ kcal mol}^{-1}$ (based on iodination measurements¹⁷⁵) for Cr(CO)₆(c).

The ΔH_f° values listed in NBS 270-4 correspond to $\Delta H^\circ = 17.0 \text{ kcal mol}^{-1}$ for sublimation of Cr(CO)₆(c). This value is consistent with vapor pressure results cited by Cotton et al.¹⁷³ and also with more recent results.¹⁷⁶ We use this value with the ΔH_f° above for Cr(CO)₆(c) to obtain $\Delta H_f^\circ = -217 \text{ kcal mol}^{-1}$ for Cr(CO)₆(g). Statistical calculations have led to $S^\circ = 116.5 \text{ cal K}^{-1} \text{ mol}^{-1}$ (ref 177) and to an apparently better $S^\circ = 111.6 \text{ cal K}^{-1} \text{ mol}^{-1}$ (ref 178) for Cr(CO)₆(g). We adopt the latter value and combine with the ΔH_f° of Cr(CO)₆(g) to obtain its $\Delta G_f^\circ = -202 \text{ kcal mol}^{-1}$.

The vapor pressure results cited above are also consistent with $\Delta S^\circ = 40 \text{ cal K}^{-1} \text{ mol}^{-1}$ for sublimation of $\text{Cr}(\text{CO})_6(\text{c})$ at 298 K. Combination of this ΔS° with our adopted S° for $\text{Cr}(\text{CO})_6(\text{g})$ leads to $S^\circ = 72 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{Cr}(\text{CO})_6(\text{c})$, for which a value based on C_p measurements would be useful. Combination of this value with the ΔH_f° adopted above leads to $\Delta G_f^\circ = -207 \text{ kcal mol}^{-1}$ for $\text{Cr}(\text{CO})_6(\text{c})$.

Calorimetric measurements by Brown, Connor, and Skinner¹⁷⁹ lead to $\Delta H_f^\circ = -1446.4 \text{ kcal mol}^{-1}$ for $\text{Cr}(\text{PF}_3)_6(\text{c})$, based on $\Delta H_f^\circ = -228.89 \text{ kcal mol}^{-1}$ for $\text{PF}_3(\text{g})$.

G. Organic Compounds and Complexes

Thermochemical properties of a few organochromium compounds have been reviewed by Cox and Pilcher¹⁸⁰ and also by Ashcroft and Mortimer.¹⁸¹ These latter authors¹⁸¹ and also Martell^{75,112} have reviewed stabilities and other thermodynamic properties of organic complexes of Cr(III) in aqueous solution. In our present review we limit discussion of organic compounds of chromium to brief mention of some recent investigations that have not been cited in these earlier reviews.^{75,112,180,181}

Before citing particular investigations, it is important to emphasize that comparisons of quantities reported in original papers or quantities listed in various reviews must be based on the same auxiliary data in order to be useful. For example, enthalpies of formation of organochromium compounds derived from combustion measurements must be based on the same ΔH_f° of formation of $\text{Cr}_2\text{O}_3(\text{c})$.

Connor, Skinner, and Virmani¹⁸² have made calorimetric measurements leading to ΔH_f° values for $\text{Cr}(\text{C}_6\text{H}_6)_2$, $\text{Cr}(\text{C}_6\text{H}_6)(\text{CO})_3$, $\text{Cr}[\text{C}_6(\text{CH}_3)_6](\text{CO})_3$, $\text{Cr}(\text{C}_6\text{H}_6)_2\text{l}$, $\text{Cr}(\text{C}_7\text{H}_8)(\text{CO})_3$, and $\text{Cr}(\text{C}_6\text{H}_5\text{CH}_3)(\text{CO})_3$.

Calorimetric measurements by Bradley and Hillyer¹⁸³ lead to ΔH_f° of $\text{Cr}(\text{OC}_4\text{H}_9)_4$.

Tel'noi et al.¹⁸⁴ have determined heats of combustion of $\text{Cr}(\text{C}_6\text{H}_6)_2$, $\text{Cr}(\text{C}_6\text{H}_6)_2\text{Cl}$, $\text{Cr}(\text{C}_6\text{H}_6)_2\text{Br}$, $\text{Cr}(\text{C}_6\text{H}_5\text{CH}_3)_2\text{l}$, $\text{Cr}[\text{C}_6\text{H}_4(\text{CH}_3)_2]_2\text{l}$, and $\text{Cr}(\text{C}_6\text{H}_5\text{C}_6\text{H}_5)_2\text{l}$.

Sorai, Tachiki, Suga, and Seki¹⁸⁵ have made C_p measurements on $[\text{Cr}_3\text{O}(\text{CH}_3\text{COO})_6(\text{H}_2\text{O})_3]\text{Cl}\cdot 6\text{H}_2\text{O}(\text{c})$ from 1.5 to 280 K. Sorai and Seki¹⁸⁶ have made C_p measurements from 1.4 to 200 K on $[\text{Cr}_4(\text{OH})_6(\text{en})_6](\text{SO}_4)_3\cdot 10\text{H}_2\text{O}(\text{c})$.

Andrews, Westrum, and Bjerrum¹⁸⁷ have determined C_p from 5 to 350 K and vapor pressures over a range of temperature for $\text{Cr}(\text{C}_6\text{H}_6)_2$. They have calculated S° values and the thermodynamics of vaporization.

Hill and Irving¹⁸⁸ have made solution calorimetric measurements leading to ΔH_f° of tris(acetylacetonato)chromium(III). Melia and Merrifield¹⁸⁹ have made C_p measurements on this compound from 80 to 300 K and also evaluated the enthalpy of vaporization. Fontaine, Pommier, and Guiochon¹⁹⁰ have reported enthalpies of vaporization of this compound and some derivative compounds.

III. Molybdenum and Tungsten

A. Descriptive Chemistry

Compounds and ions containing molybdenum or tungsten in oxidation states ranging from -2 to $+6$ are known. In general, compounds and ions containing Mo(VI) or W(VI) are more stable (weaker oxidizing agents) than corresponding Cr(VI) species. There is a much greater tendency for Mo(VI) and W(VI) toward formation of polynuclear species than there is for Cr(VI) for which such reactions ordinarily stop at dichromate. Similarly, the important species of lower oxidation states of molybdenum and tungsten are often more complex than those of chromium in the same oxidation state. In contrast to Cr(III), there are relatively few stable complexes of Mo(III) and W(III).

Cotton and Wilkinson²⁰ have recently summarized several aspects of the chemistry of molybdenum and tungsten, with particular emphasis on structures of various compounds and ions. Here we will focus on identification of aqueous species and qualitative statements about stabilities that will furnish background appropriate to further discussion of thermodynamic and electrochemical equilibria involving aqueous solutions.

Most molybdenum is obtained from molybdenite, MoS_2 , although some is obtained from various molybdates. Tungsten is found in nature chiefly as tungstates of iron, calcium, and a few other metals.

The oxides MoO_3 and WO_3 react at high temperatures with various metal oxides to form corresponding molybdates and tungstates. Both oxides dissolve (WO_3 only slowly at room temperature) in aqueous alkali to yield aqueous molybdate and tungstate ions represented by $\text{MoO}_4^{2-}(\text{aq})$ and $\text{WO}_4^{2-}(\text{aq})$. These ions have been shown to be tetrahedral (slightly distorted in some compounds) in the solid state²⁰ and also in aqueous solution.¹⁹¹ Normal solid molybdates and tungstates can be crystallized from neutral or alkaline solutions. Molybdates and tungstates of the alkali metals and ammonium ion are quite soluble in water, while those of most other metals are only slightly soluble.

The solid isopoly compounds that may be obtained from acidified solutions of molybdates and tungstates are varied and complicated. Structures of several have been determined.²⁰ Examples of isopoly ions that are well characterized in the solid state are the "paramolybdate" $\text{Mo}_7\text{O}_{24}^{6-}$ and the "octamolybdate" $\text{Mo}_8\text{O}_{26}^{4-}$.

When alkaline solutions containing $\text{MoO}_4^{2-}(\text{aq})$ or $\text{WO}_4^{2-}(\text{aq})$ are acidified under such conditions that no precipitate forms, various condensation reactions take place to yield species that are still inadequately characterized after a large number of investigations employing a variety of experimental methods. Among the aqueous isopoly species that have been postulated are $\text{Mo}_3\text{O}_{11}^{4-}$, $\text{HMo}_3\text{O}_{11}^{3-}$, $\text{Mo}_4\text{O}_{13}^{2-}$, $\text{H}_n\text{Mo}_6\text{O}_{20}^{-4+n}$, $\text{H}_n\text{Mo}_6\text{O}_{21}^{-6+n}$, various ions related to paramolybdate and octamolybdate above, and still larger ions involving 12 and 24 Mo with various numbers of hydrogen and oxygen atoms per ion. Because of uncertain identifications of the various aqueous species and the absence of activity coefficients for these highly charged ions, all numerical values of equilibrium "constants" or "quotients" involving isopoly species are open to question.

Acidification of molybdate or tungstate solutions can also lead to precipitation of $\text{MO}_3\cdot n\text{H}_2\text{O}$ compounds that are commonly called molybdic acid and tungstic acid.

Heteropoly ions and solids are obtained when molybdate and tungstate solutions that also contain other oxo ions (such as PO_4^{3-}) or certain metal ions are acidified. The acids and most salts of heteropoly anions are quite soluble in water and also in many polar organic solvents. Structures and formulas of a number of heteropoly anions have been reviewed by Cotton and Wilkinson.²⁰ Results of some structural studies have proven useful in connection with studies of aqueous solutions cited later.

Connor and Ebsworth¹⁹ have reviewed the chemistry of peroxy compounds of molybdenum and tungsten, all apparently derived from the $+6$ oxidation state. Salts prepared by action of excess H_2O_2 on normal molybdates appear to contain four peroxy groups per atom of molybdenum. It seems likely that the $\text{Mo}(\text{O}_2)_4^{2-}$ ion exists in the solid state and that either this ion or its hydrolysis products exist in aqueous solutions. Many compounds containing fewer than four peroxy groups per Mo (or W) have been reported.

The sulfides MoS_3 and WS_3 , various thiomolybdates, and thiotungstates can be precipitated from aqueous solutions. Although these compounds are important in analytical chem-

istry, solid state science, and heterogeneous catalysis, we have few quantitative data to cite later in this review.

There are several hexahalides and oxyhalides of formulas MX_6 , MOX_4 , and MO_2X_2 . There is also the volatile compound that has been described as $MoO_3 \cdot 2HCl$ or $MoO_2Cl_2 \cdot H_2O$ or (probably most realistically) $MoO(OH)_2Cl_2$.

Reduction of Mo(VI) and W(VI) compounds or aqueous ions can lead to a large number of compounds or aqueous species, many of which have been described by Cotton and Wilkinson.²⁰ We shall later cite thermodynamic data for some of these compounds (especially binary compounds with oxygen, sulfur, and halogens) and also potentials that relate various aqueous species.

High-temperature reduction of alkali metal tungstates with hydrogen or reaction of alkali metals with WO_3 leads to "tungsten bronzes" that have semimetallic properties. These substances are mostly unreactive toward water and aqueous solutions, but some are readily oxidized by $Ag^+(aq)$ and other oxidizing agents in acidic solution and by O_2 in alkaline solution. Similar "bronzes" of molybdenum and other metals are less well known, but do exist.

The interest in complexes formed between Mo(VI) [also W(VI)] and organic ligands is partly due to the fact that molybdenum is an essential element in various biological systems. Of particular interest is the role of molybdenum in the fixation of nitrogen by some bacteria commonly associated with legumes. Cotton and Wilkinson²⁰ have provided a useful introduction with leading references to this important subject.

B. Molybdenum, Molybdenum Oxides, and Molybdic Acid

As for chromium, the best general source of thermodynamic data for molybdenum and its compounds and ions is NBS 270-4. From this source we have $\Delta H_f^\circ = 0$, $\Delta G_f^\circ = 0$, and $S^\circ = 6.85 \text{ cal K}^{-1} \text{ mol}^{-1}$ for Mo(c) as listed in our Table IV. This entropy is in good agreement with the value derived by Kelley and King²² from C_p data and is in still better agreement with the value in the JANAF tables¹⁵¹ calculated from the same experimental results.

For Mo(g) we have $S^\circ = 43.461 \text{ cal K}^{-1} \text{ mol}^{-1}$ in NBS 270-4, a value in good agreement with earlier statistical calculations.^{22,151} The $\Delta H_f^\circ = 157.3 \text{ kcal mol}^{-1}$ in NBS 270-4 for Mo(g) is the same as the value in the JANAF tables,¹⁵¹ based on high temperature vapor pressure data. This value, which we adopt for our Table IV, is also in satisfactory agreement with the more recent results of Koch and Anable.¹⁹² Combination of ΔH_f° and S° leads to the tabulated $\Delta G_f^\circ = 146.4 \text{ kcal mol}^{-1}$ for Mo(g).

Earlier investigations of the high-temperature thermodynamic properties of Mo(c) and Mo(liq) have been reviewed in the JANAF tables.¹⁵¹ Here we call attention to four recent investigations¹⁹³ of C_p , $(H^\circ_T - H^\circ_{298})$, and related thermodynamic properties up to nearly 3000 K.

Now we consider the oxides of molybdenum, progressing from MoO to MoO_3 .

The NBS 270-4 lists $\Delta H_f^\circ = 101 \text{ kcal mol}^{-1}$ for MoO(g). Our calculations with the results of DeMaria et al.¹⁹⁴ are in agreement with those summarized in the JANAF tables,¹⁵¹ leading to our adopted $\Delta H_f^\circ \approx 93 \text{ kcal mol}^{-1}$ as listed in Table IV. The uncertainty in this value is large enough to overlap the value listed in NBS 270-4. We take $S^\circ = (57) \text{ cal K}^{-1} \text{ mol}^{-1}$ from the JANAF tables¹⁵¹ and calculate $\Delta G_f^\circ \approx 85 \text{ kcal mol}^{-1}$ for MoO(g).

Combustion calorimetry measurements by Staskiewicz, Tucker, and Snyder¹⁹⁵ and also by Mah¹⁹⁶ have led to $\Delta H_f^\circ = -140.86$ and $-140.64 \text{ kcal mol}^{-1}$ for $MoO_2(c)$. We adopt a mean value, $\Delta H_f^\circ = -140.76 \text{ kcal mol}^{-1}$, which is also the value listed in NBS 270-4. The JANAF tables¹⁵¹ list ΔH_f°

$= -140.5 \text{ kcal mol}^{-1}$, based on these results^{195,196} and others that we consider to have sufficiently large uncertainties that it is now better to adopt $\Delta H_f^\circ = -140.76 \text{ kcal mol}^{-1}$ as given above. We also note that some recent high-temperature emf results¹⁹⁷ lead to a calculated $\Delta H_f^\circ = -139.6 \text{ kcal mol}^{-1}$ for this compound.

We adopt $S^\circ = 11.06 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $MoO_2(c)$ as listed in NBS 270-4 and also by Kelley and King,²² based on C_p values down to 50 K and an extrapolation to absolute zero. It has been pointed out¹⁵¹ that there is a possibility of "extra" entropy that might justify a higher value of $S^\circ = 11.95 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $MoO_2(c)$. In the absence of needed C_p measurements at $T < 50 \text{ K}$, we continue to use the first S° cited above.

The NBS 270-4 lists $\Delta H_f^\circ = 3 \text{ kcal mol}^{-1}$ for $MoO_2(g)$, which is consistent with the results of two investigations.^{194,198} We also adopt $S^\circ = (66) \text{ cal K}^{-1} \text{ mol}^{-1}$ from the JANAF tables¹⁵¹ and calculate $\Delta G_f^\circ \approx 0 \text{ kcal mol}^{-1}$ for $MoO_2(g)$.

Syntheses, heat capacities, and emf results have been reported¹⁹⁹ for solids with compositions MoO_n ($2 < n < 3$).

Combustion calorimetry measurements by Staskiewicz, Tucker, and Snyder¹⁹⁵ and by Mah¹⁹⁶ have led to $\Delta H_f^\circ = -178.01$ and $-178.16 \text{ kcal mol}^{-1}$ for $MoO_3(c)$. We adopt the mean value, $\Delta H_f^\circ = -178.08 \text{ kcal mol}^{-1}$, which has also been adopted in NBS 270-4. Heat capacity measurements cited by Kelley and King²² lead to $S^\circ = 18.58 \text{ cal K}^{-1} \text{ mol}^{-1}$ and thence by combination with the ΔH_f° to $\Delta G_f^\circ = -159.66 \text{ kcal mol}^{-1}$ as also listed in NBS 270-4. Some high-temperature properties of $MoO_3(c)$ and $MoO_3(liq)$ have been reported.^{199,200}

We note that Kirshenbaum and Beardell²⁰¹ have made DTA measurements leading to ΔH values for various reactions involving MoO_2 , MoO_3 , and Mo with other metals and their oxides. These reported ΔH values are in generally poor agreement with values calculated from ΔH_f° values. The calculated quantities appear to be preferable.

Several investigators^{194,198,202-204} have made high-temperature measurements leading to data for $(MoO_3)_n(g)$ species. On the basis of their results, we adopt the uncertain thermodynamic properties listed in Table IV.

The NBS 270-4 has listed $\Delta H_f^\circ = -250.0 \text{ kcal mol}^{-1}$ for $H_2MoO_4(c)$, which is sometimes called "molybdic acid" and has also been represented as $MoO_3 \cdot H_2O(c)$. This value, which we adopt, is exactly consistent with the calorimetric results of Graham and Hepler.²⁰⁵ We estimate $S^\circ = (29) \text{ cal K}^{-1} \text{ mol}^{-1}$ and combine with the ΔH_f° to obtain $\Delta G_f^\circ = -218.1 \text{ kcal mol}^{-1}$ for $H_2MoO_4(c)$. This free energy is consistent with the experience that $H_2MoO_4(c)$ is stable with respect to $MoO_3(c)$ and $H_2O(liq)$ near 298 K. We also use the ΔG_f° to calculate that the equilibrium vapor pressure of $H_2O(g)$ over $H_2MoO_4(c)$ and $MoO_3(c)$ is $\sim 1 \text{ mmHg}$. In the absence²⁰⁵ of reliable vapor pressure data, all we can say is that this calculated value seems "reasonable".

As previously discussed,²⁰⁵ the results of Pechard and of Hüttig and Kurre, who may have worked with " $MoO_3 \cdot 2H_2O$ " or " $H_2MoO_4 \cdot H_2O$ ", are not sufficiently reliable to justify tabulating properties for this substance.

The NBS 270-4 lists $\Delta H_f^\circ = -212 \text{ kcal mol}^{-1}$ for $H_2MoO_4(g)$, while the JANAF tables¹⁵¹ list $\Delta H_f^\circ = -203.4 \text{ kcal mol}^{-1}$. Our assessment of the experimental results (cited in the JANAF tables¹⁵¹) leads us to adopt $\Delta H_f^\circ = -203 \text{ kcal mol}^{-1}$ and combine with an estimated¹⁵¹ $S^\circ = (85) \text{ cal K}^{-1} \text{ mol}^{-1}$ to obtain $\Delta G_f^\circ = -188 \text{ kcal mol}^{-1}$ for $H_2MoO_4(g)$.

C. $MoO_4^{2-}(aq)$ and Molybdates

Calorimetric measurements of the enthalpy of reaction of

TABLE IV. ^a Thermodynamic Properties of Molybdenum Compounds at 298 K


Compound	ΔH_f° kcal mol ⁻¹	ΔG_f° kcal mol ⁻¹	S° cal K ⁻¹ mol ⁻¹	Compound	ΔH_f° kcal mol ⁻¹	ΔG_f° kcal mol ⁻¹	S° cal K ⁻¹ mol ⁻¹
Mo(c)	0	0	6.85 ^{22,151}	MoF ₆ (liq)	-378.92 ^{237,240,241}	-352.05	62.06 ^{237,240}
Mo(g)	157.3 ^{151,192}	146.4	43.46 ^{122,151}	MoF ₆ (g)	-372.29 ²⁴¹	-351.88	83.75 ^{237,238}
Mo ⁺ (g)	322.5			MoOF ₄ (c)	-329.8 ²⁴⁴		
Mo ²⁺ (g)	696.5			MoO ₂ F ₂ (g)	-268 ²⁴⁸		
Mo ³⁺ (g)	1323.8			MoCl ₂ (c)	-68 ²⁴⁹	-59	(30) ²⁴⁹
Mo ⁴⁺ (g)	2395.2			MoCl ₃ (c)	-96.7 ^{249,251}	-83	(41) ²⁴⁹
Mo ⁵⁺ (g)	3807.2			MoCl ₄ (c)	-114 ²⁴⁹	-96	(53) ¹⁵¹
Mo ⁶⁺ (g)	5378			MoCl ₄ (g)	-92 ^{249,250}	-85	(89) ¹⁵¹
Mo ⁷⁺ (g)	8298			MoCl ₅ (c)	-126 ²⁴⁹	-101	(57) ¹⁵¹
Mo ⁸⁺ (g)	11 830			MoCl ₅ (g)	-107 ^{151,249}	-94	(95) ¹⁵¹
MoO(g)	~93 ^{151,194}	~85	(57) ¹⁵¹	MoOCl ₂ (c)	-126.2 ²⁵⁰	-110	(31) ²⁵⁰
MoO ₂ (c)	-140.76 ^{195,196}	-127.40	11.06 ²²	MoOCl ₃ (c)	-150.7 ²⁵¹	-130	(42) ²⁵¹
MoO ₂ (g)	~3 ^{194,198}	~0	(66) ¹⁵¹	MoOCl ₃ (g)	-123.1 ²⁵¹	-116	(88) ²⁵¹
MoO ₃ (c)	-178.08 ^{195,196}	-159.66	18.58 ²²	MoOCl ₄ (c)	-157.6 ^{249,251}	-133	(55) ²⁵¹
MoO ₃ (g)	~-85 ^{194,198,202}	~-81	(67) ¹⁵¹	MoOCl ₄ (g)	-140.3 ²⁵¹	-126	(91) ²⁵¹
(MoO ₃) ₂ (g)	~-28 ^{198,203,204}			MoO ₂ Cl ₂ (c)	-170.5 ²⁵²	-148	(33)
(MoO ₃) ₃ (g)	~-460 ^{198,203,204}			MoO ₂ Cl ₂ (g)	-150.4 ^{254,255}	-142	(82) ^{151,254}
(MoO ₃) ₄ (g)	~-626 ^{203,204}			MoO ₂ Cl ₂ ·H ₂ O(c)	-245.9 ²⁵²	-210	(43)
(MoO ₃) ₅ (g)	~-79 ^{1203,204}			MoBr ₂ (c)	-51.0 ^{257,258}	-48	(33) ²⁵⁸
H ₂ MoO ₄ (c)	-250.0 ²⁰⁵	-218.1	(29)	MoBr ₂ (g)	-20.8 ²⁵⁸	-24	(53) ²⁵⁸
H ₂ MoO ₄ (g)	-203 ¹⁵¹	-188	(85) ¹⁵¹	MoBr ₃ (c)	-62.7 ^{257,258}	-60	(52) ²⁵⁸
MoO ₄ ²⁻ (aq)	-238.3 ^{205,206}	-200.4	9	MoBr ₄ (c)	-70.4 ²⁵⁹	-68	(70)
Li ₂ MoO ₄ (c)	-363.3 ²⁰⁹	-337.1 ²⁰⁹	(31) ²⁰⁹	MoBr ₄ (g)	-36.7 ²⁵⁸	-46	(110) ²⁵⁸
Li ₂ MoO ₄ (g)	~-244 ²¹⁷			MoOBr ₃ (c)	-111.2 ²⁶¹	-99	(45) ²⁶¹
Na ₂ MoO ₄ (c)	-350.8 ^{205,215}	-323.6	38.1 ²¹⁰	MoOBr ₃ (g)	-75.2 ²⁶¹	-80	(101) ²⁶¹
Na ₂ MoO ₄ ·2H ₂ O(c)		-437.3 ²¹⁶		MoO ₂ Br ₂ (c)	-152.3 ^{254,260}	-135	(34) ²⁵⁴
Na ₂ Mo ₂ O ₇ (c)	-536.6 ²¹⁵	-492.0	59.9 ^{218a}	MoO ₂ Br ₂ (g)	-126.4 ²⁵⁴	-125	(86) ²⁵⁴
K ₂ MoO ₄ (c)	-358.0 ²¹⁹	-331.0	(45) ²⁰⁹	MoI ₂ (c)	(-25.5) ^{262,263}		
K ₂ MoO ₄ (g)	-272 ²²⁰			MoI ₂ (g)	32 ²⁶⁴		
KHMoO ₄ (g)	-242 ²²⁰			MoI ₃ (c)	-27.0 ^{262,263}		
Rb ₂ MoO ₄ (c)	-357.0 ²⁰⁶	-331.5	(56) ²⁰⁶	MoO ₂ I ₂ (g)	~-100 ²⁶⁵	~-101	(86) ²⁶⁵
Cs ₂ MoO ₄ (c)	-362.0 ²⁰⁷	-336.3	59.35 ²²¹	CaMoO ₃ (c)	-296 ^{266a,b}		
Cs ₂ MoO ₄ (g)	-293 ²⁰⁹			SrMoO ₃ (c)	-306 ²⁶⁷		
BeMoO ₄ (c)	~-330 ⁷³			SrMoO ₃ (g)	~-14 ¹²²²		
MgMoO ₄ (c)	-334.8 ²¹¹	-309.7 ₀	28.4 ²¹⁰	BaMoO ₃ (c)	-295 ²¹³		
MgMoO ₄ (g)	~-224 ²²²			BaMoO ₃ (g)	~-156 ²²³		
CaMoO ₄ (c)	-369.5 ²¹¹	-344.0	29.3 ²¹⁰	MoS ₂ (c)	-65.8 ²⁷⁰	-63.7	14.96 ²⁶⁸
CaMoO ₄ (ppt)	-367.3 ²¹²	-341.8	(29.3)	Mo ₂ S ₃ (c)	~-97 ²⁷⁰		
CaMoO ₄ (g)	-225 ²²²			Mo ₂ N(c)	-19.5 ¹⁸⁴		
SrMoO ₄ (c)	-369	-343.1 ²⁰⁹	(32)	MoC(c)	-2.4(?) ²⁷³		
SrMoO ₄ (g)	~-249 ²²²			Mo ₂ C(c)	-10.9 ²⁷³	-11.1	15.74 ²⁷⁴
BaMoO ₄ (c,ppt)	-369.4 ²¹³	-344.0	34.5 ²¹³	MoSi ₂ (c)	-31.5 ¹⁶⁶	-31.7	16.2 ¹⁶⁶
BaMoO ₄ (g)	~-242 ²²³			Mo ₃ Si(c)	-27.8 ¹⁶⁶	-27.7	25.4 ¹⁶⁶
Ag ₂ MoO ₄ (c)	-200.6 ²²⁴	-179.4 ^{209,224}	54.2	Mo ₅ Si ₃ (c)	-74.1 ¹⁶⁶	-74.7	49.6 ¹⁶⁶
Tl ₂ MoO ₄ (c)		-226.8 ⁷⁹		Mo ₃ Ge(c)	-5 ²⁷⁶		
MnMoO ₄ (c)	-284.7 ₆ ²²⁵			KMoF ₆ (c)	-496 ²⁸²		
FeMoO ₄ (c)	-257	-233	30.9 ²²⁷	RbMoF ₆ (c)	-498 ²⁸²		
PbMoO ₄ (c)			39.7 ²²⁸	CsMoF ₆ (c)	-500 ²⁸²		
Mo ₇ O ₂₄ ⁶⁻ (aq)	-1451 ²³³	-1255 ²³³	72 ²³³	Na ₂ MoCl ₆ (c)	-328 ²⁸⁴		
MoF ₅ (c)	-331.5 ²⁴⁴			K ₂ MoCl ₆ (c)	-350 ²⁸⁴		
MoF ₅ (g)			78.3 ²⁴⁵	Mo(CO) ₆ (c)	-236.5 ²⁶²	-211.5	78.0 ^{22,291}
				Mo(CO) ₆ (g)	-219.7 ^{173,290}	-206.5	117.8 ²⁹⁰

^a Italicized values agree with values in NBS 270-4 or NBS 270-6. Values in parentheses are estimates.

MoO₃(c) with excess OH⁻(aq) by Graham and Hepler²⁰⁵ and by O'Hare and Hoekstra²⁰⁶ both lead to $\Delta H_f^\circ = -238.3$ kcal mol⁻¹ for MoO₄²⁻(aq). This value, which we adopt for our Table IV, is in good agreement with earlier results of O'Hare and Hoekstra²⁰⁷ and is 0.2 kcal mol⁻¹ less negative than the value listed in NBS 270-4.

The NBS 270-4 lists $\Delta G_f^\circ = -199.0$ kcal mol⁻¹ and $S^\circ = 6.5$ cal K⁻¹ mol⁻¹ for MoO₄²⁻(aq). Hepler²⁰⁸ has calculated $\Delta G_f^\circ = -200.9$ kcal mol⁻¹ and $S^\circ = 11 \pm 2$ cal K⁻¹ mol⁻¹ for this ion. Most recently, O'Hare, Jensen, and Hoekstra²⁰⁹ have selected $\Delta G_f^\circ = -200.0 \pm 0.3$ kcal mol⁻¹ for MoO₄²⁻(aq), which corresponds to $S^\circ = 7.6$ cal K⁻¹ mol⁻¹. We adopt $\Delta G_f^\circ = -200.4$ kcal mol⁻¹ and $S^\circ = 9$ cal K⁻¹ mol⁻¹ for MoO₄²⁻(aq) on the basis of our new calculations that are described below in connection with various mol-

updates. This adopted ΔG_f° corresponds to the following potential:




The NBS 270-6 lists $S^\circ = 29.3$ cal K⁻¹ mol⁻¹ for CaMoO₄(c), as calculated by Weller and King²¹⁰ from results of their C_p measurements. Calorimetric measurements by Barany²¹¹ have provided a ΔH_f° of reaction that leads to $\Delta H_f^\circ = -369.5$ kcal mol⁻¹ for CaMoO₄(c). We emphasize that their "CaMoO₄(c)" was well-defined crystalline material. Muldrow and Hepler²¹² have made calorimetric measurements leading to the enthalpy of precipitation and thence to $\Delta H_f^\circ = -367.3$ kcal mol⁻¹ for material we designate CaMoO₄(ppt). The NBS

270-6 has listed the mean value $\Delta H_f^\circ = -368.4 \text{ kcal mol}^{-1}$ for $\text{CaMoO}_4(\text{c})$, but we believe it is better to regard the crystalline material denoted by (c) and the precipitated material denoted by (ppt) as different substances and therefore list both experimental ΔH_f° values^{211,212} cited above in our Table IV.

Combination of the S° from Weller and King²¹⁰ with the ΔH_f° from Barany²¹¹ leads to $\Delta G_f^\circ = -344.0 \text{ kcal mol}^{-1}$ for $\text{CaMoO}_4(\text{c})$, which is the value we adopt for crystalline calcium molybdate. We have no third law entropy for precipitated calcium molybdate, but estimate that it is the same as for $\text{CaMoO}_4(\text{c})$ and combine this value with the ΔH_f° from the enthalpy of precipitation²¹² to obtain $\Delta G_f^\circ = -341.8 \text{ kcal mol}^{-1}$ for $\text{CaMoO}_4(\text{ppt})$.

Graham and Hepler²⁰⁵ have calculated $K_{\text{sp}} \approx 1 \times 10^{-8}$ from older solubility data for calcium molybdate, while O'Hare, Jensen, and Hoekstra²⁰⁹ have cited $K_{\text{sp}} = 3.16 \times 10^{-9}$ from more recent work. These K_{sp} values correspond to $\Delta G^\circ = 10.9$ and $11.6 \text{ kcal mol}^{-1}$ for the reaction represented by



in which $\text{CaMoO}_4(\text{solid})$ might refer to either $\text{CaMoO}_4(\text{c})$ or $\text{CaMoO}_4(\text{ppt})$. We now describe two extreme calculations. Using the larger K_{sp} and the ΔG_f° for $\text{CaMoO}_4(\text{c})$, we obtain $\Delta G_f^\circ = -200.8 \text{ kcal mol}^{-1}$ for $\text{MoO}_4^{2-}(\text{aq})$ and thence $S^\circ = 10.3 \text{ cal K}^{-1} \text{ mol}^{-1}$ for this ion. Using the smaller K_{sp} and the ΔG_f° for $\text{CaMoO}_4(\text{ppt})$ leads to $\Delta G_f^\circ = -197.9 \text{ kcal mol}^{-1}$ and $S^\circ = 0.5 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{MoO}_4^{2-}(\text{aq})$. Other combinations lead to ΔG_f° and S° values intermediate between the values just quoted here.

The more negative ΔG_f° and more positive S° values above are most nearly consistent with other values we report below.

O'Hare²¹³ has made calorimetric measurements leading to the enthalpy of precipitation of barium molybdate, with considerable attention paid to experimental conditions in an effort to make the resulting $\Delta H_f^\circ = -369.4 \text{ kcal mol}^{-1}$ apply to $\text{BaMoO}_4(\text{c})$. O'Hare²¹³ has cited $S^\circ = 34.5 \pm 1.5 \text{ cal K}^{-1} \text{ mol}^{-1}$ for this substance and we calculate its $\Delta G_f^\circ = -344.0 \text{ kcal mol}^{-1}$. O'Hare²¹³ has also cited two older (1891 and 1953) investigations that are in good agreement with $K_{\text{sp}} = 3.6 \times 10^{-8}$ for barium molybdate. More recent measurements by Jost²¹⁴ lead to $K_{\text{sp}} = 3.16 \times 10^{-9}$. These K_{sp} values correspond to $\Delta G^\circ = 10.1_5$ and $11.5_9 \text{ kcal mol}^{-1}$ for the solution reaction. The larger K_{sp} in combination with the ΔG_f° of $\text{BaMoO}_4(\text{c})$ above leads to $\Delta G_f^\circ = -199.8 \text{ kcal mol}^{-1}$ and $S^\circ = 7.0 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{MoO}_4^{2-}(\text{aq})$, while the smaller K_{sp} leads to $\Delta G_f^\circ = -198.4 \text{ kcal mol}^{-1}$ and $S^\circ = 2.3 \text{ cal K}^{-1} \text{ mol}^{-1}$ for this ion. If, by analogy with calcium molybdate, we assume that the ΔH_f° and ΔG_f° values for "truly crystalline" barium molybdate are more negative than the corresponding values for the precipitated substance, then we obtain a more negative ΔG_f° and a more positive S° for $\text{MoO}_4^{2-}(\text{aq})$.

Graham and Hepler²⁰⁵ and Koehler, Pankratz, and Barany²¹⁵ have made calorimetric measurements that have led to reported values for ΔH_f° of $\text{Na}_2\text{MoO}_4(\text{c})$ in good agreement with each other. Recalculation of their^{205,215} results in terms of the NBS 270-3 auxiliary data and properties already cited in this review leads us now to $\Delta H_f^\circ = -350.8 \text{ kcal mol}^{-1}$ for $\text{Na}_2\text{MoO}_4(\text{c})$. We combine this value with $S^\circ = 38.1 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{Na}_2\text{MoO}_4(\text{c})$ from Weller and King²¹⁰ to obtain $\Delta G_f^\circ = -323.6 \text{ kcal mol}^{-1}$ for this compound. Zhidkova et al.²¹⁶ have determined the activity of sodium molybdate and water in saturated solution in equilibrium with $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}(\text{c})$, and also the vapor pressure of $\text{H}_2\text{O}(\text{g})$ in equilibrium with $\text{Na}_2\text{MoO}_4(\text{c})$ and $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}(\text{c})$. Combi-

nation of all these results with the ΔG_f° above for $\text{Na}_2\text{MoO}_4(\text{c})$ leads to $\Delta G_f^\circ = -437.3 \text{ kcal mol}^{-1}$ for $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}(\text{c})$, to $\Delta G_f^\circ = -200.0 \text{ kcal mol}^{-1}$ for $\text{MoO}_4^{2-}(\text{aq})$, and to $S^\circ = 7.8 \text{ cal K}^{-1} \text{ mol}^{-1}$ for this ion.

Consideration of uncertainties associated with the various ΔG_f° and S° values for $\text{MoO}_4^{2-}(\text{aq})$ has led us to the "best" values cited earlier in this section and listed in our Table IV. Our ΔG_f° value is slightly more negative than the values listed in NBS 270-4 and calculated by O'Hare, Jensen, and Hoekstra.²⁰⁹ Our S° is correspondingly larger than their values. We note that the second path cited by O'Hare, Jensen, and Hoekstra²⁰⁹ (their Table 5) is partly based on an estimated entropy, that their fourth path involving $\text{Ag}_2\text{MoO}_4(\text{c})$ is itself based on a prior selection of ΔG_f° of $\text{MoO}_4^{2-}(\text{aq})$, and that the uncertainty in their fifth path is sufficient to make it agree with all the values that have been cited.

For $\text{Li}_2\text{MoO}_4(\text{c})$ we adopt $\Delta H_f^\circ = -363.3 \text{ kcal mol}^{-1}$ from the calorimetric results of O'Hare, Jensen, and Hoekstra²⁰⁹ and the properties of $\text{LiOH}(\text{aq})$ cited by O'Hare et al.²⁰⁹ We agree with their²⁰⁹ conclusion that the results of an earlier investigation are erroneous. Combination of an estimated²⁰⁹ $S^\circ = (31) \text{ cal K}^{-1} \text{ mol}^{-1}$ with this ΔH_f° leads to $\Delta G_f^\circ = -337.1 \text{ kcal mol}^{-1}$ for $\text{Li}_2\text{MoO}_4(\text{c})$.

Yamdagni, Pupp, and Porter²¹⁷ have made high-temperature mass spectral investigations of the dissociation of $\text{Li}_2\text{MoO}_4(\text{g})$ and have calculated its $\Delta H_f^\circ \approx -244 \text{ kcal mol}^{-1}$. They have also determined the ΔH° of vaporization of $\text{Li}_2\text{MoO}_4(\text{liq})$, but absence of $(H_T - H_{298})$ data for the $\text{Li}_2\text{MoO}_4(\text{c})$ and $\text{liq})$ prevents us from using this result with the above ΔH_f° of $\text{Li}_2\text{MoO}_4(\text{c})$ to obtain another value of ΔH_f° for $\text{Li}_2\text{MoO}_4(\text{g})$.

We have already cited data for $\text{Na}_2\text{MoO}_4(\text{c})$ and $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}(\text{c})$ in connection with the properties of $\text{MoO}_4^{2-}(\text{aq})$. Here we note that there have also been high-temperature investigations²¹⁸ of $\text{Na}_2\text{MoO}_4\text{-MoO}_3$ mixtures.

Recalculation (using the most recent auxiliary data) of the calorimetric results of Koehler, Pankratz, and Barany²¹⁵ leads to $\Delta H_f^\circ = -536.6 \text{ kcal mol}^{-1}$ for $\text{Na}_2\text{Mo}_2\text{O}_7(\text{c})$. We take $S^\circ = 59.9 \text{ cal K}^{-1} \text{ mol}^{-1}$ from Weller and Kelley^{218a} and calculate $\Delta G_f^\circ = -492.0 \text{ kcal mol}^{-1}$ for $\text{Na}_2\text{Mo}_2\text{O}_7(\text{c})$.

The calorimetric results of Nelson, Moss, and Hepler²¹⁹ lead to $\Delta H_f^\circ = -358.0 \text{ kcal mol}^{-1}$ for $\text{K}_2\text{MoO}_4(\text{c})$. O'Hare et al.²⁰⁹ have estimated $S^\circ = (45) \text{ cal K}^{-1} \text{ mol}^{-1}$, which leads to $\Delta G_f^\circ = -331.0 \text{ kcal mol}^{-1}$ for $\text{K}_2\text{MoO}_4(\text{c})$. High-temperature measurements by Farber and Srivastava²²⁰ have provided ΔH° values that lead us to $\Delta H_f^\circ = -272 \text{ kcal mol}^{-1}$ for $\text{K}_2\text{MoO}_4(\text{g})$ and $\Delta H_f^\circ = -242 \text{ kcal mol}^{-1}$ for $\text{KHM}_2\text{O}_4(\text{g})$.

Recent calorimetric measurements by O'Hare and Hoekstra²⁰⁶ in combination with properties of $\text{RbOH}(\text{aq})$ cited by these authors lead to $\Delta H_f^\circ = -357.0 \text{ kcal mol}^{-1}$ for $\text{Rb}_2\text{MoO}_4(\text{c})$. Combination of the estimated²⁰⁶ $S^\circ = (56) \text{ cal K}^{-1} \text{ mol}^{-1}$ with this ΔH_f° leads to $\Delta G_f^\circ = -331.5 \text{ kcal mol}^{-1}$ for $\text{Rb}_2\text{MoO}_4(\text{c})$.

Calorimetric measurements by O'Hare and Hoekstra²⁰⁷ in combination with properties of $\text{CsOH}(\text{aq})$ cited by them lead to $\Delta H_f^\circ = -362.0 \text{ kcal mol}^{-1}$ for $\text{Cs}_2\text{MoO}_4(\text{c})$. We also have $S^\circ = 59.35 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{Cs}_2\text{MoO}_4(\text{c})$ from the heat capacity measurements of Osborne, Flotow, and Hoekstra.²²¹ The ΔH_f° and S° above lead to $\Delta G_f^\circ = -336.3 \text{ kcal mol}^{-1}$ for this compound.

The enthalpy of sublimation of $\text{Cs}_2\text{MoO}_4(\text{c})$ cited by O'Hare et al.²⁰⁹ leads to $\Delta H_f^\circ = -293 \text{ kcal mol}^{-1}$ for $\text{Cs}_2\text{MoO}_4(\text{g})$.

Old measurements cited in NBS Circular 500⁷³ lead to $\Delta H_f^\circ \approx -330 \text{ kcal mol}^{-1}$ for $\text{BeMoO}_4(\text{c})$.

Calorimetric results from Barany²¹¹ lead us to $\Delta H_f^\circ = -334.8_2 \text{ kcal mol}^{-1}$ for $\text{MgMoO}_4(\text{c})$, in agreement with the value in NBS 270-6. The $S^\circ = 28.4 \text{ cal K}^{-1} \text{ mol}^{-1}$ for this compound that is listed in NBS 270-6 is the same as the value calculated by Weller and King²¹⁰ from their C_p results.

The derived $\Delta G_f^\circ = -309.7_0$ kcal mol⁻¹ is also consistent with the NBS 270-6.

The NBS 270-6 lists ΔH_f° (at 0 K) = -224 kcal mol⁻¹ for MgMoO₄(g). This value is within the limits set by investigations of two reactions by Verhaegen et al.²²²

We have already discussed properties of CaMoO₄(c) and CaMoO₄(ppt) in connection with MoO₄²⁻(aq). The NBS 270-6 lists ΔH_f° (at 0 K) = -197 kcal mol⁻¹ for CaMoO₄(g). This value is consistent with the experimental results of Verhaegen et al.²²² and the NBS 270-4 ΔH_f° of MoO₃(g), but use of the ΔH_f° we have adopted for MoO₃(g) leads to our $\Delta H_f^\circ = -225$ kcal mol⁻¹ for CaMoO₄(g).

O'Hare et al.²⁰⁹ have quoted $K_{sp} = 2.58 \times 10^{-7}$ for SrMoO₄(c). We use this K_{sp} with our ΔG_f° of MoO₄²⁻(aq) to calculate $\Delta G_f^\circ = -343.1$ kcal mol⁻¹ for SrMoO₄(c). Combination of this value with $S^\circ = (32)$ cal K⁻¹ mol⁻¹ leads to $\Delta H_f^\circ = -369$ kcal mol⁻¹ for this compound. The NBS 270-6 has listed $\Delta H_f^\circ = -370$ kcal mol⁻¹ for SrMoO₄(c), taken from a source that is unknown to us.


The NBS 270-6 lists $\Delta H_f^\circ = -235$ kcal mol⁻¹ for SrMoO₄(g), which is intermediate between values calculated from results of two investigations by Verhaegen et al.²²² in combination with the NBS 270-4 ΔH_f° of MoO₃(g). A similar calculation with our ΔH_f° for MoO₃(g) leads to better agreement between the different reactions and thence to $\Delta H_f^\circ \approx -249$ kcal mol⁻¹ for SrMoO₄(g) as listed in our Table IV.

We have already discussed BaMoO₄(c) in connection with MoO₄²⁻(aq). Results of Pupp, Yamdagni, and Porter²²³ lead to $\Delta H_f^\circ \approx -242$ kcal mol⁻¹ for BaMoO₄(g).

The NBS 270-4 lists $\Delta H_f^\circ = -225$ kcal mol⁻¹ for CuMoO₄(c), apparently based on results of Tammann and Westerhold previously cited in NBS Circular 500⁷³ and by Bichowsky and Rossini.⁴⁵ Our interpretation of these uncertain results leads to $\Delta H_f^\circ = -242$ kcal mol⁻¹, which is close to the value listed by Bichowsky and Rossini.⁴⁵ We omit this uncertain value from our Table IV.

Results²²⁴ of a calorimetric determination of the enthalpy of precipitation and of solubilities at several temperatures are all in accord with $\Delta H_f^\circ = -200.6$ kcal mol⁻¹ for Ag₂MoO₄(c). Taking $K_{sp} = 2.2 \times 10^{-12}$ from the investigations cited above²²⁴ and another cited by O'Hare et al.²⁰⁹ leads to $\Delta G_f^\circ = -179.4$ kcal mol⁻¹ for Ag₂MoO₄(c) and thence by combination with the ΔH_f° to $S^\circ = 54.2$ cal K⁻¹ mol⁻¹ for this compound, whereas $S^\circ = 51$ cal K⁻¹ mol⁻¹ is given in NBS 270-4. A third law entropy based on C_p measurements is needed.

We use the ΔG_f° above to calculate the potential:



Gaultier and Pannetier⁷⁹ have made DTA measurements leading to data for phase transitions in Tl₂MoO₄(c) and have also made solubility measurements that lead to $K_{sp} = 1 \times 10^{-8}$ from which we calculate $\Delta G_f^\circ = -226.8$ kcal mol⁻¹ for this compound.

We calculate $\Delta H_f^\circ = -284.7_6$ kcal mol⁻¹ for MnMoO₄(c) from the calorimetric results of Barany.²²⁵ This value, which we adopt, is nearly identical with the value listed in NBS 270-4. The recent results of Ziolkowski and Courtine²²⁶ are in fair agreement with this value.

The NBS 270-4 has listed $S^\circ = 30.9$ cal K⁻¹ mol⁻¹ for FeMoO₄(c), which is the same as the value reported by Weller²²⁷ on the basis of C_p measurements. The NBS 270-4 has also listed $\Delta H_f^\circ = -257$ kcal mol⁻¹ for this compound, apparently based on the results of Tammann and Westerhold cited in NBS Circular 500.⁷³ Combination of ΔH_f° and S° gives the ΔG_f° . The NBS 270-4 has listed $\Delta H_f^\circ = -702$ kcal mol⁻¹ for Fe₂(MoO₄)₃(c), also apparently based on the results of Tammann and Westerhold mentioned above. We consider

this value to be quite uncertain and do not list it in our Table IV.

For PbMoO₄(c) the NBS 270-4 lists $S^\circ = 39.7$ cal K⁻¹ mol⁻¹, which is the same as the third law value found by Weller and Kelley.²²⁸ As pointed out by O'Hare et al.,²⁰⁹ the ΔG_f° in NBS 270-4 corresponds to a K_{sp} for PbMoO₄(c) smaller than any published value they (or we) know of. The $K_{sp} = 3.3 \times 10^{-12}$ cited by O'Hare et al.²⁰⁹ leads to $\Delta G_f^\circ = -221.9$ kcal mol⁻¹ for PbMoO₄(c). Combination of this ΔG_f° with the S° leads to $\Delta H_f^\circ = -246$ kcal mol⁻¹ for PbMoO₄(c). This value differs considerably from $\Delta H_f^\circ = -251.4$ kcal mol⁻¹ listed in NBS 270-4 and $\Delta H_f^\circ = -250.9$ kcal mol⁻¹ calculated from the calorimetric enthalpy of precipitation reported by Muldrow and Hepler.²¹² Because of the uncertainties as to which ΔH_f° and ΔG_f° values are best, we list only the entropy of lead molybdate.

Giauque et al.²²⁹ have investigated magnetic and thermodynamic properties of Gd₂(MoO₄)₃(c) at low temperatures.

It has long been known that acidification of solutions containing molybdate ions results in various complicated reactions involving polynuclear species that are commonly called isopolymolybdates. On the basis of many investigations cited by Sillén,^{75,112} it now appears certain that the principal species in weakly acidic solutions are heptamolybdates and octamolybdates, whereas in strongly acidic solutions there are depolymerization reactions leading to various cationic species. In addition to the summaries of results tabulated by Sillén,^{75,112} we call attention to a review by Souchay²³⁰ and a few recent investigations cited below.

Two recent spectroscopic investigations²³¹ and one very thorough kinetic investigation²³² are in general agreement with the statement above that the principal isopolymolybdate species in dilute acid are the heptamolybdate and octamolybdate ions that are commonly represented by Mo₇O₂₄⁶⁻(aq) and Mo₈O₂₆⁴⁻(aq), and also protonated forms related to these species.

We also have four calorimetric investigations²³³ of the formation of isopolymolybdates from molybdate ion or of the formation of molybdate from isopolymolybdates.

It is only for the heptamolybdate ion Mo₇O₂₄⁶⁻(aq) that equilibrium and calorimetric results of different investigations are in general agreement. We therefore limit our listings in Table IV to this ion.

Further acidification of solutions containing heptamolybdate and octamolybdate ions leads to depolymerization reactions and formation of cationic species. Reactions and equilibrium constants that refer to concentrated solutions have been summarized by Sillén^{75,112} and also recently investigated by Krumenacker,²³⁴ whose results are interpreted in terms of equilibrium constants for reactions involving such species as HMoO₃⁺(aq), H₂MoO₆²⁺(aq), etc.

Many heteropolymolybdates have been investigated and more or less well characterized. Some examples are H₃CrMo₆O₂₁, H₈CeMo₁₂O₄₂, H₃PMo₁₂O₄₀, and various related salts. Sillén^{75,112} has cited a number of determinations of solubility products and acid ionization constants for heteropolymolybdates. We call attention to recent polarographic investigations²³⁵ of oxidation-reduction properties and also determinations of acid ionization constants.²³⁶ Jespersen²³³ has made calorimetric measurements on H₃PMo₁₂O₄₀(aq).

D. Halides and Oxyhalides of Molybdenum

Heat capacity and entropy of vaporization data from Osborne et al.²³⁷ lead to $S^\circ = 62.06$ cal K⁻¹ mol⁻¹ for MoF₆(liq) and $S^\circ = 83.75$ cal K⁻¹ mol⁻¹ for MoF₆(g), both at 298 K. These are the values listed in NBS 270-4 and adopted for our Table IV. This S° of MoF₆(g) agrees with the results of a statistical thermodynamic calculation by Weinstock and Goodman.²³⁸ We also call attention to two earlier investiga-

tions. Brady et al.²³⁹ have made heat capacity measurements leading to a reported $S^\circ = 60.6 \pm 0.7 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{MoF}_6(\text{liq})$, which has been recalculated²³⁷ to be $61.96 \text{ cal K}^{-1} \text{ mol}^{-1}$. There are also vapor pressure data and derived enthalpy and entropy of vaporization values from Cady and Hargreaves.²⁴⁰

We adopt $\Delta H_f^\circ = -372.29 \text{ kcal mol}^{-1}$ for $\text{MoF}_6(\text{g})$, based on the fluorine combustion calorimetry of Settle, Feder, and Hubbard.²⁴¹ Combination of this value with the enthalpy of vaporization²⁴⁰ leads to $\Delta H_f^\circ = -378.92 \text{ kcal mol}^{-1}$ for $\text{MoF}_6(\text{liq})$. We adopt these ΔH_f° values and combine with the entropies to obtain $\Delta G_f^\circ = -351.88 \text{ kcal mol}^{-1}$ for $\text{MoF}_6(\text{g})$ and $\Delta G_f^\circ = -352.05 \text{ kcal mol}^{-1}$ for $\text{MoF}_6(\text{liq})$ as listed in Table IV.

We also call attention to some other paths to these enthalpies of formation. Myers and Brady²⁴² have measured the enthalpy of reaction of $\text{MoF}_6(\text{liq})$ with excess $\text{OH}^-(\text{aq})$ with results that lead us to $\Delta H_f^\circ = -393.2 \text{ kcal mol}^{-1}$ for $\text{MoF}_6(\text{liq})$. More recently, Nuttall, Churney, and Kilday²⁴³ have made similar measurements with results that lead us to $\Delta H_f^\circ = -376.4 \text{ kcal mol}^{-1}$ for $\text{MoF}_6(\text{liq})$. Both of these ΔH_f° values are based on the ΔH_f° of $\text{F}^-(\text{aq})$ from NBS 270-3. Allowing for six times the uncertainty in ΔH_f° of $\text{F}^-(\text{aq})$ (as discussed by Nuttall et al.²⁴³), we see that this latter ΔH_f° for $\text{MoF}_6(\text{liq})$ is in reasonable agreement with the value we have adopted from Settle, Feder, and Hubbard,²⁴¹ based on direct combination of the elements. It appears that the work of Myers and Brady²⁴² contains substantial errors. We calculate $\Delta H_f^\circ = -373.8 \text{ kcal mol}^{-1}$ for $\text{MoF}_6(\text{g})$ from the results of Burgess, Haigh, and Peacock²⁴⁴ on the reaction of $\text{MoF}_6(\text{g})$ with excess alkali, which is in fair agreement with our adopted value.

Burgess et al.²⁴⁴ have made similar experiments with $\text{MoF}_5(\text{c})$ and $\text{MoOF}_4(\text{c})$, leading to our listed ΔH_f° values for these compounds. The entropy of $\text{MoF}_5(\text{g})$ has been calculated by statistical thermodynamics.²⁴⁵

Paine and McDowell²⁴⁶ have studied the evaporation of $\text{MoOF}_4(\text{c})$ at 25°C with mass spectrometry and have shown that the vapor contains small amounts of oligomeric species, possibly $(\text{MoOF}_4)_4(\text{g})$. Therefore, we cannot calculate a reliable ΔH_f° of $\text{MoOF}_4(\text{g})$ from the reported²⁴⁷ heat of sublimation of $\text{MoOF}_4(\text{c})$.

Results of Zmbov, Uy, and Margrave²⁴⁸ lead to $\Delta H_f^\circ = -268 \text{ kcal mol}^{-1}$ for $\text{MoO}_2\text{F}_2(\text{g})$.

The NBS 270-4 lists thermodynamic properties for $\text{MoCl}_2(\text{c})$, $\text{MoCl}_3(\text{c})$, $\text{MoCl}_4(\text{c})$, $\text{MoCl}_4(\text{g})$, $\text{MoCl}_5(\text{c})$, $\text{MoCl}_5(\text{g})$, $\text{MoO}_2\text{Cl}_2(\text{c})$, $\text{MoO}_2\text{Cl}_2(\text{g})$, $\text{MoO}_2\text{Cl}_2 \cdot \text{H}_2\text{O}(\text{c})$, and $\text{MoOCl}_4(\text{c})$. In addition to all of these compounds, we also have thermodynamic data for $\text{MoOCl}_2(\text{c})$, $\text{MoOCl}_3(\text{c})$, $\text{MoOCl}_3(\text{g})$, and $\text{MoOCl}_4(\text{g})$. The thermodynamic properties of these 14 substances are related by way of thermodynamic data for about twice as many reactions. There is no set of thermodynamic properties that is consistent with all of the published results, which shows that at least some of the real uncertainties and errors in thermodynamic data are larger than quoted in the original publications. We have therefore not tried to undertake any purely statistical analysis of the published results in order to obtain a set of thermodynamic properties that leads to a minimum difference between published and calculated reaction thermodynamics. Instead, we have made tentative choices of what appear to be the best experimental results and calculated a few thermodynamic properties that have then been combined in various cycles to obtain still further thermodynamic properties. Recycling and repetition from different starting points have finally led us to the thermodynamic properties that are given in Table IV for the 14 substances listed in the beginning of this paragraph. Rather than describe in detail how we settled on these particular values, we show in following paragraphs how these values are consistent with

results of particular measurements, and also how some of them are inconsistent with results of other measurements.

Shchukarev et al.²⁴⁹ have summarized the results of a large number of previous calorimetric measurements. Some of these results lead to $\Delta H_f^\circ = -114 \text{ kcal mol}^{-1}$ for $\text{MoCl}_4(\text{c})$ and to $\Delta H_f^\circ = -126 \text{ kcal mol}^{-1}$ for $\text{MoCl}_5(\text{c})$, as listed for both of these compounds in Table IV. We note that these ΔH_f° values are based on ΔH_f° for $\text{MoO}_3(\text{c})$ and do not involve ΔH_f° values for any other molybdenum halides. The JANAF tables¹⁵¹ cite equilibrium results for the $\text{MoCl}_5(\text{c})$ - $\text{MoCl}_4(\text{c})$ - $\text{Cl}_2(\text{g})$ system that are in good agreement with the ΔH_f° values given above. The enthalpy of vaporization^{151,249} of $\text{oCl}_5(\text{c})$ leads to our tabulated ΔH_f° of $\text{MoCl}_5(\text{g})$.

We obtain $\Delta H_f^\circ = -92 \text{ kcal mol}^{-1}$ for $\text{MoCl}_4(\text{g})$ from the average of ΔH_f° values from Shchukarev et al.²⁴⁹ and Oppermann and Stöver²⁵⁰ for the decomposition of $\text{MoCl}_3(\text{c})$ to $\text{MoCl}_2(\text{c})$ and $\text{MoCl}_4(\text{g})$. For this calculation we have used the ΔH_f° values for $\text{MoCl}_2(\text{c})$ and $\text{MoCl}_3(\text{c})$ that are discussed below.

Calorimetric results summarized by Shchukarev et al.²⁴⁹ lead to $\Delta H_f^\circ = -68 \text{ kcal mol}^{-1}$ for $\text{MoCl}_2(\text{c})$. This value is dependent on ΔH_f° of $\text{MoO}_3(\text{c})$ and also on the ΔH_f° of $\text{MoO}_2\text{Cl}_2(\text{c})$, which is discussed below.

For $\text{MoCl}_3(\text{c})$ we have taken $\Delta H_f^\circ = -96.7 \text{ kcal mol}^{-1}$ from the calorimetric results of Shchukarev et al.²⁴⁹ and the equilibrium results of Oppermann, Stöver, and Kunze.²⁵¹ Calculation of ΔH_f° of $\text{MoCl}_3(\text{c})$ from these equilibrium results²⁵¹ is dependent on ΔH_f° values for various other molybdenum halides and oxyhalides.

There have been three independent investigations²⁵²⁻²⁵⁴ of the enthalpy of reaction of $\text{MoO}_2\text{Cl}_2(\text{c})$ with excess $\text{OH}^-(\text{aq})$. These results lead to $\Delta H_f^\circ = -170.5$, -173.2 , and $-175.4 \text{ kcal mol}^{-1}$ for $\text{MoO}_2\text{Cl}_2(\text{c})$. Because of the troublesome reaction of $\text{MoO}_2\text{Cl}_2(\text{c})$ with atmospheric water, we expect that the most exothermic enthalpy of reaction is best²⁵² and therefore adopt $\Delta H_f^\circ = -170.5 \text{ kcal mol}^{-1}$ for this compound.

Our tabulated ΔH_f° of $\text{MoO}_2\text{Cl}_2(\text{g})$ is consistent with the above ΔH_f° for $\text{MoO}_2\text{Cl}_2(\text{c})$ and the enthalpy of sublimation reported by Oppermann²⁵⁴ and by Shchukarev et al.²⁵⁵ The equilibrium data from Shchukarev et al.²⁵⁵ and from Hultgren and Brewer²⁵⁶ on the reaction between $\text{MoO}_3(\text{c})$ and $\text{HCl}(\text{g})$ to form $\text{MoO}_2\text{Cl}_2(\text{g})$ and $\text{H}_2\text{O}(\text{g})$ lead to a slightly less negative ΔH_f° for $\text{MoO}_2\text{Cl}_2(\text{g})$. All of these results support our choice for ΔH_f° of $\text{MoO}_2\text{Cl}_2(\text{c})$ in the paragraph above.

Our tabulated $\Delta H_f^\circ = -150.7 \text{ kcal mol}^{-1}$ for $\text{MoOCl}_3(\text{c})$ is consistent with the calorimetric results of Oppermann, Stöver, and Kunze.²⁵¹ The $\Delta H_f^\circ = -123.1 \text{ kcal mol}^{-1}$ for $\text{MoOCl}_3(\text{g})$ is based on the above ΔH_f° for $\text{MoOCl}_3(\text{c})$ and the enthalpy of sublimation.²⁵¹

For $\text{MoOCl}_4(\text{c})$ we adopt $\Delta H_f^\circ = -157.6 \text{ kcal mol}^{-1}$ from results of two calorimetric investigations^{249,251} that differ from each other by 6 kcal mol^{-1} . The enthalpy of vaporization that leads to $\Delta H_f^\circ = -140.3 \text{ kcal mol}^{-1}$ for $\text{MoOCl}_4(\text{g})$ is from Oppermann et al.²⁵¹


Our $\Delta H_f^\circ = -126.2 \text{ kcal mol}^{-1}$ for $\text{MoOCl}_2(\text{c})$ is based on the equilibrium results of Oppermann and Stöver²⁵⁰ for decomposition of $\text{MoOCl}_2(\text{c})$ to $\text{MoCl}_2(\text{c})$ and $\text{MoO}_2\text{Cl}_2(\text{g})$.

We also list entropies for the molybdenum halides and oxyhalides. These estimated entropies are intended to be consistent with results of various equilibrium investigations referred to above. Entropies based on measured heat capacities and the third law are needed.


We adopt $\Delta H_f^\circ = -245.9 \text{ kcal mol}^{-1}$ for the compound listed in Table IV as $\text{MoO}_2\text{Cl}_2 \cdot \text{H}_2\text{O}(\text{c})$, based on a calorimetric measurement of the enthalpy of reaction with excess $\text{OH}^-(\text{aq})$.²⁵² This compound, which is commonly prepared by reaction of $\text{MoO}_3(\text{c})$ with $\text{HCl}(\text{g})$ at about 250°C , has also been represented as $\text{MoO}(\text{OH})_2\text{Cl}_2(\text{c})$ and $\text{MoO}_3 \cdot 2\text{HCl}(\text{c})$. We

estimate that the entropy of this compound is $10 \text{ cal K}^{-1} \text{ mol}^{-1}$ greater than that of $\text{MoO}_2\text{Cl}_2(\text{c})$ and calculate the ΔG_f° . This ΔG_f° in combination with that of $\text{MoO}_3(\text{c})$ leads to a calculated 0.04 atm (at 298 K) for the pressure of $\text{HCl}(\text{g})$ in equilibrium with $\text{MoO}_3 \cdot 2\text{HCl}(\text{c})$ and $\text{MoO}_3(\text{c})$. A similar calculation with the ΔG_f° of $\text{MoO}_2\text{Cl}_2(\text{c})$ leads to $2 \times 10^{-5} \text{ atm}$ for the pressure of $\text{H}_2\text{O}(\text{g})$ in equilibrium with $\text{MoO}_2\text{Cl}_2 \cdot \text{H}_2\text{O}(\text{c})$ and $\text{MoO}_2\text{Cl}_2(\text{c})$. Both of these calculated equilibrium pressures are in accord with the properties of these compounds.

Our Table IV lists $\Delta H_f^\circ = -51.0 \text{ kcal mol}^{-1}$ for $\text{MoBr}_2(\text{c})$ and $\Delta H_f^\circ = -62.7 \text{ kcal mol}^{-1}$ for $\text{MoBr}_3(\text{c})$. A calorimetric determination by Shukurov et al.²⁵⁷ of the heats of combustion of these compounds leads to $\Delta H_f^\circ = -54 \pm 5 \text{ kcal mol}^{-1}$ for $\text{MoBr}_2(\text{c})$ and to $\Delta H_f^\circ = -64 \pm 5 \text{ kcal mol}^{-1}$ for $\text{MoBr}_3(\text{c})$. Equilibrium data from Oppermann²⁵⁸ lead to $\Delta H^\circ_{\text{reaction}} = 74.8 \text{ kcal}$ for



and to $\Delta H^\circ_{\text{reaction}} = 25.5 \text{ kcal}$ for



Our selected values have been chosen to be reasonably consistent with both the calorimetric and the equilibrium data.

Oppermann²⁵⁸ has determined the enthalpy of sublimation of $\text{MoBr}_2(\text{c})$, and we have calculated $\Delta H_f^\circ = -20.8 \text{ kcal mol}^{-1}$ for $\text{MoBr}_2(\text{g})$ from his data.

The results of two equilibrium studies by Oppermann²⁵⁸ in combination with our adopted values of ΔH_f° for $\text{MoBr}_2(\text{c})$ and $\text{MoBr}_3(\text{c})$ give $\Delta H_f^\circ = -36.7 \text{ kcal mol}^{-1}$ for $\text{MoBr}_4(\text{g})$. Shchukarev et al.²⁵⁹ have determined the enthalpies of solution of $\text{MoBr}_4(\text{c})$ and $\text{MoCl}_5(\text{c})$, leading to our $\Delta H_f^\circ = -70.4 \text{ kcal mol}^{-1}$ for $\text{MoBr}_4(\text{c})$. This value depends on our adopted ΔH_f° for $\text{MoCl}_5(\text{c})$.

Table IV lists $\Delta H_f^\circ = -152.3 \text{ kcal mol}^{-1}$ for $\text{MoO}_2\text{Br}_2(\text{c})$, which is the average of two calorimetric determinations,^{254,260} and also $\Delta H_f^\circ = -126.4 \text{ kcal mol}^{-1}$ for $\text{MoO}_2\text{Br}_2(\text{g})$ from the heat of sublimation determined by Oppermann.²⁵⁴ Oppermann et al.²⁶¹ have made similar studies on $\text{MoOBr}_3(\text{c})$ and $\text{MoOBr}_3(\text{g})$, leading to our $\Delta H_f^\circ = -111.2 \text{ kcal mol}^{-1}$ for $\text{MoOBr}_3(\text{c})$ and $\Delta H_f^\circ = -75.2 \text{ kcal mol}^{-1}$ for $\text{MoOBr}_3(\text{g})$.

Our estimated entropies for the molybdenum bromides and oxybromides are consistent with the equilibrium measurements of Oppermann et al.^{258,261} The uncertainty in these entropies is considerable (maybe $\pm 10 \text{ cal K}^{-1} \text{ mol}^{-1}$), but the internal consistency of the values is (probably) much better.

The $\Delta H_f^\circ = -27.0 \text{ kcal mol}^{-1}$ for $\text{MoI}_3(\text{c})$ has been calculated by Skinner et al.^{262,263} from the heat of reaction between $\text{Mo}(\text{CO})_6(\text{c})$ and $\text{I}_2(\text{g})$ and the ΔH_f° of $\text{Mo}(\text{CO})_6(\text{c})$, which we will discuss later. They also estimate ΔH_f° for $\text{MoI}_2(\text{c})$ to be $1.5 \text{ kcal mol}^{-1}$ less negative than ΔH_f° for $\text{MoI}_3(\text{c})$. These estimated and measured values are adopted for Table IV.

NBS 270-4 lists $\Delta H_f^\circ = 32 \text{ kcal mol}^{-1}$ for $\text{MoI}_2(\text{g})$, which is consistent with our recalculation of Allen's results²⁶⁴ for the equilibrium between $\text{Mo}(\text{c})$, $\text{I}(\text{g})$, and $\text{MoI}_2(\text{g})$ at 1200 K .

Oppermann²⁶⁵ claims to have shown that $\text{MoO}_2\text{I}_2(\text{g})$ is formed in the reaction between $\text{MoO}_2(\text{c})$ and $\text{I}_2(\text{g})$ at about 1000 K , and he has derived the uncertain values of ΔH_f° and S° we have included in Table IV for $\text{MoO}_2\text{I}_2(\text{g})$.

E. Other Molybdenum Compounds

Barancheeva and Zharkova²⁶⁶ have determined ΔH_f° for calcium molybdate, $\text{CaMoO}_4(\text{c})$, by combustion calorimetry. We calculate $\Delta H_f^\circ = -280 \text{ kcal mol}^{-1}$ from the published heat of combustion, using our ΔH_f° for $\text{CaMoO}_4(\text{c})$. Earlier high-temperature equilibrium^{266a} and high-temperature emf^{266b} investigations lead to $\Delta H_f^\circ = -296 \text{ kcal mol}^{-1}$ for

$\text{CaMoO}_4(\text{c})$, in agreement with the value listed in NBS 270-6. We accept the latter value.

Our $\Delta H_f^\circ = -306 \text{ kcal mol}^{-1}$ for $\text{SrMoO}_3(\text{c})$ is calculated from combustion calorimetry results from Zharkova and Barancheeva,²⁶⁷ and is the same as the value listed in NBS 270-6.

NBS 270-6 also lists $\Delta H_f^\circ = -130 \text{ kcal mol}^{-1}$ (at 0 K) for $\text{SrMoO}_3(\text{g})$, which is the mean of two investigations by Verhaegen et al.,²²² calculated with the NBS 270-4 ΔH_f° for $\text{MoO}_3(\text{g})$. With our adopted ΔH_f° for $\text{MoO}_3(\text{g})$ we obtain $\Delta H_f^\circ \approx -141 \text{ kcal mol}^{-1}$ for $\text{SrMoO}_3(\text{g})$ and a better agreement between the two values of ΔH_f° that can be calculated from Verhaegen's results.

For $\text{BaMoO}_3(\text{c})$ a calorimetric and a high-temperature equilibrium investigation (discussed by O'Hare²¹³) are in good agreement with $\Delta H_f^\circ = -295 (\pm 2) \text{ kcal mol}^{-1}$, as adopted in NBS 270-6 and in our Table IV.

Our $\Delta H_f^\circ \approx -156 \text{ kcal mol}^{-1}$ for $\text{BaMoO}_3(\text{g})$ is calculated from the high-temperature equilibrium data of Pupp et al.²²³ and our adopted ΔH_f° for $\text{MoO}_2(\text{g})$.

For $\text{MoS}_2(\text{c})$, NBS 270-4 lists $\Delta H_f^\circ = -56.2 \text{ kcal mol}^{-1}$ and $S^\circ = 14.96 \text{ cal K}^{-1} \text{ mol}^{-1}$. The entropy is from C_p measurements by Westrum and McBride,²⁶⁸ while the ΔH_f° value is consistent with the results of two old investigations.²⁶⁹ O'Hare et al.²⁷⁰ have made calorimetric measurements that lead to $\Delta H_f^\circ = -65.8 \text{ kcal mol}^{-1}$. This value, which we adopt, is in reasonable agreement with some high-temperature studies, as discussed by O'Hare et al.²⁷⁰ We agree with the analysis by O'Hare et al.²⁷⁰ of high-temperature equilibrium and emf data for $\text{Mo}_2\text{S}_3(\text{c})$, leading to an approximate $\Delta H_f^\circ \approx -97 \text{ kcal mol}^{-1}$ for this compound.

Parravano and Malquori²⁶⁹ have made equilibrium studies on the system $\text{MoS}_3\text{--MoS}_2\text{--S}_2$, but their conclusion regarding the stability of MoS_3 is at variance with that of another investigation.²⁷¹ Therefore, we do not list $\text{MoS}_3(\text{c})$ in Table IV.

The enthalpy of decomposition of $(\text{NH}_4)_2\text{MoS}_4(\text{c})$ is known from equilibrium measurements on its thermal decomposition,²⁷² but the absence of a ΔH_f° value for $\text{MoS}_3(\text{c})$ prevents us from calculating a ΔH_f° for $(\text{NH}_4)_2\text{MoS}_4(\text{c})$.

For the nitride, $\text{Mo}_2\text{N}(\text{c})$, we have $\Delta H_f^\circ = -19.5 \pm 0.3 \text{ kcal mol}^{-1}$ from combustion calorimetry by Mah.¹⁶⁴ This value is listed in NBS 270-4.

Mah²⁷³ has also determined $\Delta H_f^\circ = -2.4 \text{ kcal mol}^{-1}$ for "MoC" and $\Delta H_f^\circ = -10.9 \text{ kcal mol}^{-1}$ for Mo_2C . As it was not known by Mah that the high-carbon carbide is carbon deficient with an actual composition of $\text{MoC}_{0.67}$, the heat of formation of "MoC" might be considerably in error.²⁷ For $\text{Mo}_2\text{C}(\text{c})$ we have $S^\circ = 15.74 \text{ cal K}^{-1} \text{ mol}^{-1}$ from C_p measurements.²⁷⁴ Our ΔG_f° in Table IV is consistent with the S° and ΔH_f° values.

High-temperature heat contents²⁷⁴ combined with the adopted ΔH_f° and S° lead to a free energy of formation of $\text{Mo}_2\text{C}(\text{c})$ which at 800 K is about $1.2 \text{ kcal mol}^{-1}$ more negative than the value calculated by Solbakken and Emmett²⁷⁵ from equilibrium measurements on the reaction of $\text{Mo}(\text{c})$ and $\text{CH}_4(\text{g})$ to form $\text{Mo}_2\text{C}(\text{c})$ and $\text{H}_2(\text{g})$. We have no explanation for this discrepancy.

We adopt enthalpies of formation and entropies for $\text{Mo}_3\text{Si}(\text{c})$, $\text{MoSi}_2(\text{c})$, and $\text{Mo}_5\text{Si}_3(\text{c})$ from Chart's recent review.¹⁶⁶ The slightly less negative ΔH_f° values listed in NBS 270-4 for these compounds are within the uncertainty limits assigned by Chart. The tabulated enthalpies and entropies lead to our free energy values.

NBS 270-4 lists $\Delta H_f^\circ = -5 \text{ kcal mol}^{-1}$ for $\text{Mo}_3\text{Ge}(\text{c})$, consistent with the results of vapor pressure measurements by Peavler and Searcy.²⁷⁶

Potential measurements by El-Shamy and El-Aggan²⁷⁷ on the $\text{Mo}(\text{VI})\text{--Mo}(\text{V})$ couple in aqueous HCl lead to $E^\circ = 0.48 \text{ V}$ at 30°C , but the molybdenum species are ill-defined and we

do not try to write a redox equation. Haight²⁷⁸ has shown that MoOCl_5^{2-} is the most important Mo(V) species in concentrated HCl, and that the dinuclear ion $(\text{MoOCl}_4)_2\text{O}^{4-}$ is the initial hydrolysis product. Haight²⁷⁸ has evaluated the equilibrium constants at several temperatures and thence the enthalpy of reaction for this dimerization process. His results are in general agreement with more recent work, in which the formation of dinuclear cationic species in more dilute acid solution has also been demonstrated.²⁷⁹ Saha et al.²⁸⁰ have prepared the acids $\text{H}_2\text{MoOCl}_5 \cdot 2\text{H}_2\text{O}$ and HMoCl_4 in the solid state and have also shown that the corresponding bromides have similar properties. Marov et al.²⁸¹ have reported equilibrium constants for the reactions between MoOX_4^- ($X = \text{NCS}^-$ or Cl^-) and Br^- , I^- , or H_3PO_4 in strongly acidic solutions.


Solution calorimetry by Burgess et al.²⁸² has led to the ΔH_f° values we list in Table IV for $\text{KMoF}_6(\text{c})$, $\text{RbMoF}_6(\text{c})$, and $\text{CsMoF}_6(\text{c})$.

The complex formation between CH_3O^- and Mo(V) in absolute methanol has been investigated by Gut, Schmid, and Serrallach.²⁸³

Efimov and Belorukova²⁸⁴ have described the synthesis of Mo(IV) compounds of the type M_2MoCl_6 ($M = \text{Na}, \text{K}, \text{Rb},$ or Cs) and evaluated ΔH_f° for the sodium and potassium compounds by solution calorimetry.

The stable complex ion $\text{Mo}(\text{CN})_8^{4-}$ forms soluble complexes with trivalent metal ions, and the stability constant of the Fe^{3+} complex has been estimated.²⁸⁵

Potentials have been reported²⁸⁶ for the half-reaction



but it is necessary to realize that evaluation of activity coefficient and thence of a standard potential is uncertain.

It has been shown that the green solutions of Mo(III) in HCl contain a mixture of Mo(III) species.²⁸⁷ Recently, the green dinuclear ion $(\text{H}_2\text{O})_4\text{Mo}(\text{OH})_2\text{Mo}(\text{H}_2\text{O})_4^{4+}$ and the yellow hexaquo ion $\text{Mo}(\text{H}_2\text{O})_6^{3+}$ have been identified.^{287,288}

We have no quantitative data on Mo(II) in aqueous solution. The identification of some aqueous Mo(II) species is discussed in recent papers.²⁸⁹

NBS 270-4 reports $\Delta H_f^\circ = -234.9 \text{ kcal mol}^{-1}$ for $\text{Mo}(\text{CO})_6(\text{c})$, in accordance with bomb calorimetric results of Cotton, Fischer, and Wilkinson.¹⁷³ More recently, Skinner et al.²⁶² propose $\Delta H_f^\circ = -236.5 \text{ kcal mol}^{-1}$ as a weighted average of their results for the heat of combustion and heat of decomposition of $\text{Mo}(\text{CO})_6(\text{c})$. Cotton et al.¹⁷³ report $\Delta H_{\text{vaporization}}^\circ = 16.3 \text{ kcal mol}^{-1}$, while Monchamp and Cotton²⁹⁰ find $\Delta H_{\text{vaporization}}^\circ = 17.3 \text{ kcal mol}^{-1}$. We adopt $\Delta H_f^\circ = -236.5 \text{ kcal mol}^{-1}$ for $\text{Mo}(\text{CO})_6(\text{c})$ and $\Delta H_f^\circ = -219.7 \text{ kcal mol}^{-1}$ for $\text{Mo}(\text{CO})_6(\text{g})$, using the average value of the quoted enthalpies of vaporization. From the results of their C_p measurements, Astrov et al.²⁹¹ have calculated $S^\circ = 78.17 \pm 0.25 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{Mo}(\text{CO})_6(\text{c})$. On the basis of the same data, Kelley and King²² give $S^\circ = 77.9 \pm 0.8 \text{ cal K}^{-1} \text{ mol}^{-1}$. We adopt $S^\circ = 78.0 \text{ cal K}^{-1} \text{ mol}^{-1}$ in Table IV. Monchamp and Cotton²⁹⁰ have discussed earlier spectroscopic investigations on $\text{Mo}(\text{CO})_6(\text{g})$ and have made statistical calculations on the basis of these spectroscopic data, leading to values of S° for $\text{Mo}(\text{CO})_6(\text{g})$ from 117.6 to 122.6 $\text{cal K}^{-1} \text{ mol}^{-1}$. From the vapor pressure measurements of Monchamp and Cotton²⁹⁰ we calculate and adopt $S^\circ = 117.8 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{Mo}(\text{CO})_6(\text{g})$.

There are few thermodynamic investigations on organic molybdenum compounds. In addition to data covered by earlier reviews,^{180,181} we note that the heats of formation of adducts $\text{MoCl}_4 \cdot 2\text{L}$ ($\text{L} = \text{pyridine}, \text{tetrahydrofuran},$ or $\text{tetrahydrothiophene}$) have been determined.²⁹²

F. Tungsten, Tungsten Oxides, and Tungstic Acid

We take $\Delta H_f^\circ = 0$, $\Delta G_f^\circ = 0$, and $S^\circ = 7.80 \text{ cal K}^{-1}$

mol^{-1} from NBS 270-4 for W(c). The entropy is the same as listed by Kelley and King²² and in the JANAF tables.¹⁵¹

For W(g) we have $S^\circ = 41.549 \text{ cal K}^{-1} \text{ mol}^{-1}$, also in agreement with the value given by Kelley and King²² and in NBS 270-4. On the basis of vapor pressure measurements by Plante and Sessoms²⁹³ and earlier results cited by them, we adopt $\Delta H_f^\circ = 204.5 \text{ kcal mol}^{-1}$ for W(g). Combination of ΔH_f° and S° leads to $\Delta G_f^\circ = 194.4 \text{ kcal mol}^{-1}$.

The heat capacity, enthalpy, and related thermodynamic properties of tungsten at high temperatures have been the subject of a recent investigation;²⁹⁴ earlier results are discussed in the JANAF tables.¹⁵¹

For $\text{WO}(\text{g})$ we adopt $\Delta H_f^\circ \simeq 102 \text{ kcal mol}^{-1}$ from the work of DeMaria et al.,¹⁹⁴ as also discussed in the JANAF tables.¹⁵¹ The uncertainty in this value overlaps the NBS 270-4 value of $\Delta H_f^\circ = 108 \text{ kcal mol}^{-1}$. We use an estimated¹⁵¹ $S^\circ = (59) \text{ cal K}^{-1} \text{ mol}^{-1}$ to calculate $\Delta G_f^\circ \simeq 94 \text{ kcal mol}^{-1}$.

The $\Delta H_f^\circ = -140.94 \text{ kcal mol}^{-1}$ for $\text{WO}_2(\text{c})$ given in NBS 270-4 is based on combustion calorimetry by Mah.²⁹⁵ Charlu and Kleppa²⁹⁶ obtained $\Delta H_f^\circ = -140.2 \pm 0.7 \text{ kcal mol}^{-1}$ by this same method. Our recalculations of high-temperature emf results^{197,297} lead to $\Delta H_f^\circ = -139.5$ and $-140.4 \text{ kcal mol}^{-1}$, respectively. Earlier equilibrium measurements discussed in the JANAF tables¹⁵¹ lead us to $\Delta H_f^\circ = -140.6 \pm 0.7 \text{ kcal mol}^{-1}$. We adopt the value from Mah in our Table V, but note that there is some evidence for a slightly less negative value. We adopt $S^\circ = 12.08 \text{ cal K}^{-1} \text{ mol}^{-1}$ as given by Kelley and King²² and listed in NBS 270-4, and calculate $\Delta G_f^\circ = -127.6 \text{ kcal mol}^{-1}$ for $\text{WO}_2(\text{c})$.

We have $\Delta H_f^\circ \simeq 15 \text{ kcal mol}^{-1}$ for $\text{WO}_2(\text{g})$ as a mean value of third law recalculations of the results from DeMaria et al.¹⁹⁴ and Chupka, Berkowitz, and Giese.²⁹⁸ NBS 270-4 lists $\Delta H_f^\circ = 11 \text{ kcal mol}^{-1}$. We adopt $S^\circ = (68) \text{ cal K}^{-1} \text{ mol}^{-1}$ from the JANAF tables¹⁵¹ and thence calculate $\Delta G_f^\circ \simeq 12 \text{ kcal mol}^{-1}$ for $\text{WO}_2(\text{g})$.

For $\text{WO}_{2.72}(\text{c})$ ($= \frac{1}{18}\text{W}_{18}\text{O}_{49}$) we adopt $\Delta H_f^\circ = -186.3 \text{ kcal mol}^{-1}$ from two investigations.^{296,297} The analysis of older data in the JANAF tables¹⁵¹ also supports a more negative value than the $\Delta H_f^\circ = -185 \text{ kcal mol}^{-1}$ listed in NBS 270-4. The ΔG_f° value in Table V has been calculated from our ΔH_f° and an estimated¹⁵¹ $S^\circ = (16.4) \text{ cal K}^{-1} \text{ mol}^{-1}$.

For $\text{WO}_{2.90}(\text{c})$ ($= \frac{1}{20}\text{W}_{20}\text{O}_{58}$) we adopt $\Delta H_f^\circ = -196.0 \text{ kcal mol}^{-1}$ as listed in NBS 270-4 and in the JANAF tables.¹⁵¹ This value is consistent with two investigations.^{296,297} Table V gives $\Delta G_f^\circ = -177.7 \text{ kcal mol}^{-1}$, calculated from our ΔH_f° and an estimated¹⁵¹ $S^\circ = (17.5) \text{ cal K}^{-1} \text{ mol}^{-1}$.

Our values for $\text{WO}_{2.96}(\text{c})$ are the same as listed in the JANAF tables.¹⁵¹

For $\text{WO}_3(\text{c})$ we adopt $\Delta H_f^\circ = -201.45 \text{ kcal mol}^{-1}$ from NBS 270-4, in agreement with bomb calorimetry results of Mah.²⁹⁵ This value also agrees well with $\Delta H_f^\circ = -201.84 \text{ kcal mol}^{-1}$ from an earlier investigation.²⁹⁹ The NBS 270-4 lists $S^\circ = 18.14 \text{ cal K}^{-1} \text{ mol}^{-1}$, in good agreement with the value given by Kelley and King,²² and thence $\Delta G_f^\circ = -182.62 \text{ kcal mol}^{-1}$ for $\text{WO}_3(\text{c})$.

A third law analysis of high-temperature data from three investigations^{194,298,300} leads to $\Delta H_f^\circ \simeq -70 \text{ kcal mol}^{-1}$ for $\text{WO}_3(\text{g})$. Combining this value with an estimated¹⁵¹ $S^\circ = (68) \text{ cal K}^{-1} \text{ mol}^{-1}$ gives $\Delta G_f^\circ \simeq -66 \text{ kcal mol}^{-1}$.

Our enthalpies of formation for $(\text{WO}_3)_n(\text{g})$ ($n = 2, 3, 4$) and for $\text{W}_3\text{O}_8(\text{g})$ are all based on the high-temperature vapor pressure and mass spectrometric measurements by Ackermann and Rauh.³⁰¹ They have been combined with estimated¹⁵¹ entropies to give ΔG_f° values. The values listed in NBS 270-4 for ΔH_f° of $(\text{WO}_3)_3(\text{g})$ and $(\text{WO}_3)_4(\text{g})$ are $\sim 20 \text{ kcal mol}^{-1}$ less negative than our values and are in better agreement with the results of Norman and Staley,³⁰² whose results are inconsistent with the free energy functions in the JANAF

TABLE V.^a Thermodynamic Properties of Tungsten Compounds at 298 K

Compound	ΔH_f° kcal mol ⁻¹	ΔG_f° kcal mol ⁻¹	S° cal K ⁻¹ mol ⁻¹	Compound	ΔH_f° kcal mol ⁻¹	ΔG_f° kcal mol ⁻¹	S° cal K ⁻¹ mol ⁻¹
W(c)	0	0	7.80 ²²	WO ₂ F ₂ (g)	~-215 ²⁴⁸		
W(g)	204.5 ²⁹³	194.4	41.549 ²²	WCl ₂ (c)	-67 ²⁴⁹	-52	(31) ¹⁵¹
W ⁺ (g)	390.1			WCl ₄ (c)	~-108 ²⁴⁹	~-88	(47) ¹⁵¹
WO(g)	~102 ¹⁹⁴	~94	(59) ¹⁵¹	WCl ₄ (g)	~-77 ²⁴⁹	~-70	(91) ¹⁵¹
WO ₂ (c)	-140.94 ²⁹⁵	-127.6	12.08 ²²	WCl ₅ (c)	-122 ²⁴⁹	-95	(52) ¹⁵¹
WO ₂ (g)	~15 ^{194,298}	~12	(68) ¹⁵¹	WCl ₅ (g)	-98 ²⁴⁹	-85	(97) ¹⁵¹
WO _{2.72} (c)	-186.3 ^{296,297}	-169.0	(16.4) ¹⁵¹	(WCl ₅) ₂ (g)	-206 ²⁴⁹	-173	(170) ¹⁵¹
WO _{2.90} (c)	-196.0 ^{151,296,297}	-177.7	(17.5) ¹⁵¹	WCl ₆ (c,α)	-141.9 ²⁴⁹	-109	(57) ¹⁵¹
WO _{2.96} (c)	-199.6 ¹⁵¹	-180.9	(17.9) ¹⁵¹	WCl ₆ (g)	-118.0 ¹⁵¹	-98	(100) ¹⁵¹
WO ₃ (c)	-201.45 ²⁹⁵	-182.62	18.14	WCl ₆ (g)	-147 ³⁴⁴	-131	(32) ³⁴⁴
WO ₃ (g)	~-70 ^{194,298,300}	~-66	(68) ¹⁵¹	WCl ₃ (c)	-165.9 ³⁴⁴	-145	(43) ³⁴⁴
(WO ₃) ₂ (g)	~-277 ³⁰¹	~-258	(99) ¹⁵¹	WCl ₄ (c)	-172.4 ³⁴¹	-143	(41) ¹⁵¹
(WO ₃) ₃ (g)	~-484 ³⁰¹	~-447	(121) ¹⁵¹	WCl ₄ (g)	-152.4 ^{151,342}	-138	92.1 ³⁴³
(WO ₃) ₄ (g)	~-671 ³⁰¹	~-617	(145) ¹⁵¹	WO ₂ Cl ₂ (c)	-189 ^{249,341}	-170	(48) ¹⁵¹
W ₃ O ₈ (g)	~-409 ³⁰¹	~-379	(118) ¹⁵¹	WO ₂ Cl ₂ (g)	-162 ^{151,341}	-155	(85) ¹⁵¹
H ₂ WO ₄ (c)	-270.0 ³⁰⁵	-239.0	(33)	WBr ₅ (c)	-75.4 ³⁴⁵	-65	(65) ¹⁵¹
H ₂ WO ₄ (g)	~-217 ³⁰⁶	~-201	(84) ¹⁵¹	WBr ₅ (g)	-52 ³⁴⁶	-27	(110) ¹⁵¹
WO ₄ ²⁻ (aq)	-256.5 ³⁰⁴	-219	(11)	WBr ₆ (c)	-83.2 ³⁴⁵	-71	(75) ¹⁵¹
HW ₆ O ₂₁ ⁵⁻ (aq)	-1396.6 ³¹³	-1226.8 ^{311,312}	86	WBr ₆ (g)	-126.0 ³⁵¹	-114	(46) ³⁴¹
Li ₂ WO ₄ (c)	-383.2 ²⁰⁹			WBr ₄ (c)	-131.8 ^{347,348}	-118	(55) ^b
Li ₂ WO ₄ (g)	~-240 ²¹⁷			WBr ₄ (g)	-101.9 ³⁴⁸⁻³⁵⁰	-103	(105) ^b
Na ₂ WO ₄ (c)	-369.6 ^{215,322}	-342.2	38.5 ³²¹	WO ₂ Br ₂ (c)	-170.2 ^{347,348}		
Na ₂ W ₂ O ₇ (c)	-573.8 ²¹⁵	-528.8	60.8 ^{218a}	WO ₂ Br ₂ (g)	-133.0 ³⁴⁸⁻³⁵⁰	-131	(86) ³⁴⁹
Na ₂ W ₄ O ₁₃ (c)	-990.9 ²¹⁵			WI ₂ (c)	(-10) ²⁶³		
K ₂ WO ₄ (c)	-377.9 ²⁰⁹			WI ₃ (c)	~-10 ²⁶³		
K ₂ WO ₄ (g)	~-278 ²²⁰			WO ₂ (l)	~-137 ³⁵³		
KHWO ₄ (g)	~-257 ²²⁰			WO ₂ l ₂ (c)	-145.4 ³⁵³		
MgWO ₄ (c)	-362.7 ³²⁴	-336.0	24.18 ³²¹	WO ₂ l ₂ (g)	-102.3 ^{349,353}	-104	(89) ³⁵³
MgWO ₄ (g)	~-215 ²²²			WF ₅ Cl(liq)	-390 ³³⁸		
CaWO ₄ (c)	-392.2 ²¹¹	-366.7	30.27 ³²¹	WF ₄ Cl ₂ (liq)	-352 ³³⁸		
CaWO ₄ (g)	~-226 ²²²			CaWO ₃ (g)	~-118 ²²²		
SrWO ₄ (c)	~-391	-366 ³²⁵	(33)	KWF ₆ (c)	-541 ³⁵⁴		
SrWO ₄ (g)	~-250 ²²²			KWCl ₆ (c)	-238 ³⁵⁵		
BaWO ₄ (c)	-407 ⁴⁵			K ₂ WCl ₇ (c)	-342 ³⁵⁵		
SnWO ₄ (g)	~-417 ²²²			K ₃ W ₂ Cl ₉ (c)	~-530 ³⁵⁶		
Sn ₂ WO ₅ (g)	~-472 ²²²			Rb ₃ W ₂ Cl ₉ (c)	~-571 ³⁵⁶		
MnWO ₄ (c)	-311.9 ³²⁸			Cs ₃ W ₂ Cl ₉ (c)	~-578 ³⁵⁶		
FeWO ₄ (c)	-276 ⁴⁵	-252	31.61 ³³⁰	WF ₅ (CH ₃ O)(c)	-417 ³³⁸		
CoWO ₄ (c)	-270.7 ³²⁴			cis-WF ₄ (CH ₃ O) ₂ (liq)	-404 ³³⁸		
NiWO ₄ (c)	-269.9 ³²⁴			cis-WF ₄ (CH ₃ O) ₄ (c)	-362 ³³⁸		
CuWO ₄ (c)	-245.5 ³²⁴			WS ₂ (c)	-54 ³⁵⁷	-56	29 ³⁵⁷
ZnWO ₄ (c)	-294.6 ³²⁴	-268.5	28.34 ³³⁰	WC(c)	-9.6 ₃ ^{273,358}		
CdWO ₄ (c)	-281.8 ³²⁴			W ₂ C(c)	-6.3 ²⁷³		
PbWO ₄ (c)	-271(?)	-247	40.2 ²²⁸	W(CO) ₆ (c)	-227.9 ¹⁷³		
Ag ₂ WO ₄ (c)	-220.7 ³²²	-198.9	(53)	W(CO) ₆ (g)	-210.2 ¹⁷³		
WF(g)	~86 ³⁴⁰			WSi ₂ (c)	-22 ¹⁶⁶	-21	15 ¹⁶⁶
WF ₂ (g)	~-25 ³⁴⁰			W ₅ Si ₃ (c)	-32 ¹⁶⁶	-33	55 ¹⁶⁶
WF ₃ (g)	~-127 ³⁴⁰			WTe ₂ (c)	-15.3 ³⁶¹	-12.1	20.9 ³⁶¹
WF ₄ (g)	-246 ³⁴⁰			Na _{0.53} WO ₃ (c)	-232.5 ³⁶⁴		
WF ₅ (c)	-345.9 ³³⁶			Na _{0.679} WO ₃ (c)	-240.9 ³⁶⁴	-221.0	22.77 ³⁶⁴
(WF ₅) ₄ (g)	-1316 ³³⁹			Na _{0.77} WO ₃ (c)	-245.9 ³⁶⁴		
WF ₆ (liq)	-417.7 ^{240,337}	-390.0	60.7 ^{240,337}	H _{0.18} WO ₃ (c)	-202.6 ³⁶⁵		
WF ₆ (g)	-411.5 ³³⁵	-390.1	81.49 ¹⁵¹	H _{0.35} WO ₃ (c)	-203.7 ³⁶⁵		
WOF ₄ (c)	-359 ³³⁸	-333	(42) ¹⁵¹	(NH ₄) _{0.25} WO ₃ (c)	-224.4 ³⁶⁵		
WOF ₄ (g)	~-316 ²⁴⁸	~-301	(80.4) ¹⁵¹				

^a Italicized values agree with values in NBS 270-4 or NBS 270-6. Values in parentheses are estimates. ^b Oppermann and Stöver³⁴⁸ have estimated $S^\circ = 62 \pm 6$ cal K⁻¹ mol⁻¹ for WBr₄(c) and $S^\circ = 112 \pm 7$ cal K⁻¹ mol⁻¹ for WBr₄(g). We have adopted their entropy of sublimation of WBr₄(c), but not their entropy values. These seem to be too high, compared with the estimated entropies of other related compounds.

tables¹⁵¹ and have not been adopted here.

The existence of W₃O(c) is contested^{297,303} and we do not include any data reported³⁰³ for W₃O(c) in our Table V.


For the compound WO₃·H₂O(c), sometimes written as H₂WO₄(c), the standard enthalpy of solution in aqueous sodium hydroxide has been determined as -13.7 and -13.2 kcal mol⁻¹, respectively.^{304,305} We combine the least exothermic heat of solution³⁰⁵ with our adopted ΔH_f° for WO₄²⁻(aq), which we will discuss later, to obtain $\Delta H_f^\circ = -270.0$ kcal mol⁻¹ for WO₃·H₂O(c). We estimate $S^\circ = (33)$ cal K⁻¹

mol⁻¹. With these values, a vapor pressure of H₂O(g) over WO₃·H₂O(c) and WO₃(c) of about 40 mmHg can be calculated. This vapor pressure "should" be of the same order of magnitude as for MoO₃·H₂O(c), ~1 mmHg. In the absence of reliable vapor pressure data and third law entropies for MoO₃·H₂O(c) and WO₃·H₂O(c), all we can say is that we suspect minor errors in the thermodynamic data for WO₄²⁻(aq) and WO₃·H₂O(c). We note that use of the more exothermic enthalpy of solution³⁰⁴ leads to a still higher calculated vapor pressure.

For $\text{WO}_3\text{-H}_2\text{O}(\text{g})$ we adopt $\Delta H_f^\circ \simeq -217 \text{ kcal mol}^{-1}$ from equilibrium measurements by Belton and McCarron.³⁰⁶ This value is in good agreement with the results of Glemser et al.³⁰⁷ The $\Delta H_f^\circ = -229 \text{ kcal mol}^{-1}$ in NBS 270-4 is consistent with the earlier results of Meyer et al.³⁰⁸

G. $\text{WO}_4^{2-}(\text{aq})$ and Tungstates

The most direct way to determine ΔH_f° of $\text{WO}_4^{2-}(\text{aq})$ is to measure the enthalpy of solution of $\text{WO}_3(\text{c})$ in excess base. This is, however, beset with experimental difficulties. Two different investigations of the enthalpy of solution of $\text{WO}_3(\text{c})$ in aqueous sodium hydroxide lead to $\Delta H_f^\circ = -257.1 \text{ kcal mol}^{-1}$ (ref 305), as adopted in NBS 270-4, and to $\Delta H_f^\circ = -256.5 \text{ kcal mol}^{-1}$ (ref 304), respectively, for $\text{WO}_4^{2-}(\text{aq})$. For reasons discussed in ref 304, we accept this latter value. We also note that Sherfey and Brenner³⁰⁹ employed electrochemical calorimetry to obtain $\Delta H^\circ = -7.5 \pm 1.5 \text{ kcal}$ for the reaction



and thence $\Delta H_f^\circ \simeq -253.6 \text{ kcal mol}^{-1}$ for $\text{WO}_4^{2-}(\text{aq})$.

Whereas several routes were available to calculate thermodynamic properties of $\text{MoO}_4^{2-}(\text{aq})$, similar data for tungstates are comparatively scarce, and we are unable to present any reliable experimental values for ΔG_f° or S° for $\text{WO}_4^{2-}(\text{aq})$. From $K_{\text{sp}} = 1 \times 10^{-8}$ for $\text{CaWO}_4(\text{c})$ ⁷⁵ and the ΔG_f° for $\text{CaWO}_4(\text{c})$ to be discussed later, we calculate $\Delta G_f^\circ = -223.5 \text{ kcal mol}^{-1}$ for $\text{WO}_4^{2-}(\text{aq})$ and thence $S^\circ = 26 \text{ cal K}^{-1} \text{ mol}^{-1}$ for this ion. As this seems to be an unreasonably high entropy value, we prefer to estimate $S^\circ = (11) \text{ cal K}^{-1} \text{ mol}^{-1}$ and combine with the above ΔH_f° to find $\Delta G_f^\circ = -219 \text{ kcal mol}^{-1}$ for $\text{WO}_4^{2-}(\text{aq})$.

An alkaline tungstate solution contains monomeric $\text{WO}_4^{2-}(\text{aq})$, but on acidification complicated polymerization reactions occur. These reactions are not yet well understood. Equilibrium constants for postulated reactions are to be found in Sillén's compilations;^{75,112} Kepert³¹⁰ has reviewed the literature up to 1961 and some later work has been discussed by Cotton and Wilkinson.²⁰

Two recent investigations^{311,312} are in general agreement on the equilibrium constant for the formation of the "paratungstate A" ion, $\text{HW}_6\text{O}_{21}^{5-}(\text{aq})$, for which the enthalpy of formation is also known.³¹³ However, these investigators^{311,312} have come to quite different conclusions regarding other thermodynamic and kinetic characteristics of tungstate solutions, clearly showing that our knowledge of the properties of these solutions is far from complete. We therefore limit our listings in Table V to $\text{HW}_6\text{O}_{21}^{5-}(\text{aq})$. Stopped-flow kinetic experiments³¹⁴ have shown that $\text{HW}_6\text{O}_{21}^{5-}(\text{aq})$ is formed rapidly (within $\sim 1 \text{ s}$) upon acidification of a tungstate solution and that the intermediate species are not well defined.

Further acidification of solutions containing polytungstate ions leads to depolymerization reactions in which cationic species are formed. Sillén^{75,112} quotes some equilibrium constants for such reactions; Nazarenko and Poluektova³¹⁵ have investigated the dissociation constant of tungstic acid in solutions of varying ionic strength.

Sillén^{75,112} cites an impressive number of determinations of solubility products and ionization constants for various heteropolytungstates. We call attention to recent determinations of reduction potentials,³¹⁶ an acid dissociation constant,³¹⁷ and heats of solution and neutralization^{318,319} for some heteropolytungstates. Weakley³²⁰ has described the preparation of some polytungstate ions containing two heteroatoms and has also measured some redox potentials and stability constants related to such species.

O'Hare et al.²⁰⁹ cite $\Delta H_f^\circ = -383.2 \text{ kcal mol}^{-1}$ for

$\text{Li}_2\text{WO}_4(\text{c})$, but we cannot assess the reliability of this value.

King and Weller³²¹ have made C_p measurements on $\text{Na}_2\text{WO}_4(\text{c})$, leading to $S^\circ = 38.5 \pm 0.5 \text{ cal K}^{-1} \text{ mol}^{-1}$ as adopted in our Table V. Another extrapolation of the specific heat from 51 to 0 K has led to¹⁵¹ $S^\circ = 38.32 \text{ cal K}^{-1} \text{ mol}^{-1}$. The calorimetric measurements of Koehler et al.²¹⁵ [recalculated with our adopted ΔH_f° for $\text{H}_2\text{WO}_4(\text{c})$] lead to $\Delta H_f^\circ = -369.5 \text{ kcal mol}^{-1}$ for $\text{Na}_2\text{WO}_4(\text{c})$, and those of Graham and Hepler³²² give $\Delta H_f^\circ = -369.7 \text{ kcal mol}^{-1}$. We adopt the mean value, $\Delta H_f^\circ = -369.6 \text{ kcal mol}^{-1}$, for $\text{Na}_2\text{WO}_4(\text{c})$.

The ΔH_f° values listed in Table V for $\text{Na}_2\text{W}_2\text{O}_7(\text{c})$ and $\text{Na}_2\text{W}_4\text{O}_{13}(\text{c})$ are recalculated by us from the work of Koehler et al.²¹⁵ Our ΔG_f° for $\text{Na}_2\text{W}_2\text{O}_7(\text{c})$ is calculated from the adopted ΔH_f° and the third law entropy,^{218a} $S^\circ = 60.8 \text{ cal K}^{-1} \text{ mol}^{-1}$.

We cannot account for any experimental values of ΔH_f° for $\text{K}_2\text{WO}_4(\text{c})$, $\text{Rb}_2\text{WO}_4(\text{c})$, or $\text{Cs}_2\text{WO}_4(\text{c})$. O'Hare et al.²⁰⁹ cite $\Delta H_f^\circ = -377.9 \text{ kcal mol}^{-1}$ for $\text{K}_2\text{WO}_4(\text{c})$ and have reported estimated ΔH_f° values for $\text{Rb}_2\text{WO}_4(\text{c})$ and $\text{Cs}_2\text{WO}_4(\text{c})$.

Yamdagni, Pupp, and Porter²¹⁷ have reported the value we list for ΔH_f° (at 0 K) for $\text{Li}_2\text{WO}_4(\text{g})$, while the values listed for ΔH_f° for $\text{K}_2\text{WO}_4(\text{g})$ and $\text{KHWO}_4(\text{g})$ are due to Farber and Srivastava.²²⁰

The NBS 270-6 gives $\Delta H_f^\circ = -366.3 \text{ kcal mol}^{-1}$ and $S^\circ = 24.18 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{MgWO}_4(\text{c})$. The entropy is from C_p measurements by King and Weller.³²¹ The high-temperature emf data of Rezukhina and Levitskii³²³ give $\Delta H_f^\circ = -362.3 \text{ kcal mol}^{-1}$ (third law analysis¹⁵¹) or $\Delta H_f^\circ = -372.7 \text{ kcal mol}^{-1}$ (second law analysis¹⁵¹). Our recalculation of calorimetric data from Navrotsky and Kleppa³²⁴ leads to $\Delta H_f^\circ = -362.7 \text{ kcal mol}^{-1}$. We consider this last value to be the best one and adopt it in Table V. Combination of this ΔH_f° with the S° leads to our ΔG_f° .

For $\text{CaWO}_4(\text{c})$, $S^\circ = 30.21 \text{ cal K}^{-1} \text{ mol}^{-1}$ is listed in NBS 270-6, in close agreement with the value from King and Weller.³²¹ Barany's calorimetric data²¹¹ permit us to calculate $\Delta H_f^\circ = -392.2 \text{ kcal mol}^{-1}$ for $\text{CaWO}_4(\text{c})$. The ΔH_f° value in NBS 270-6 is 1 kcal mol⁻¹ more negative, partly due to different choices of ΔH_f° for $\text{H}_2\text{WO}_4(\text{c})$. From our ΔH_f° value and the S° value we obtain $\Delta G_f^\circ = -366.7 \text{ kcal mol}^{-1}$. The solubility product for $\text{CaWO}_4(\text{c})$ is then calculated to be $K_{\text{sp}} = 4 \times 10^{-12}$, as compared to an old value,⁷⁵ $K_{\text{sp}} = 1 \times 10^{-8}$, based on conductance measurements.

NBS 270-6 lists $\Delta H_f^\circ = -391.9 \text{ kcal mol}^{-1}$, $\Delta G_f^\circ = -366 \text{ kcal mol}^{-1}$, and $S^\circ = 33 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{SrWO}_4(\text{c})$. The ΔG_f° value is the same as reported by Levitskii and Rezukhina³²⁵ from high-temperature emf data, but these latter authors report $\Delta H_f^\circ = -393.5 \text{ kcal mol}^{-1}$ and $S^\circ = 28.3 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{SrWO}_4(\text{c})$. This latter entropy seems to be unreasonably small. As the experimental ΔG_f° is more likely to be correct than the derived ΔH_f° , we follow NBS 270-6 and adopt $\Delta G_f^\circ = -366 \text{ kcal mol}^{-1}$, $S^\circ = (33) \text{ cal K}^{-1} \text{ mol}^{-1}$, and calculate $\Delta H_f^\circ \simeq -391 \text{ kcal mol}^{-1}$.

For $\text{BaWO}_4(\text{c})$, NBS 270-6 lists $\Delta H_f^\circ = -407 \text{ kcal mol}^{-1}$, which is consistent with the results of Tammann and Westergold, as cited by Bichowsky and Rossini.⁴⁵

The NBS 270-6 lists ΔH_f° (at 0 K) for $\text{MgWO}_4(\text{g})$, $\text{CaWO}_4(\text{g})$, and $\text{SrWO}_4(\text{g})$, while NBS 270-4 lists ΔH_f° (at 0 K) for $\text{SnWO}_4(\text{g})$ and $\text{Sn}_2\text{WO}_5(\text{g})$. The values are all consistent with the results of Verhaegen et al.²²² We have recalculated their results, using our properties of $\text{WO}_2(\text{g})$ and $\text{WO}_3(\text{g})$, and list these recalculated values in Table V. The uncertainty in these ΔH_f° values is about $\pm 10 \text{ kcal mol}^{-1}$.

Our calculations with the high-temperature data of Rezukhina et al.^{326,327} lead to $\Delta H_f^\circ \simeq -317 \text{ kcal mol}^{-1}$ for $\text{MnWO}_4(\text{c})$. Two different calorimetric investigations³²⁸ lead to $\Delta H_f^\circ = -311.9 \text{ kcal mol}^{-1}$ for $\text{MnWO}_4(\text{c})$, which is the value listed in NBS 270-4 and adopted for our Table V.

The ΔH_f° values listed in NBS 270-4 for $\text{FeWO}_4(\text{c})$, $\text{FeWO}_4 \cdot$


$3\text{H}_2\text{O}(c)$, and $\text{Fe}_2(\text{WO}_4)_3 \cdot 8\text{H}_2\text{O}(c)$ are apparently derived from old measurements by Tammann and Westerhold, as cited by Bichowsky and Rossini.⁴⁵ We adopt $\Delta H_f^\circ = -276 \text{ kcal mol}^{-1}$ for $\text{FeWO}_4(c)$ in Table V, but do not include ΔH_f° values for the other two compounds. The entropy of $\text{FeWO}_4(c)$ has recently been determined by Lyon and Westrum.³³⁰ We adopt their $S^\circ = 31.61 \text{ cal K}^{-1} \text{ mol}^{-1}$, in good agreement with the value in NBS 270-4, $S^\circ = 31.5 \text{ cal K}^{-1} \text{ mol}^{-1}$, which is from Weller.²²⁷

The enthalpies of formation of $\text{CoWO}_4(c)$, $\text{NiWO}_4(c)$, $\text{CuWO}_4(c)$, $\text{ZnWO}_4(c)$, and $\text{CdWO}_4(c)$ can all be calculated from the measured heats of reaction of the corresponding oxides with $\text{WO}_3(c)$ at 970 K.³²⁴ We have recalculated ΔH_f° for $\text{CoWO}_4(c)$, using C_p data from Yakovleva and Rezukhina,³²⁷ and for the other compounds estimated that the enthalpy of reaction is 0.2 kcal mol⁻¹ less negative at 298 K than it is at 970 K to obtain the ΔH_f° values listed in our Table V. We note that the data from Tammann and Westerhold (as cited by Bichowsky and Rossini⁴⁵) lead to $\Delta H_f^\circ = -251 \text{ kcal mol}^{-1}$ for $\text{CuWO}_4(c)$ instead of $-245.5 \text{ kcal mol}^{-1}$ as chosen by us. NBS 270-4 lists $\Delta H_f^\circ = -264 \text{ kcal mol}^{-1}$ for $\text{CuWO}_4(c)$, from a source unknown to us. An older calorimetric³²⁸ ΔH_f° for $\text{NiWO}_4(c)$ is 1.8 kcal mol⁻¹ more negative than our adopted value.

The value in Table V for ΔH_f° of $\text{ZnWO}_4(c)$ agrees well with the value deduced from high-temperature equilibrium data, as discussed by Lyon and Westrum.³²⁹ The C_p measurements of Lyon and Westrum³³⁰ on $\text{ZnWO}_4(c)$ lead to $S^\circ = 28.34 \pm 0.1 \text{ cal K}^{-1} \text{ mol}^{-1}$. From the calculated $\Delta G_f^\circ = -268.5 \text{ kcal mol}^{-1}$, the solubility product of $\text{ZnWO}_4(c)$ is found to be $K_{sp} = 2.5 \times 10^{-11}$. We are not aware of any experimental determination of this quantity.

For $\text{PbWO}_4(c)$ the NBS 270-4 lists $S^\circ = 40.2 \text{ cal K}^{-1} \text{ mol}^{-1}$, which is the third law value found by Weller and Kelley.²²⁸ The $\Delta H_f^\circ = -277.0 \text{ kcal mol}^{-1}$ for $\text{PbWO}_4(c)$ has been reported by Zharkova and Gerasimov³³¹ on the basis of earlier high-temperature equilibrium results. The solubility product is³³² $K_{sp} = 8.6 \times 10^{-17}$. We cannot assess the reliability of any of these values. They lead to an improbable $S^\circ = 32 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{WO}_4^{2-}(aq)$ [using the quoted S° for $\text{PbWO}_4(c)$ and data for $\text{Pb}^{2+}(aq)$ from NBS 270-3]. If we take our adopted $S^\circ = (11) \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{WO}_4^{2-}(aq)$ and $\Delta H_f^\circ = -277.0 \text{ kcal mol}^{-1}$ for $\text{PbWO}_4(c)$, we calculate $K_{sp} = 2.5 \times 10^{-21}$ and $\Delta H^\circ(\text{ppt}) = -20.1 \text{ kcal mol}^{-1}$. If we instead begin with $K_{sp} = 8.6 \times 10^{-17}$, then $\Delta H_f^\circ = -270.8 \text{ kcal mol}^{-1}$ and $\Delta H^\circ(\text{ppt}) = -13.9 \text{ kcal mol}^{-1}$ are calculated. As this last value is closer to the heats of precipitation of $\text{PbMoO}_4(c)$, $\text{CdWO}_4(c)$, $\text{ZnWO}_4(c)$, and $\text{CuWO}_4(c)$ than the more exothermic heat of precipitation is, we hesitantly adopt $\Delta H_f^\circ = -271 \text{ kcal mol}^{-1}$ for $\text{PbWO}_4(c)$, noting that the discrepancies discussed here make a calorimetric determination of the ΔH_f° for $\text{PbWO}_4(c)$ desirable.

The enthalpy of solution of $\text{Ag}_2\text{WO}_4(c)$ has been determined as 14.7 kcal mol⁻¹ from calorimetric measurements³²² (in which the experimental conditions were chosen to avoid the formation of polytungstates), and as 12.1 kcal mol⁻¹ from the temperature variation of the solubility product.³³³ We adopt the former heat of solution, and estimate $S^\circ = (53) \text{ cal K}^{-1} \text{ mol}^{-1}$. We can then calculate for $\text{Ag}_2\text{WO}_4(c)$: $\Delta H_f^\circ = -220.7 \text{ kcal mol}^{-1}$, $\Delta G_f^\circ = -198.9 \text{ kcal mol}^{-1}$, $K_{sp} = 4 \times 10^{-13}$, and also



Pan's³³³ electrochemical measurements have led to $K_{sp} = 5 \times 10^{-12}$ and $E^\circ = 0.466 \text{ V}$. The differences between the values calculated by us and the experimental values exceed the stated uncertainties. A third law entropy for $\text{Ag}_2\text{WO}_4(c)$ is needed.

The free energies of formation of lanthanum and neodym-

ium tungstates have been investigated by emf methods at high temperatures.³³⁴ We cannot make calculations to relate the published data to our standard state at 298 K.

H. Halides and Oxyhalides of Tungsten

Fluorine combustion calorimetry by O'Hare and Hubbard³³⁵ has led to $\Delta H_f^\circ = -411.5 \text{ kcal mol}^{-1}$ for $\text{WF}_6(g)$. This value, which is in excellent agreement with the results of two more recent calorimetric investigations,³³⁶ is adopted in NBS 270-4 and our Table V. We also follow NBS 270-4 in adopting $S^\circ = 81.49 \text{ cal K}^{-1} \text{ mol}^{-1}$ from statistical calculations cited in the JANAF tables.¹⁵¹ Our calculated $\Delta G_f^\circ = -390.1 \text{ kcal mol}^{-1}$ is consistent with the listed ΔH_f° and S° values.

The NBS 270-4 gives $\Delta H_f^\circ = -417.7 \text{ kcal mol}^{-1}$ and $S^\circ = 60.1 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{WF}_6(\text{liq})$, values that are consistent with the above ΔH_f° and vapor pressures from Cady et al.^{240,337} We adopt these values, which lead to $\Delta G_f^\circ = -390.0 \text{ kcal mol}^{-1}$ for $\text{WF}_6(\text{liq})$.

The mass spectrometric investigation by Zmbov, Uy, and Margrave²⁴⁸ leads to our adopted $\Delta H_f^\circ \approx -215 \text{ kcal mol}^{-1}$ for $\text{WO}_2\text{F}_2(g)$ and $\Delta H_f^\circ \approx -316 \text{ kcal mol}^{-1}$ for $\text{WOF}_4(g)$. Burgess et al.³³⁸ have reported a calorimetric $\Delta H_f^\circ = -359 \text{ kcal mol}^{-1}$ for $\text{WOF}_4(c)$. The enthalpy of sublimation of $\text{WOF}_4(c)$ reported by Cady and Hargreaves,²⁴⁷ $\Delta H^\circ(\text{subl}) = 17.0 \text{ kcal mol}^{-1}$, cannot be used to calculate ΔH_f° for $\text{WOF}_4(g)$, as it has been shown²⁴⁶ that oligomeric species are formed in the gas phase over $\text{WOF}_4(c)$. We combine the heats of formation with estimated entropies to calculate the free energies of formation in Table V.

Calorimetric measurements³³⁶ lead to $\Delta H_f^\circ = -345.9 \text{ kcal mol}^{-1}$ for $\text{WF}_5(c)$ on the basis of our adopted value of ΔH_f° for $\text{WF}_6(g)$. The enthalpy of sublimation has been determined,³³⁹ leading to our adopted $\Delta H_f^\circ = -1316 \text{ kcal mol}^{-1}$ for $(\text{WF}_5)_4(g)$.

Dittmer et al.³⁴⁰ have made equilibrium experiments at various temperatures leading to ΔH_f° values for the lower valent gaseous tungsten fluorides. The enthalpy of formation of $\text{WF}_4(g)$ is probably determined with good accuracy from these experiments, while the uncertainties in the values we list for $\text{WF}(g)$, $\text{WF}_2(g)$, and $\text{WF}_3(g)$ probably are considerable.

The enthalpies of formation listed in Table V for the tungsten chlorides are mainly derived from measurements by Shchukarev et al.²⁴⁹ Solution calorimetry leads to $\Delta H_f^\circ = -142.9 \text{ kcal mol}^{-1}$ for $\text{WCl}_6(c,\alpha)$, based on our adopted ΔH_f° for $\text{WO}_3 \cdot \text{H}_2\text{O}(c)$. Three investigations, cited in the JANAF tables,¹⁵¹ are in good agreement with the enthalpy of sublimation of $\text{WCl}_6(c,\alpha)$ and thence $\Delta H_f^\circ = -118.0 \text{ kcal mol}^{-1}$ for $\text{WCl}_6(g)$.

The uncertain ΔH_f° values listed in Table V for $\text{WCl}_5(c)$, $\text{WCl}_5(g)$, $(\text{WCl}_5)_2(g)$, $\text{WCl}_4(c)$, and $\text{WCl}_4(g)$ are all derived from the equilibrium measurements by Shchukarev et al.²⁴⁹ while ΔH_f° for $\text{WCl}_2(c)$ is from combustion calorimetry.²⁴⁹

There are two calorimetric determinations of the enthalpy of reaction of $\text{WO}_2\text{Cl}_2(c)$ with aqueous hydroxide. We calculate $\Delta H_f^\circ = -187.0 \text{ kcal mol}^{-1}$ from Shchukarev's data²⁴⁹ and $\Delta H_f^\circ = -190.6 \text{ kcal mol}^{-1}$ from Oppermann's data.³⁴¹ We adopt $\Delta H_f^\circ = -189 \text{ kcal mol}^{-1}$ for $\text{WO}_2\text{Cl}_2(c)$. The enthalpy of sublimation is 26 kcal mol⁻¹ (at 298 K) according to a third law recalculation of the data of Shchukarev and Suvorov as cited in the JANAF tables¹⁵¹ or 28 kcal mol⁻¹ according to Oppermann.³⁴¹ We adopt the mean value and then obtain $\Delta H_f^\circ = -162 \text{ kcal mol}^{-1}$ for $\text{WO}_2\text{Cl}_2(g)$.

For $\text{WOCl}_4(c)$ the NBS 270-4 lists $\Delta H_f^\circ = -161.7 \text{ kcal mol}^{-1}$, in good agreement with the results of Shchukarev et al.²⁴⁹ Oppermann's data³⁴¹ lead to $\Delta H_f^\circ = -172.4 \text{ kcal mol}^{-1}$, which we adopt, as this value seems to be more reasonable in relation to the ΔH_f° of $\text{MoOCl}_4(c)$.

It has been observed that the enthalpy of sublimation of

$\text{WOCl}_4(\text{c})$ varies considerably with the temperature,³⁴² which might partly explain the difference between the value we calculate from the data of Enghag and Staffansson³⁴² [$\Delta H^\circ(\text{sublimation}) = 19.0 \text{ kcal mol}^{-1}$] and that calculated¹⁵¹ from the data of Shchukarev and Suvorov [$\Delta H^\circ(\text{subl}) = 23.4 \text{ kcal mol}^{-1}$]. We weight the more recent value more heavily, adopt $\Delta H^\circ(\text{subl}) = 20.0 \text{ kcal mol}^{-1}$, and calculate $\Delta H_f^\circ = -152.4 \text{ kcal mol}^{-1}$ for $\text{WOCl}_4(\text{g})$.

An apparently reliable $S^\circ = 92.1 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{WOCl}_4(\text{g})$ has been calculated from spectroscopically determined molecular parameters.³⁴³


Table V lists $\Delta H_f^\circ = -165.9 \text{ kcal mol}^{-1}$ for $\text{WOCl}_3(\text{c})$ from calorimetric measurements by Oppermann, Stöver, and Kunze³⁴⁴ on the reaction between $\text{WOCl}_3(\text{c})$ and hydrogen peroxide in alkaline solution. We have used our adopted ΔH_f° value for $\text{Na}_2\text{WO}_4(\text{c})$ in this calculation, as well as for the corresponding calculations for $\text{WO}_2\text{Cl}_2(\text{c})$ and $\text{WOCl}_4(\text{c})$ described above.

Oppermann et al.³⁴⁴ also calculate $\Delta H^\circ = 32.5 \text{ kcal mol}^{-1}$ of $\text{WOCl}_2(\text{c})$ for the reaction between $\text{WOCl}_3(\text{c})$ and $\text{WOCl}_2(\text{c})$ to form $\text{WOCl}_4(\text{g})$, whence $\Delta H_f^\circ = -147 \text{ kcal mol}^{-1}$ for $\text{WOCl}_2(\text{c})$. This equilibrium study also leads to $S^\circ = 110 \pm 5 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{WOCl}_4(\text{g})$, in substantial disagreement with the value listed above, which casts suspicion also on the accuracy of the value quoted for ΔH_f° of $\text{WOCl}_2(\text{c})$.


Burgess et al.³³⁸ have reported enthalpies of alkaline hydrolysis of $\text{WF}_5\text{Cl}(\text{liq})$ and $\text{WF}_4\text{Cl}_2(\text{liq})$, from which we calculate the ΔH_f° values listed in Table V for these compounds.

Shchukarev and Kokovin³⁴⁵ have made calorimetric measurements on the enthalpy of solution of $\text{WBr}_6(\text{c})$ and $\text{WBr}_5(\text{c})$, leading to our adopted ΔH_f° values. The values in NBS 270-4 for these compounds are apparently also derived from this source. Shchukarev et al.³⁴⁶ have studied the sublimation of $\text{WBr}_5(\text{c})$. Our interpretation of their results leads to $\Delta H_f^\circ = -52 \text{ kcal mol}^{-1}$ for $\text{WBr}_5(\text{g})$. The uncertainty in this value overlaps the value given in NBS 270-4, $\Delta H_f^\circ = -55.4 \text{ kcal mol}^{-1}$.

Two calorimetric investigations^{347,348} are in good agreement with $\Delta H_f^\circ = -170.2 \text{ kcal mol}^{-1}$ for $\text{WO}_2\text{Br}_2(\text{c})$. The enthalpy of sublimation,³⁴⁸⁻³⁵⁰ $36.5 \pm 1.0 \text{ kcal mol}^{-1}$, leads to $\Delta H_f^\circ = -133.7 \text{ kcal mol}^{-1}$ for $\text{WO}_2\text{Br}_2(\text{g})$. Gupta³⁴⁹ has studied the equilibria



and



with results that lead to $\Delta H_f^\circ = -131 \pm 2 \text{ kcal mol}^{-1}$ for $\text{WO}_2\text{Br}_2(\text{g})$. We adopt $\Delta H_f^\circ = -133.0 \text{ kcal mol}^{-1}$ for $\text{WO}_2\text{Br}_2(\text{g})$. The entropy of $\text{WO}_2\text{Br}_2(\text{g})$ has been calculated by Gupta³⁴⁹ from estimated molecular parameters.

From the heat of reaction of $\text{WOBBr}_4(\text{c})$ with aqueous hydroxide as reported by Oppermann and Stöver³⁴⁸ and by Shchukarev et al.,³⁴⁷ we obtain $\Delta H_f^\circ = -131.8 \text{ kcal mol}^{-1}$ for $\text{WOBBr}_4(\text{c})$. The value listed in NBS 270-4 is consistent with the results of the latter investigation.³⁴⁷ We adopt $\Delta H^\circ(\text{subl}) = 30.0 \pm 0.5 \text{ kcal mol}^{-1}$ as an average of three investigations³⁴⁸⁻³⁵⁰ to obtain $\Delta H_f^\circ = -101.8 \text{ kcal mol}^{-1}$ for $\text{WOBBr}_4(\text{g})$.

One calorimetric investigation by Oppermann et al.³⁵¹ leads to $\Delta H_f^\circ = -126.0 \text{ kcal mol}^{-1}$ for $\text{WOBBr}_3(\text{c})$, as listed in Table V. They³⁵¹ have also studied the decomposition of $\text{WOBBr}_3(\text{c})$ and calculated ΔH_f° for $\text{WOBBr}_2(\text{c})$ from their data. As it has been suggested³⁵² that their measurements refer to nonequilibrium conditions, we have not included thermodynamic properties of $\text{WOBBr}_2(\text{c})$ in Table V.

The reaction between $\text{W}(\text{CO})_6$ and l_2 at 530 K has been studied calorimetrically by Virmani, Barnes, and Skinner.²⁶³

The values of ΔH_f° for $\text{Wl}_2(\text{c})$ and $\text{Wl}_3(\text{c})$ listed in Table V are derived from their results, using our adopted ΔH_f° for $\text{W}(\text{CO})_6(\text{c})$, which we discuss later. As the products formed in the calorimetric reaction are not well defined, the uncertainties in the listed ΔH_f° values are fairly large ($\sim 3 \text{ kcal mol}^{-1}$).

Gupta^{349,353} has investigated the thermal stabilities of tungsten oxyiodides and discussed earlier work. On the basis of his results and earlier data quoted by him, we adopt $\Delta H_f^\circ = -145.4 \text{ kcal mol}^{-1}$ for $\text{WO}_2\text{l}_2(\text{c})$, $\Delta H_f^\circ = -102.3 \text{ kcal mol}^{-1}$ for $\text{WO}_2\text{l}_2(\text{g})$, and $S^\circ = (89) \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{WO}_2\text{l}_2(\text{g})$. The entropy is calculated from estimated molecular parameters.³⁵³ The measurements also permit calculation of $\Delta H_f^\circ = -137 \pm 5 \text{ kcal mol}^{-1}$ for $\text{WO}_2\text{l}(\text{c})$.


I. Other Tungsten Compounds

The enthalpies of formation of the gaseous calcium and strontium tungstites can be deduced from the mass spectrometric results of Verhaegen et al.²²² We list ΔH_f° (at 0 K) $\simeq -118 \text{ kcal mol}^{-1}$ for $\text{CaWO}_3(\text{g})$, but omit $\text{SrWO}_3(\text{g})$, as the values for this compound seem to be erroneous.

The enthalpies of formation of $\text{KWF}_6(\text{c})$, $\text{KWCl}_6(\text{c})$, and $\text{K}_2\text{WCl}_7(\text{c})$, listed in our Table V, have been determined calorimetrically.^{354,355} Kudryashova et al.³⁵⁶ have reported enthalpies of combustion and alkaline oxidative hydrolysis for the compounds $\text{M}_3\text{W}_2\text{Cl}_9(\text{c})$, where $\text{M} = \text{K}, \text{Rb}, \text{or Cs}$. We calculate $\Delta H_f^\circ = -524 \text{ kcal mol}^{-1}$ or $-536 \text{ kcal mol}^{-1}$ for $\text{K}_3\text{W}_2\text{Cl}_9(\text{c})$ from the two experiments, using our adopted ΔH_f° values for $\text{K}_2\text{WO}_4(\text{c})$, $\text{WCl}_5(\text{c})$, and $\text{WCl}_6(\text{c})$. Our listed ΔH_f° values for the other two compounds are calculated from the results of the hydrolysis experiments. Although it is possible to use the bomb calorimetric results for calculation of ΔH_f° values for $\text{Rb}_2\text{WO}_4(\text{c})$ and $\text{Cs}_2\text{WO}_4(\text{c})$, the discordant ΔH_f° values listed above for $\text{K}_3\text{W}_2\text{Cl}_9(\text{c})$ indicate that the errors are too great to make such calculated ΔH_f° values useful.

Our Table V also includes ΔH_f° values for $\text{WF}_5(\text{OMe})(\text{c})$, *cis*- $\text{WF}_4(\text{OMe})_2(\text{liq})$, and *cis*- $\text{WF}_2(\text{OMe})_4(\text{c})$, where OMe denotes the methoxide group. These enthalpies have been determined via alkaline hydrolysis.³³⁸

Emf measurements³⁵⁷ on the equilibrium



have led to $\Delta H_f^\circ = -54 \text{ kcal mol}^{-1}$ and $S^\circ = 29 \text{ cal K}^{-1} \text{ mol}^{-1}$ for $\text{WS}_2(\text{c})$.

The enthalpies of formation of the tungsten carbides have been determined by combustion calorimetry. Our adopted $\Delta H_f^\circ = -6.3 \text{ kcal mol}^{-1}$ for $\text{W}_2\text{C}(\text{c})$ is from Mah,²⁷³ while $\Delta H_f^\circ = -9.6_3 \text{ kcal mol}^{-1}$ for $\text{WC}(\text{c})$ is the mean of the values from Mah²⁷³ and from McGraw, Seltz, and Snyder.³⁵⁸

For $\text{W}(\text{CO})_6(\text{c})$ we adopt the same value as listed in NBS 270-4, $\Delta H_f^\circ = -227.9 \text{ kcal mol}^{-1}$, calculated from the heat of combustion of $\text{W}(\text{CO})_6(\text{c})$ reported by Cotton, Fischer, and Wilkinson.¹⁷³ These authors¹⁷³ also quote from earlier measurements the enthalpy of sublimation, $17.7 \text{ kcal mol}^{-1}$. We calculate $\Delta H_f^\circ = -210.2 \text{ kcal mol}^{-1}$ for $\text{W}(\text{CO})_6(\text{g})$. We are not able to account for the value ($-208.3 \text{ kcal mol}^{-1}$) listed in NBS 270-4, which corresponds to $19.6 \text{ kcal mol}^{-1}$ for the enthalpy of sublimation.

Virmani, Barnes, and Skinner²⁶³ have recently proposed $\Delta H_f^\circ = -229.0 \pm 0.8 \text{ kcal mol}^{-1}$ for $\text{W}(\text{CO})_6(\text{c})$. This value has not been considered here, as it is based on calorimetric work that (as far as we know) is not yet published.

The most important W(IV) complex is the stable octacyanide, $\text{W}(\text{CN})_8^{4-}$. Baadsgaard and Treadwell³⁵⁹ have made measurements leading to



- Chemical Thermodynamic Properties", National Bureau of Standards Technical Note 270-6, U.S. Government Printing Office, Washington, D.C., 1971.
- (6) R. N. Goldberg and L. G. Hepler, *Chem. Rev.*, **68**, 229 (1968).
 - (7) I. M. Klotz and R. M. Rosenberg, "Chemical Thermodynamics", 3d ed, W. A. Benjamin, Menlo Park, Calif., 1972; G. N. Lewis and M. Randall (revised by K. S. Pitzer and L. Brewer), "Thermodynamics", McGraw-Hill, New York, N.Y., 1961.
 - (8) E. A. Heintz, *J. Inorg. Nucl. Chem.*, **21**, 262 (1961).
 - (9) B. D. Nash, T. T. Campbell, and F. E. Block, Bureau of Mines Report of Investigations No. 7112, U.S. Government Printing Office, Washington, D.C., 1968.
 - (10) I. Bernal and S. E. Harrison, *J. Chem. Phys.*, **34**, 102 (1961).
 - (11) L. Petrakis, *J. Phys. Chem.*, **66**, 433 (1962).
 - (12) A. J. Deyrup, *Inorg. Chem.*, **3**, 1645 (1964).
 - (13) R. L. de Beauchamp and T. A. Sullivan, Bureau of Mines Report of Investigations No. 7194, U.S. Government Printing Office, Washington, D.C., 1968.
 - (14) E. Kyuno, M. Kamada, and N. Tanaka, *Bull. Chem. Soc. Jpn.*, **40**, 1848 (1967).
 - (15) T. W. Swaddle, J. H. Lipton, G. Guastalla, and P. Bayliss, *Can. J. Chem.*, **49**, 2433 (1971).
 - (16) N. C. Tombs, W. J. Croft, J. R. Carter, and J. F. Fitzgerald, *Inorg. Chem.*, **3**, 1791 (1964); H. C. Clark and Y. N. Sadana, *Can. J. Chem.*, **42**, 50, 702 (1964).
 - (17) H. C. Mishra and M. C. R. Symons, *J. Chem. Soc.*, 4490 (1963).
 - (18) N. Bailey and M. C. R. Symons, *J. Chem. Soc.*, 203 (1957).
 - (19) J. A. Connor and E. A. V. Ebsworth in "Advances in Inorganic Chemistry and Radiochemistry", Vol. 6, H. J. Emeléus and A. G. Sharpe, Ed., Academic Press, New York, N.Y., 1964.
 - (20) F. A. Cotton and G. Wilkinson, "Advanced Inorganic Chemistry", 3d ed, Interscience, New York, N.Y., 1972.
 - (21) B. L. Bartlett and D. Quane, *Inorg. Chem.*, **12**, 1925 (1973).
 - (22) K. K. Kelley and E. G. King, "Contributions to the Data on Theoretical Metallurgy. XIV. Entropies of the Elements and Inorganic Compounds", Bureau of Mines Bulletin 592, U.S. Government Printing Office, Washington, D.C., 1961.
 - (23) K. Clusius and P. Franzosini, *Z. Naturforsch. A*, **17**, 525 (1962).
 - (24) G. I. Kalishevich, P. V. Gel'd, and R. P. Krentsis, *Russ. J. Phys. Chem. (Engl. Transl.)*, **39**, 1602 (1965).
 - (25) A. B. Bonilla and C. W. Garland, *J. Phys. Chem. Solids*, **35**, 871 (1974).
 - (26) D. R. Stull and G. C. Sinke, "Thermodynamic Properties of the Elements", *Adv. Chem. Ser.*, **No. 18** (1956).
 - (27) E. K. Storms, "The Refractory Carbides", Academic Press, New York, N.Y., 1967.
 - (28) D. S. Dickson, J. R. Myers, and R. K. Saxer, *J. Phys. Chem.*, **69**, 4044 (1965).
 - (29) R. T. Grimley, R. P. Burns, and M. G. Inghram, *J. Chem. Phys.*, **34**, 664 (1961).
 - (30) D. M. Chizhikov, Y. V. Tsvetkov, E. K. Kazenas, and V. K. Tagirov, *Russ. J. Phys. Chem. (Engl. Transl.)*, **46**, 468 (1972).
 - (31) M. Farber and R. D. Srivastava, *Combust. Flame*, **20**, 43 (1973).
 - (32) G. Nagarajan, *Aust. J. Chem.*, **16**, 908 (1963).
 - (33) S. M. Ariya, S. A. Shchukarev, and V. B. Glushkova, *J. Gen. Chem. USSR (Engl. Transl.)*, **23**, 1307 (1953).
 - (34) A. D. Mah, *J. Am. Chem. Soc.*, **76**, 3363 (1954).
 - (35) A. D. Mah, Bureau of Mines Report of Investigations No. 7217, U.S. Government Printing Office, Washington, D.C., 1969.
 - (36) I. A. Novokhatskii and L. M. Lenev, *Russ. J. Inorg. Chem. (Engl. Transl.)*, **11**, 1078 (1966).
 - (37) K. K. Kelley, "Contributions to the Data on Theoretical Metallurgy. XIII. High-Temperature Heat-Content, Heat-Capacity, and Entropy Data for the Elements and Inorganic Compounds", Bureau of Mines Bulletin 584, U.S. Government Printing Office, Washington, D.C., 1960.
 - (38) J. N. Ramsey, D. Caplan, and A. A. Burr, *J. Electrochem. Soc.*, **103**, 135 (1956).
 - (39) J. D. Tretjakov and H. Schmalzried, *Ber. Bunsenges. Phys. Chem.*, **69**, 396 (1965).
 - (40) F. N. Mazandarany and R. D. Pehlke, *J. Electrochem. Soc.*, **121**, 711 (1974).
 - (41) Y. Jeannin, C. Mannerskantz, and R. D. Richardson, *Trans. Metall. Soc. AIME*, **227**, 300 (1963).
 - (42) A. D. Kulkarni and W. L. Worrell, *Metall. Trans.*, **3**, 2363 (1972).
 - (43) L. G. Hepler, *J. Am. Chem. Soc.*, **80**, 6181 (1958); C. N. Muldrow, Jr., and L. G. Hepler, *ibid.*, **79**, 4045 (1957). Comparison of quantities calculated in these two papers clearly illustrates the importance of basing thermodynamic calculations for aqueous solutions on reaction equations that correctly indicate the important species in solution. The calculations in the 1958 paper are as reliable as the available data permit, except for a change in ΔH_f° of $\text{CrO}_3(\text{c})$ discussed elsewhere in this review. But some of the calculations in the 1957 paper yielded incorrect results because of neglect of one important aqueous species.
 - (44) C. A. Neugebauer and J. L. Margrave, *J. Phys. Chem.*, **61**, 1429 (1957).
 - (45) F. R. Bichowsky and F. D. Rossini, "The Thermochemistry of the Chemical Substances", Reinhold, New York, N.Y., 1936.
 - (46) R. Richelmi and M. Laffitte, *C. R. Acad. Sci., Ser. C*, **265**, 541 (1967).
 - (47) J. D. McDonald and J. L. Margrave, *J. Inorg. Nucl. Chem.*, **30**, 665 (1968).
 - (48) W. V. Smith, K. S. Pitzer, and W. M. Latimer, *J. Am. Chem. Soc.*, **59**, 2642 (1937).
 - (49) K. Pan, *J. Chinese Chem. Soc. (Taipei)*, **1**, 1 (1954).
 - (50) J. Y. Cann and G. B. Mueller, *J. Am. Chem. Soc.*, **57**, 2525 (1935).
 - (51) J. R. Howard and G. H. Nancollas, *Trans. Faraday Soc.*, **53**, 1449 (1957).
 - (52) H. G. Linge and A. L. Jones, *Aust. J. Chem.*, **21**, 2189 (1968).
 - (53) J. D. Neuss and W. Rieman, *J. Am. Chem. Soc.*, **56**, 2238 (1934).
 - (54) J. R. Howard, V. S. K. Nair, and G. H. Nancollas, *Trans. Faraday Soc.*, **54**, 1034 (1958).
 - (55) N. Bailey, A. Carrington, K. A. K. Lott, and M. C. R. Symons, *J. Chem. Soc.*, 290 (1960).
 - (56) Y. Sasaki, *Acta Chem. Scand.*, **16**, 719 (1962).
 - (57) J. Galea and J. Haladjian, *Rev. Chim. Miner.*, **7**, 623 (1970).
 - (58) R. Arnek and S. R. Johansson, *Acta Chem. Scand.*, **26**, 2903 (1972).
 - (59) H. G. Linge and A. L. Jones, *Aust. J. Chem.*, **21**, 1445 (1968).
 - (60) W. G. Davies and J. E. Prue, *Trans. Faraday Soc.*, **51**, 1045 (1955).
 - (61) J. Y.-P. Tong and E. L. King, *J. Am. Chem. Soc.*, **75**, 6180 (1953).
 - (62) A. Lifshitz and B. Perlmuter-Hayman, *J. Phys. Chem.*, **65**, 2098 (1961); B. Perlmuter-Hayman and Y. Weissmann, *Isr. J. Chem.*, **6**, 17 (1968).
 - (63) J. G. Mason and A. D. Kowalak, *Inorg. Chem.*, **3**, 1248 (1964).
 - (64) J. Lagrange and J. Bye, *Bull. Soc. Chim. Fr.*, 1490 (1967).
 - (65) J. Y. Tong and R. L. Johnson, *Inorg. Chem.*, **5**, 1902 (1966).
 - (66) G. P. Haight, Jr., D. C. Richardson, and N. H. Coburn, *Inorg. Chem.*, **3**, 1777 (1964).
 - (67) F. Holloway, *J. Am. Chem. Soc.*, **74**, 224 (1952).
 - (68) J. H. Espenson and S. R. Helzer, *Inorg. Chem.*, **8**, 1051 (1969).
 - (69) O. Glemser and A. Mueller, *Z. Anorg. Chem.*, **334**, 150 (1964).
 - (70) M. J. Ferrante, J. M. Stuve, and M. P. Krug, Bureau of Mines Report of Investigations No. 7691, U.S. Government Printing Office, Washington, D.C., 1972.
 - (71) T. Nelson, C. Moss, and L. G. Hepler, *J. Phys. Chem.*, **64**, 376 (1960).
 - (72) L. F. Shmagin and A. A. Shidlovskii, *Russ. J. Phys. Chem.*, **45**, 310 (1971).
 - (73) F. D. Rossini, D. D. Wagman, W. H. Evans, S. Levine, and I. Jaffe, "Selected Values of Chemical Thermodynamic Properties", National Bureau of Standards Circular 500, U.S. Government Printing Office, Washington, D.C., 1952.
 - (74) A. A. Shidlovskii, A. A. Voskresenskii, and T. N. Balakireva, *Russ. J. Phys. Chem.*, **45**, 1067 (1971).
 - (75) L. G. Silén and A. E. Martell, "Stability Constants of Metal-Ion Complexes", *Chem. Soc., Spec. Publ.*, **No. 17**, (1964).
 - (76) G. L. Beyer and W. Rieman, *J. Am. Chem. Soc.*, **65**, 971 (1943).
 - (77) I. M. Kolthoff, R. W. Perlich, and W. Weiblen, *J. Phys. Chem.*, **46**, 561 (1942).
 - (78) S. Suzuki, *J. Chem. Soc. Jpn., Pure Chem. Sect.*, **74**, 219 (1952); *Chem. Abstr.*, **47**, 10970b (1953).
 - (79) M. Gaultier and G. Pannetier, *Rev. Chim. Miner.*, **9**, 271 (1972); M. Natarajan and E. E. Secco, *Can. J. Chem.*, **52**, 712 (1974).
 - (80) A. Engelbrecht and A. V. Grosse, *J. Am. Chem. Soc.*, **74**, 5262 (1952).
 - (81) A. J. Edwards, W. E. Falconer, and W. A. Sunder, *J. Chem. Soc., Dalton Trans.*, 541 (1974).
 - (82) W. H. Hartford and M. Darrin, *Chem. Rev.*, **58**, 1 (1958).
 - (83) J. G. Spencer and L. G. Hepler, unpublished enthalpy of reaction of $\text{CrO}_2\text{Cl}_2(\text{liq})$ with excess $\text{OH}^-(\text{aq})$, 1959.
 - (84) W. N. Hansen and M. Griffel, *J. Chem. Phys.*, **28**, 902 (1958).
 - (85) L. Brewer, L. A. Bromley, P. W. Gilles, and N. L. Lofgren in "The Chemistry and Metallurgy of Miscellaneous Materials: Thermodynamics", L. L. Quill, Ed., McGraw-Hill, New York, N.Y., 1950.
 - (86) H. Tanaka, A. Yamaguchi, and J. Moriyama, *J. Jpn. Metall. Soc.*, **35**, 1161 (1971). We are grateful to Dr. K. Nakamura for his translation.
 - (87) R. A. Kent and J. L. Margrave, *J. Am. Chem. Soc.*, **87**, 3582 (1965).
 - (88) K. F. Zmbov and J. L. Margrave, *J. Inorg. Nucl. Chem.*, **29**, 673 (1967).
 - (89) J. W. Stout and R. C. Chisholm, *J. Chem. Phys.*, **36**, 979 (1962).
 - (90) M. Papapietro and G. Monnier, *C. R. Acad. Sci., Ser. C*, **273**, 703 (1971).
 - (91) C. G. Maier, Bureau of Mines Bulletin 436, U.S. Government Printing Office, Washington, D.C., 1942.
 - (92) R. Shaw and R. F. Porter, personal communication to L. G. H., 1973.
 - (93) R. C. Schoonmaker, A. H. Friedman, and R. F. Porter, *J. Chem. Phys.*, **31**, 1586 (1959).
 - (94) K. Sano, *J. Chem. Soc. Jpn.*, **59**, 17 (1938); *Chem. Abstr.*, **32**, 2813 (1938).
 - (95) I. V. Vasil'kova, A. I. Efimov, and B. Z. Pitirimov, *Russ. J. Inorg. Chem.*, **9**, 417 (1964); S. A. Shchukarev, I. V. Vasil'kova, A. I. Efimov, and B. Z. Pitirimov, *ibid.*, **11**, 268 (1966).
 - (96) N. W. Gregory and T. R. Burton, *J. Am. Chem. Soc.*, **75**, 6054 (1953).
 - (97) S. A. Shchukarev, T. A. Tolmacheva, M. A. Oranskaya, and E. A. Ukrainitseva, *Zh. Neorg. Khim.*, **1**, 1697 (1956).
 - (98) R. J. Sime and N. W. Gregory, *J. Am. Chem. Soc.*, **82**, 800 (1960).
 - (99) V. A. Rupcheva, T. V. Romanova, and S. A. Amirova, *Russ. J. Inorg. Chem.*, **15**, 170 (1970).
 - (100) T. L. Allen, *J. Am. Chem. Soc.*, **78**, 5476 (1956).
 - (101) L. L. Handy and N. W. Gregory, *J. Am. Chem. Soc.*, **72**, 5049 (1950).
 - (102) C.-F. Shieh and N. W. Gregory, *J. Phys. Chem.*, **77**, 2346 (1973).
 - (103) L. L. Handy and N. W. Gregory, *J. Am. Chem. Soc.*, **74**, 2050 (1952).
 - (104) K. Lascelles, R. A. J. Shelton, and H. Schafer, *J. Less-Common Met.*, **29**, 109 (1972).
 - (105) I. V. Vasil'kova, A. I. Efimov, P. S. Shapkin, and A. A. Vishnyakov, *Russ. J. Inorg. Chem.*, **14**, 219 (1969).
 - (106) N. V. Galitskii, K. P. Minina, and G. M. Borisovskaya, *Russ. J. Inorg. Chem.*, **12**, 1386 (1967).
 - (107) C. M. Cook, Jr., *J. Inorg. Nucl. Chem.*, **25**, 123 (1963).
 - (108) H. W. Jenkins, G. Mamantov, and D. L. Manning, *J. Electrochem. Soc.*, **117**, 183 (1970).
 - (109) M. W. Evans in "The Transuranium Elements", G. T. Seaborg, J. J. Katz, and W. M. Manning, Ed., McGraw-Hill, New York, N.Y., 1949.
 - (110) P. Gross, G. L. Wilson, and W. A. Gutteridge, *J. Chem. Soc. A*, 1908 (1970).

- (111) G. Grube and L. Schlecht, *Z. Electrochem.*, **32**, 178 (1926).
- (112) L. G. Sillén and A. E. Martell, "Stability Constants of Metal-Ion Complexes", Supplement No. 1, *Chem. Soc. Spec. Publ.*, **No. 25** (1971).
- (113) K. M. Sancier and J. S. Mills, *J. Phys. Chem.*, **67**, 1438 (1963); B. A. Zabin and H. Taube, *Inorg. Chem.*, **3**, 963 (1964); J. Levitan, *J. Electrochem. Soc.*, **111**, 286 (1964); G. Thompson, *Univ. Calif. Radiat. Lab. Rep.*, 11410 (1964); J. E. Finholt, *ibid.*, 8879 (1960).
- (114) U. von Meyenburg, O. Siroky, and G. Schwarzenbach, *Helv. Chim. Acta*, **56**, 1099 (1973).
- (115) T. W. Swaddle and P. C. Kong, *Can. J. Chem.*, **48**, 3223 (1970).
- (116) C. Postmus and E. L. King, *J. Phys. Chem.*, **59**, 1208 (1955).
- (117) K. Schug and E. L. King, *J. Am. Chem. Soc.*, **80**, 1089 (1958).
- (118) T. W. Swaddle and E. L. King, *Inorg. Chem.*, **4**, 532 (1965).
- (119) J. H. Espenson and O. J. Parker, *J. Am. Chem. Soc.*, **90**, 3689 (1968).
- (120) C. F. Hale and E. L. King, *J. Phys. Chem.*, **71**, 1779 (1967).
- (121) R. J. Baltisberger and E. L. King, *J. Am. Chem. Soc.*, **86**, 795 (1964).
- (122) H. S. Gates and E. L. King, *J. Am. Chem. Soc.*, **80**, 5011 (1958).
- (123) D. F. C. Morris and S. D. Hammond, *Electrochim. Acta*, **13**, 545 (1968).
- (124) A. B. Lamb and G. R. Fonda, *J. Am. Chem. Soc.*, **43**, 1155 (1921); R. E. Hamm and C. M. Shull, Jr., *ibid.*, **73**, 1240 (1951); G. Schwarzenbach, *Pure Appl. Chem.*, **5**, 377 (1962).
- (125) H. B. Johnson and W. L. Reynolds, *Inorg. Chem.*, **2**, 468 (1963); E. L. King, M. J. M. Woods, and H. S. Gates, *J. Am. Chem. Soc.*, **80**, 5015 (1958).
- (126) J. H. Espenson and E. L. King, *J. Phys. Chem.*, **64**, 380 (1960).
- (127) L. O. Spreer and E. L. King, *Inorg. Chem.*, **10**, 916 (1971).
- (128) T. W. Swaddle and G. Guastalla, *Inorg. Chem.*, **7**, 1915 (1968).
- (129) J. H. Espenson and D. E. Binau, *Inorg. Chem.*, **5**, 1365 (1966).
- (130) K. Nagase and N. Tanaka, *Bull. Chem. Soc. Jpn.*, **45**, 1097 (1972); Y. Kaji, A. Uehara, E. Kyuno, and R. Tsuchiya, *ibid.*, **43**, 1906 (1970).
- (131) K. M. Jones and J. Bjerrum, *Acta Chem. Scand.*, **19**, 974 (1965).
- (132) E. E. Mercer and J. A. Hormuth, *J. Inorg. Nucl. Chem.*, **31**, 2145 (1969).
- (133) M. Ardon and N. Sutin, *Inorg. Chem.*, **6**, 2268 (1967).
- (134) N. Fogel, J. M. J. Tai, and J. Yarborough, *J. Am. Chem. Soc.*, **84**, 1145 (1962); R. Tsuchiya and A. Umayahara, *Bull. Chem. Soc. Jpn.*, **36**, 554 (1963); N. Tanaka, K. Ogino, and G. Sato, *ibid.*, **39**, 366 (1966).
- (135) D. N. Hume and I. M. Kolthoff, *J. Am. Chem. Soc.*, **65**, 1897 (1943).
- (136) F. H. Guzzetta and W. B. Hadley, *Inorg. Chem.*, **3**, 259 (1964).
- (137) K. G. Poulsen, J. Bjerrum, and I. Poulsen, *Acta Chem. Scand.*, **8**, 921 (1954).
- (138) J. C. Templeton and E. L. King, *J. Am. Chem. Soc.*, **93**, 7160 (1971).
- (139) S. C. Chan and K. Y. Hui, *Aust. J. Chem.*, **21**, 3061 (1968).
- (140) A. J. Cunningham, D. A. House, and H. K. J. Powell, *Aust. J. Chem.*, **23**, 2375 (1970).
- (141) N. V. Duffy and J. E. Earley, *J. Am. Chem. Soc.*, **89**, 272 (1967).
- (142) G. Guastalla and T. W. Swaddle, *Can. J. Chem.*, **51**, 821 (1973).
- (143) J. P. Birk and J. H. Espenson, *Inorg. Chem.*, **7**, 991 (1968); D. N. Purohit and J. Bjerrum, *J. Inorg. Nucl. Chem.*, **33**, 2067 (1971); J. N. Armor and A. Haim, *J. Am. Chem. Soc.*, **93**, 867 (1971); J. H. Espenson and W. R. Bushey, *Inorg. Chem.*, **10**, 2457 (1971).
- (144) W. W. Wendlandt and J. L. Bear, *J. Inorg. Nucl. Chem.*, **22**, 77 (1961); R. Tsuchiya, Y. Kaji, A. Uehara, and E. Kyuno, *Bull. Chem. Soc. Jpn.*, **42**, 1881 (1969); W. W. Wendlandt, G. D'Ascenzo, and R. H. Gore, *Thermochim. Acta*, **1**, 488 (1970); E. L. Simmons and W. W. Wendlandt, *ibid.*, **2**, 217 (1971).
- (145) F. S. Boericke and W. M. Bangert, Bureau of Mines Report of Investigations No. 3813, U.S. Government Printing Office, Washington, D.C., 1945.
- (146) H. M. Chen and J. Chipman, *Trans. Am. Soc. Met.*, **38**, 70 (1947).
- (147) T. Katsura and A. Muan, *Trans. Metall. Soc. AIME*, **230**, 77 (1964).
- (148) F. Müller and O. J. Kleppa, *J. Inorg. Nucl. Chem.*, **35**, 2673 (1973).
- (149) N. G. Schmahl and E. Minzi, *Z. Phys. Chem.*, **47**, 358 (1965).
- (150) E. Aukrust and A. Muan, *J. Am. Ceram. Soc.*, **46**, 358 (1963).
- (151) "JANAF Thermochemical Tables", 2d ed, D. R. Stull, H. Prophet, et al., NSRDS-NBS 37, U.S. Government Printing Office, Washington, D.C., 1971.
- (152) A. G. Zalazinskii, V. F. Balakirev, N. M. Chebotaev, and G. I. Chufarov, *Russ. J. Inorg. Chem.*, **14**, 326 (1969).
- (153) H. v. Wartenberg, *Z. Anorg. Chem.*, **249**, 100 (1942).
- (154) H. A. Doerner, Bureau of Mines Technical Paper No. 577, U.S. Government Printing Office, Washington, D.C., 1937.
- (155) H. Oppermann, *Z. Anorg. Chem.*, **359**, 51 (1968).
- (156) R. J. Sime and N. W. Gregory, *J. Am. Chem. Soc.*, **82**, 93 (1960).
- (157) G. P. Haight, Jr., T. J. Huang, and H. Platt, *J. Am. Chem. Soc.*, **96**, 3137 (1974); J. Y.-P. Tong and E. L. King, *ibid.*, **82**, 3805 (1960); M. Ardon and R. A. Plane, *ibid.*, **81**, 3197 (1959); F. H. Westheimer, *Chem. Rev.*, **45**, 419 (1949); K. Pan and S.-H. Lin, *J. Chin. Chem. Soc. (Taipei)*, **7**, 75 (1960); H.-L. Krauss and G. Gnatz, *Chem. Ber.*, **92**, 2110 (1959); J. H. Espenson and R. J. Kinney, *Inorg. Chem.*, **10**, 376 (1971).
- (158) G. Wolf, *Phys. Status Solidi A*, **5**, 627 (1971).
- (159) A. Sieverts and A. Gotta, *Z. Anorg. Chem.*, **172**, 1 (1928).
- (160) A. D. Stock and K. I. Hardcastle, *J. Inorg. Nucl. Chem.*, **32**, 1183 (1970).
- (161) H. Mabuchi, N. Sano, and Y. Matsushita, *Metall. Trans.*, **2**, 1503 (1971).
- (162) H. Kleykamp, *Ber. Bunsenges. Phys. Chem.*, **73**, 354 (1969).
- (163) F. J. Kohl and C. A. Stearns, *J. Phys. Chem.*, **74**, 2714 (1970).
- (164) A. D. Mah, Bureau of Mines Report of Investigations No. 5529, U.S. Government Printing Office, Washington, D.C., 1960.
- (165) T. Mills, *J. Less-Common Met.*, **22**, 373 (1970); **23**, 317 (1971); A. U. Seybolt and R. A. Oriani, *J. Met.*, **8**, AIME Trans., **206**, 556 (1956); J. Pearson and U. J. C. Ende, *J. Iron Steel Inst., London*, **175**, 52 (1953).
- (166) T. G. Chart, National Physical Laboratory (England) Report Chem., Aug. 18, 1972.
- (167) V. N. Eremenko, G. M. Lukashenko, and V. R. Sidorko, *Russ. J. Phys. Chem.*, **45**, 1133 (1971); A. S. Bolgar, S. P. Gordienko, A. A. Lysenko, and V. V. Fesenko, *ibid.*, **45**, 1154 (1971); Y. M. Golutvin and L. Chin-K'uei, *ibid.*, **35**, 62 (1961).
- (168) G. I. Kalishevich, V. L. Zagryazhskii, L. V. Shchipanova, N. N. Serebrennikov, and P. V. Gel'd, *Russ. J. Phys. Chem.*, **43**, 132 (1969).
- (169) J. P. Hager and J. F. Elliott, *Trans. Metall. Soc. AIME*, **239**, 513 (1967).
- (170) J. Drowart, A. Pattoret, and S. Smoes, *Proc. Br. Ceram. Soc.*, **8**, 67 (1967).
- (171) F. Grönwold and E. F. Westrum, Jr., *Z. Anorg. Chem.*, **328**, 272 (1964).
- (172) F. Grönwold, *J. Chem. Thermodyn.*, **5**, 545 (1973).
- (173) F. A. Cotton, A. K. Fischer, and G. Wilkinson, *J. Am. Chem. Soc.*, **78**, 5168 (1956).
- (174) H. A. Skinner and Y. Virmani, *Rev. Roum. Chim.*, **17**, 467 (1972).
- (175) J. A. Connor, H. A. Skinner, and Y. Virmani, *J. Chem. Soc., Faraday Trans. 1*, **68**, 1754 (1972).
- (176) T. N. Rezukhina and V. V. Shvyrev, *Vestn. Mosk. Univ.*, **7**, 41 (1952); *Chem. Abstr.*, **47**, 4676h (1953); A. A. Boni, *J. Electrochem. Soc.*, **113**, 1089 (1966).
- (177) N. S. Hawkins, H. C. Matraw, W. W. Sabol, and D. R. Carpenter, *J. Chem. Phys.*, **23**, 2422 (1955).
- (178) K. Kawai and H. Murata, *Bull. Chem. Soc. Jpn.*, **33**, 1008 (1960).
- (179) D. L. S. Brown, J. A. Connor, and H. A. Skinner, *J. Chem. Soc., Faraday Trans. 1*, **70**, 1649 (1974).
- (180) J. D. Cox and G. Pilcher, "Thermochemistry of Organic and Organometallic Compounds", Academic Press, New York, N.Y., 1970.
- (181) S. J. Ashcroft and C. T. Mortimer, "Thermochemistry of Transition Metal Complexes", Academic Press, New York, N.Y., 1970.
- (182) J. A. Connor, H. A. Skinner, and Y. Virmani, *J. Chem. Soc., Faraday Trans. 1*, **69**, 1218 (1973).
- (183) D. C. Bradley and M. J. Hillyer, *Trans. Faraday Soc.*, **62**, 2382 (1966).
- (184) V. I. Tel'noi, I. B. Rabinovich, B. G. Gribov, A. S. Pashinkin, B. A. Salamatina, and V. I. Chernova, *Russ. J. Phys. Chem.*, **46**, 465 (1972).
- (185) M. Sorai, M. Tachiki, H. Suga, and S. Seki, *J. Phys. Soc. Jpn.*, **30**, 750 (1971).
- (186) M. Sorai and S. Seki, *J. Phys. Soc. Jpn.*, **32**, 382 (1972).
- (187) J. T. S. Andrews, E. F. Westrum, Jr., and N. Bjerrum, *J. Organomet. Chem.*, **17**, 293 (1969).
- (188) J. O. Hill and R. J. Irving, *J. Chem. Soc. A*, 1413 (1967).
- (189) T. P. Melia and R. Merrifield, *J. Chem. Soc. A*, 2819 (1968); *J. Inorg. Nucl. Chem.*, **32**, 1489 (1970).
- (190) R. Fontaine, C. Pommier, and G. Guiochon, *Bull. Soc. Chim. Fr.*, 3011 (1972).
- (191) L. A. Woodward and H. L. Roberts, *Trans. Faraday Soc.*, **52**, 615 (1956); R. H. Busey and O. L. Keller, *J. Chem. Phys.*, **41**, 215 (1964).
- (192) R. K. Koch and W. E. Anable, Bureau of Mines Report of Investigations No. 7063, U.S. Government Printing Office, Washington, D.C., 1968.
- (193) F. L. Oetting and J. D. Navratil, *J. Chem. Eng. Data*, **17**, 230 (1972); B. Y. Berezin, V. Y. Chekhovskoi, and A. E. Sheindlin, *High Temp.-High Pressures*, **3**, 287 (1971); J. A. Treverton and J. L. Margrave, *Proceedings of the ASTM Conference on Thermodynamic Properties*, Boston, Mass., Oct 1970, p 489; A. Cezaırlıyan, M. S. Morse, H. A. Berman, and C. W. Beckett, *J. Res. Natl. Bur. Stand., Sect. A*, **74**, 65 (1970).
- (194) G. DeMaria, R. P. Burns, J. Drowart, and M. G. Inghram, *J. Chem. Phys.*, **32**, 1373 (1960).
- (195) B. A. Staskiewicz, J. R. Tucker, and P. E. Snyder, *J. Am. Chem. Soc.*, **77**, 2987 (1955).
- (196) A. D. Mah, *J. Phys. Chem.*, **61**, 1572 (1957).
- (197) S. Berglund and P. Kierkegaard, *Acta Chem. Scand.*, **23**, 329 (1969).
- (198) R. P. Burns, G. DeMaria, J. Drowart, and R. T. Grimley, *J. Chem. Phys.*, **32**, 1363 (1960).
- (199) J. Bousquet and A. Guillon, *Bull. Soc. Chim. Fr.*, 2250 (1966); J. Bousquet, J.-C. David, M. Diot, and A. Guillon, *ibid.*, 2390 (1968); S. Zador and C. B. Alcock, *J. Chem. Thermodyn.*, **2**, 9 (1970).
- (200) E. G. King, W. W. Weller, and A. U. Christensen, Bureau of Mines Report of Investigations No. 5664, U.S. Government Printing Office, Washington, D.C., 1960; L. A. Cosgrove and P. E. Snyder, *J. Am. Chem. Soc.*, **75**, 1227 (1953).
- (201) A. D. Kirshenbaum and A. J. Beardell, *Thermochim. Acta*, **4**, 239 (1972).
- (202) E. R. Plante, Ph.D. Thesis, University of Kansas, 1960.
- (203) J. Berkowitz, M. G. Inghram, and W. A. Chupka, *J. Chem. Phys.*, **26**, 842 (1957).
- (204) E. K. Kazenas and Y. V. Tsvetkov, *Russ. J. Inorg. Chem.*, **14**, 5 (1969).
- (205) R. L. Graham and L. G. Hepler, *J. Am. Chem. Soc.*, **78**, 4846 (1956).
- (206) P. A. G. O'Hare and H. R. Hoekstra, *J. Chem. Thermodyn.*, **6**, 117 (1974).
- (207) P. A. G. O'Hare and H. R. Hoekstra, *J. Chem. Thermodyn.*, **5**, 851 (1973).
- (208) L. G. Hepler, *Can. J. Chem.*, **47**, 3469 (1969).
- (209) P. A. G. O'Hare, K. J. Jensen, and H. R. Hoekstra, *J. Chem. Thermodyn.*, **6**, 681 (1974).
- (210) W. W. Weller and E. G. King, Bureau of Mines Report of Investigations No. 6147, U.S. Government Printing Office, Washington, D.C., 1963.
- (211) R. Barany, Bureau of Mines Report of Investigations No. 6143, U.S. Government Printing Office, Washington, D.C., 1962.
- (212) C. N. Muldrow, Jr., and L. G. Hepler, *J. Phys. Chem.*, **62**, 982 (1958).
- (213) P. A. G. O'Hare, *J. Chem. Thermodyn.*, **6**, 425 (1974).
- (214) P. Post, *Bull. Soc. Chim. Fr.*, 2993 (1972).
- (215) M. F. Koehler, L. B. Pankratz, and R. Barany, Bureau of Mines Report of Investigations No. 5973, U.S. Government Printing Office, Washington, D.C., 1962.
- (216) A. P. Zhidikova, I. L. Khodakovskii, M. A. Urusova, and V. M. Valyashko, *Russ. J. Inorg. Chem.*, **18**, 612 (1973).

- (217) R. Yamdagni, C. Pupp, and R. F. Porter, *J. Inorg. Nucl. Chem.*, **32**, 3509 (1970).
- (218) A. Navrotsky and O. J. Kleppa, *Inorg. Chem.*, **6**, 2119 (1967); A. Brenner and W. H. Metzger, Jr., *J. Electrochem. Soc.*, **115**, 258 (1968).
- (218a) W. W. Weller and K. K. Kelley, Bureau of Mines Report of Investigations No. 6191, U.S. Government Printing Office, Washington, D.C., 1963.
- (219) T. Nelson, C. Moss, and L. G. Hepler, *J. Phys. Chem.*, **64**, 376 (1960).
- (220) M. Farber and R. D. Srivastava, *Combust. Flame*, **20**, 33 (1973).
- (221) D. W. Osborne, H. E. Flotow, and H. R. Hoekstra, *J. Chem. Thermodyn.*, **6**, 179 (1974).
- (222) G. Verhaegen, R. Colin, G. Exsteen, and J. Drowart, *Trans. Faraday Soc.*, **61**, 1372 (1965).
- (223) C. Pupp, R. Yamdagni, and R. F. Porter, *J. Inorg. Nucl. Chem.*, **31**, 2021 (1969).
- (224) C. N. Muldrow, Jr., and L. G. Hepler, *J. Am. Chem. Soc.*, **78**, 5989 (1956); K. Pan, *J. Chin. Chem. Soc. (Taipei)*, **1**, 16 (1954); J. E. Ricci and W. F. Linke, *J. Am. Chem. Soc.*, **73**, 3601 (1951).
- (225) R. Barany, Bureau of Mines Report of Investigations No. 6618, U.S. Government Printing Office, Washington, D.C., 1965.
- (226) J. Ziolkowski and P. Courtine, *Ann. Chim. (Paris)*, **8**, 303 (1973).
- (227) W. W. Weller, Bureau of Mines Report of Investigations No. 6782, U.S. Government Printing Office, Washington, D.C., 1966.
- (228) W. W. Weller and K. K. Kelley, Bureau of Mines Report of Investigations No. 6357, U.S. Government Printing Office, Washington, D.C., 1964.
- (229) R. A. Fisher, E. W. Hornung, G. E. Brodale, and W. F. Giauque, *J. Chem. Phys.*, **56**, 193 (1972); E. W. Hornung, G. E. Brodale, R. A. Fisher, and W. F. Giauque, *ibid.*, **56**, 5007 (1972); G. E. Brodale, R. A. Fisher, E. W. Hornung, and W. F. Giauque, *ibid.*, **56**, 6118 (1972).
- (230) P. Souchay, *Pure Appl. Chem.*, **6**, 61 (1963).
- (231) W. P. Griffith and P. J. B. Lesniak, *J. Chem. Soc. A*, 1066 (1969); E. Pungor and A. Halasz, *J. Inorg. Nucl. Chem.*, **32**, 1187 (1970).
- (232) D. S. Honig and K. Kustin, *Inorg. Chem.*, **11**, 65 (1972).
- (233) R. Arnek and I. Szilard, *Acta Chem. Scand.*, **22**, 1334 (1968); N. Kiba and T. Takeuchi, *Bull. Chem. Soc. Jpn.*, **46**, 3086 (1973); N. Kiba and T. Takeuchi, *J. Inorg. Nucl. Chem.*, **36**, 847 (1974); N. D. Jespersen, *ibid.*, **35**, 3873 (1973).
- (234) L. Krumenacker, *Ann. Chim. (Paris)*, 425 (1972).
- (235) L. McKean and M. T. Pope, *Inorg. Chem.*, **13**, 747 (1974); G. A. Tsigdinos and C. J. Hallada, *J. Less-Common Met.*, **36**, 79 (1974).
- (236) O. W. Rollins, *J. Inorg. Nucl. Chem.*, **33**, 75 (1971).
- (237) D. W. Osborne, F. Schreiner, J. G. Malm, H. Selig, and L. Rochester, *J. Chem. Phys.*, **44**, 2802 (1966).
- (238) B. Weinstock and G. L. Goodman, *Adv. Chem. Phys.*, **9**, 169 (1965).
- (239) A. P. Brady, O. E. Myers, and J. D. Clauss, *J. Phys. Chem.*, **64**, 588 (1960).
- (240) G. H. Cady and G. B. Hargreaves, *J. Chem. Soc.*, 1563 (1961).
- (241) J. L. Settle, H. M. Feder, and W. N. Hubbard, *J. Phys. Chem.*, **65**, 1337 (1961).
- (242) O. E. Myers and A. P. Brady, *J. Phys. Chem.*, **64**, 591 (1960).
- (243) R. L. Nuttall, K. L. Churney, and M. V. Kilday, National Bureau of Standards Report 10904, U.S. Department of Commerce, Washington, D.C., 1972.
- (244) J. Burgess, I. Haigh, and R. D. Peacock, *J. Chem. Soc., Dalton Trans.*, 1062 (1974).
- (245) V. T. Orekov and A. G. Rybakov, *Zh. Fiz. Khim.*, **47**, 1612 (1973).
- (246) R. T. Paine and R. S. McDowell, *Inorg. Chem.*, **13**, 2366 (1974).
- (247) G. H. Cady and G. B. Hargreaves, *J. Chem. Soc.*, 1568 (1961).
- (248) K. F. Zmbov, O. M. Uy, and J. L. Margrave, *J. Phys. Chem.*, **73**, 3008 (1969).
- (249) S. A. Shchukarev, G. I. Novikov, I. V. Vasil'kova, A. V. Suvorov, N. V. Andreeva, B. N. Sharupin, and A. K. Baev, *Russ. J. Inorg. Chem.*, **5**, 802 (1960).
- (250) H. Oppermann and G. Stöver, *Z. Anorg. Chem.*, **387**, 218 (1972).
- (251) H. Oppermann, G. Stöver, and G. Kunze, *Z. Anorg. Chem.*, **387**, 201 (1972).
- (252) R. L. Graham and L. G. Hepler, *J. Phys. Chem.*, **63**, 723 (1959).
- (253) S. A. Shchukarev, I. V. Vasil'kova, and B. N. Sharupin, *Vestn. Leningr. Univ.*, (14, No. 4), *Fiz., Khim.*, No. 1, 73 (1959); *Chem. Abstr.*, **53**, 13856d (1959).
- (254) H. Oppermann, *Z. Anorg. Chem.*, **379**, 262 (1970).
- (255) S. A. Shchukarev, I. V. Vasil'kova, and B. N. Sharupin, *Vestn. Leningr. Univ.*, (16, No. 22), *Fiz., Khim.*, No. 4, 130 (1961).
- (256) N. Hultgren and L. Brewer, *J. Phys. Chem.*, **60**, 947 (1956).
- (257) A. Shukurov, T. Nikol'skaya, I. V. Vasil'kova, and S. A. Shchukarev, *Izv. Akad. Nauk Tadzh. SSR, Otd. Geol.-Khim. Tekh. Nauk* No. 4, 23 (1961); *Chem. Abstr.*, **57**, 14498g (1962).
- (258) H. Oppermann, *Z. Anorg. Chem.*, **395**, 249 (1973).
- (259) S. A. Shchukarev, I. V. Vasil'kova, and N. D. Zaitseva, *Vestn. Leningr. Univ.*, (16, No. 22), *Fiz., Khim.*, No. 4, 127 (1961); *Chem. Abstr.*, **56**, 9510g (1962).
- (260) I. V. Vasil'kova, N. D. Zaitseva, and Y. S. Svalov, *Vestn. Leningr. Univ.*, (16, No. 16), *Fiz., Khim.*, No. 3, 140 (1961); *Chem. Abstr.*, **56**, 4168e (1962).
- (261) H. Oppermann, G. Kunze, and G. Stöver, *Z. Anorg. Chem.*, **387**, 339 (1972).
- (262) D. S. Barnes, G. Pilcher, D. A. Pittam, H. A. Skinner, D. Todd, and Y. Virmani, *J. Less-Common Met.*, **36**, 177 (1974).
- (263) Y. Virmani, D. S. Barnes, and H. A. Skinner, *J. Chem. Soc., Dalton Trans.*, 399 (1974).
- (264) T. L. Allen and D. M. Yost, *J. Chem. Phys.*, **22**, 855 (1954).
- (265) H. Oppermann, *Z. Anorg. Chem.*, 383, 285 (1971).
- (266) N. G. Barancheeva and L. A. Zharkova, *Russ. J. Inorg. Chem.*, **40**, 736 (1966).
- (266a) Y. Gerasimov, *Dokl. Mezhdunar. Kongr. Teor. Prikl. Khim.*, **13**, 77 (1953); *Chem. Abstr.*, **48**, 4941b (1954).
- (266b) V. A. Levitskii, M. Ya. Frenkel, and T. N. Rezukhina, *Elektrokimiya*, **1**, 1371 (1965); *Chem. Abstr.*, **64**, 16722g (1966).
- (267) L. A. Zharkova and N. G. Barancheeva, *Russ. J. Phys. Chem.*, **38**, 401 (1964).
- (268) E. F. Westrum, Jr., and J. J. McBride, *Phys. Rev.*, **98**, 270 (1955).
- (269) N. Parravano and G. Malquori, *Atti Accad. Naz. Lincei, Cl. Sci. Fis., Mat., Nat., Rend.*, **7**, 19, 109 (1928); *Chem. Abstr.*, **22**, 2091 (1928); I. A. Makolkin, *Russ. J. Phys. Chem.*, **14**, 110 (1940); *Chem. Abstr.*, **35**, 959^g (1941).
- (270) P. A. G. O'Hare, E. Benn, F. Y. Cheng, and G. Kuzmycz, *J. Chem. Thermodyn.*, **2**, 797 (1970).
- (271) W. Biltz and A. Kocher, *Z. Anorg. Chem.*, **248**, 172 (1941).
- (272) A. Muller, T. P. Prasad, and R. Menge, *Z. Anorg. Chem.*, **391**, 107 (1972); T. P. Prasad, E. Diemann, and A. Muller, *J. Inorg. Nucl. Chem.*, **35**, 1895 (1973).
- (273) A. D. Mah, Bureau of Mines Report of Investigations No. 6337, U.S. Government Printing Office, Washington, D.C., 1963.
- (274) L. B. Pankratz, W. W. Weller, and E. G. King, Bureau of Mines Report of Investigations No. 6861, U.S. Government Printing Office, Washington, D.C., 1966; I. E. Paukov, P. G. Streikov, and V. S. Flatkina, *Russ. J. Phys. Chem.*, **42**, 1576 (1968).
- (275) A. Solbakken and P. H. Emmett, *J. Am. Chem. Soc.*, **91**, 31 (1969).
- (276) R. J. Peavler and A. W. Searcy, *J. Am. Chem. Soc.*, **78**, 2076 (1956).
- (277) H. K. El-Shamy and A. M. El-Aggan, *J. Am. Chem. Soc.*, **75**, 1187 (1953).
- (278) G. P. Haight, Jr., *J. Inorg. Nucl. Chem.*, **24**, 663 (1962).
- (279) M. Ardon and A. Pernick, *Inorg. Chem.*, **12**, 2484 (1973).
- (280) H. K. Saha and M. C. Halder, *J. Inorg. Nucl. Chem.*, **33**, 705 (1971); H. K. Saha and A. K. Banerjee, *ibid.*, **33**, 2989 (1971).
- (281) I. N. Marov, Y. N. Dubrov, A. N. Ermakov, and G. N. Martynova, *Russ. J. Inorg. Chem.*, **14**, 224 (1969); I. N. Marov, Y. N. Dubrov, V. K. Belayeva, and A. N. Ermakov, *ibid.*, **13**, 1107 (1968).
- (282) J. Burgess, I. Haigh, R. D. Peacock, and P. Taylor, *J. Chem. Soc., Dalton Trans.*, 1064 (1974).
- (283) R. Gut, E. Schmid, and J. Serrallach, *Helv. Chim. Acta*, **54**, 593 (1971).
- (284) A. I. Efimov and L. P. Belorukova, *Russ. J. Inorg. Chem.*, **12**, 792 (1967).
- (285) W. U. Malik and S. I. Ali, *Talanta*, **8**, 737 (1961).
- (286) I. M. Kolthoff and W. J. Tomsicek, *J. Phys. Chem.*, **40**, 247 (1936); W. U. Malik and S. I. Ali, *Indian J. Chem.*, **1**, 374 (1963).
- (287) M. Ardon and A. Pernick, *Inorg. Chem.*, **13**, 2275 (1974).
- (288) Y. Sasaki and A. G. Sykes, *J. Chem. Soc., Chem. Commun.*, 767 (1973).
- (289) A. R. Bowen and H. Taube, *Inorg. Chem.*, **13**, 2245 (1974); J. A. Filipponi, Jr., H. J. Sniadoch, and R. L. Grayson, *ibid.*, **13**, 2121 (1974).
- (290) R. R. Monchamp and F. A. Cotton, *J. Chem. Soc.*, 1438 (1960).
- (291) D. N. Astrov, E. S. Itsekevich, and K. A. Sharifov, *Zh. Fiz. Khim.*, **29**, 424 (1955); *Chem. Abstr.*, **50**, 13588f (1956).
- (292) A. D. Westland and V. Uzelac, *Can. J. Chem.*, **48**, 2871 (1970).
- (293) E. R. Plante and A. B. Sessoms, *J. Res. Natl. Bur. Stand., Sect. A*, **77**, 237 (1973).
- (294) L. Leibowitz, M. G. Chasanov, and L. W. Mishler, *Trans. Metall. Soc. AIME*, **245**, 981 (1969).
- (295) A. D. Mah, *J. Am. Chem. Soc.*, **81**, 1582 (1959).
- (296) T. V. Charlou and O. J. Kleppa, *J. Chem. Thermodyn.*, **5**, 325 (1973).
- (297) F. E. Rizzo, L. R. Bidwell, and D. F. Frank, *Trans. Metall. Soc. AIME*, **239**, 1901 (1967).
- (298) W. A. Chupka, J. Berkowitz, and C. F. Giese, *J. Chem. Phys.*, **30**, 827 (1959).
- (299) G. Huff, E. Squitieri, and P. E. Snyder, *J. Am. Chem. Soc.*, **70**, 3380 (1948).
- (300) R. J. Ackermann, R. J. Thorn, C. Alexander, and M. Tetenbaum, *J. Phys. Chem.*, **64**, 350 (1960).
- (301) R. J. Ackermann and E. G. Rauh, *J. Phys. Chem.*, **67**, 2596 (1963).
- (302) J. H. Norman and H. G. Staley, *J. Chem. Phys.*, **43**, 3804 (1965).
- (303) J. Bousquet and G. Perachon, *C. R. Acad. Sci.*, **258**, 934 (1964).
- (304) H. C. Ko, T. Matsui, and L. G. Hepler, *Thermochim. Acta*, **10**, 211 (1974); see also ref 322.
- (305) V. I. Spitsyn and N. N. Patsukova, *Russ. J. Inorg. Chem.*, **10**, 1304 (1965).
- (306) G. R. Belton and R. L. McCarron, *J. Phys. Chem.*, **68**, 1852 (1964).
- (307) O. Glemser and H. Ackermann, *Z. Anorg. Chem.*, **325**, 281 (1963); O. Glemser and R. J. Haeseler, *ibid.*, **316**, 168 (1962).
- (308) G. Meyer, J. F. Oosterom, and W. J. W. Oeveren, *Recl. Trav. Chim. Pays-Bas*, **78**, 417 (1959).
- (309) J. M. Sherfey and A. Brenner, *J. Electrochem. Soc.*, **105**, 665 (1958).
- (310) D. L. Kepert, *Prog. Inorg. Chem.*, **4**, 199 (1962).
- (311) G. Goldstein, C.-M. Wolff, and J.-P. Schwing, *Bull. Soc. Chim. Fr.*, 1201 (1971).
- (312) R. Arnek and Y. Sasaki, *Acta Chem. Scand., Ser. A*, **28**, 20 (1974).
- (313) R. Arnek, *Acta Chem. Scand.*, **23**, 1986 (1969).
- (314) J. C. Dewan and D. L. Kepert, *J. Chem. Soc., Dalton Trans.*, 224 (1973).
- (315) V. A. Nazarenko and E. N. Poluektova, *Russ. J. Inorg. Chem.*, **14**, 105 (1969).
- (316) C. M. Flynn, Jr., M. T. Pope, and S. O'Donnell, *Inorg. Chem.*, **13**, 831 (1974).
- (317) E. S. Ganelina and L. A. Bubnova, *Zh. Neorg. Khim.*, **19**, 159 (1974); *Chem. Abstr.*, **80**, 74814s (1974).
- (318) V. I. Spitsyn, G. V. Kosmodemyanskaya, and M. M. Sadykova, *Zh. Neorg. Khim.*, **17**, 3331 (1972); *Chem. Abstr.*, **78**, 48884d (1973).
- (319) E. S. Ganelina, V. P. Kuzmicheva, L. A. Bubnova, and M. B. Krasnopolskaya, *Zh. Neorg. Khim.*, **18**, 1893 (1973); *Chem. Abstr.*, **79**, 142388n (1973).

- (320) T. J. R. Weakley, *J. Chem. Soc., Dalton Trans.*, 341 (1973).
- (321) E. G. King and W. W. Weller, Bureau of Mines Report of Investigations No. 5791, U.S. Government Printing Office, Washington, D.C., 1961.
- (322) R. L. Graham and L. G. Hepler, *J. Am. Chem. Soc.*, **80**, 3538 (1958).
- (323) T. N. Rezukhina and V. A. Levitskii, *Russ. J. Phys. Chem.*, **37**, 1277 (1963).
- (324) A. Navrotsky and O. J. Kleppa, *Inorg. Chem.*, **8**, 756 (1969).
- (325) V. A. Levitskii and T. N. Rezukhina, *Zh. Fiz. Khim.*, **37**, 1135 (1963); *Chem. Abstr.*, **59**, 7028h (1963).
- (326) T. N. Rezukhina, Y. I. Gerasimov, and V. A. Morozova, *Zh. Fiz. Khim.*, **25**, 93 (1951); *Chem. Abstr.*, **45**, 5005b (1951).
- (327) R. A. Yakovleva and T. N. Rezukhina, *Russ. J. Phys. Chem.*, **34**, 390 (1960).
- (328) Z. V. Proshina and T. N. Rezukhina, *Russ. J. Inorg. Chem.*, **5**, 488 (1960).
- (329) W. G. Lyon and E. F. Westrum, Jr., *J. Chem. Thermodyn.*, **6**, 781 (1974).
- (330) W. G. Lyon and E. F. Westrum, Jr., *J. Chem. Thermodyn.*, **6**, 763 (1974).
- (331) L. A. Zharkova and Y. I. Gerasimov, *Russ. J. Phys. Chem.*, **35**, 1131 (1961).
- (332) A. A. Bulatova, V. B. Aleskovskii, and M. I. Bulatov, *Zh. Prikl. Khim. (Leningrad)*, **46**, 2761 (1973); *Chem. Abstr.*, **80**, 52885p (1974).
- (333) K. Pan, *J. Chin. Chem. Soc. (Taipei)*, **1**, 26 (1954).
- (334) V. N. Chentsov, Y. Y. Skolis, V. A. Levitskii, and Y. Khekimov, *Izv. Akad. Nauk SSSR, Neorg. Mater.*, **9**, 1591 (1973); *Chem. Abstr.*, **80**, 7743e (1974); I. A. Vasileva, S. N. Mudretsova, and E. I. Malienko, *Russ. J. Phys. Chem.*, **47**, 1383 (1973).
- (335) P. A. G. O'Hare and W. N. Hubbard, *J. Phys. Chem.*, **70**, 3353 (1966).
- (336) J. Schroeder and F. J. Sieben, *Chem. Ber.*, **103**, 76 (1970); V. Y. Leonidov, V. S. Pervov, L. I. Klyuev, O. M. Gaisinskaya, V. A. Medvedev, and N. S. Nikolaev, *J. Gen. Chem. USSR (Engl. Transl.)*, **205**, 574 (1972).
- (337) E. J. Barber and G. H. Cady, *J. Phys. Chem.*, **60**, 505 (1956).
- (338) J. Burgess, C. J. W. Fraser, R. D. Peacock, P. Taylor, A. Majid, and J. M. Winfield, *J. Fluorine Chem.*, **3**, 55 (1973/74).
- (339) J. Schroeder and F. J. Grewe, *Chem. Ber.*, **103**, 1536 (1970).
- (340) G. Dittmer, A. Klopfer, D. S. Ross, and J. Schroeder, *J. Chem. Soc., Chem. Commun.*, 846 (1973).
- (341) H. Oppermann, *Z. Anorg. Chem.*, **383**, 1 (1971).
- (342) P. Enghag and L.-I. Staffansson, *Acta Chem. Scand.*, **26**, 1067 (1972).
- (343) W. Brockner, H. Hovdan, and S. J. Cyvin, *Z. Naturforsch., A*, **29**, 620 (1974).
- (344) H. Oppermann, G. Stöver, and G. Kunze, *Z. Anorg. Chem.*, **387**, 317 (1972).
- (345) S. A. Shchukarev and G. A. Kokovin, *Russ. J. Inorg. Chem.*, **9**, 715 (1964).
- (346) S. A. Shchukarev, G. I. Novikov, and G. A. Kokovin, *Zh. Neorg. Khim.*, **4**, 2184 (1959).
- (347) S. A. Shchukarev and G. A. Kokovin, *Russ. J. Inorg. Chem.*, **9**, 849 (1964).
- (348) H. Oppermann and G. Stöver, *Z. Anorg. Chem.*, **383**, 14 (1971).
- (349) S. K. Gupta, *J. Phys. Chem.*, **75**, 112 (1971).
- (350) G. A. Kokovin, *Russ. J. Inorg. Chem. (Engl. Transl.)*, **12**, 7 (1967).
- (351) H. Oppermann, G. Stöver, and G. Kunze, *Z. Anorg. Chem.*, **387**, 329 (1972).
- (352) S. K. Gupta, *Inorg. Chem.*, **12**, 1622 (1973).
- (353) S. K. Gupta, *J. Phys. Chem.*, **73**, 4086 (1969).
- (354) J. Burgess, I. Haigh, and R. D. Peacock, *Chem. Commun.*, 977 (1971).
- (355) N. D. Zaitseva, *Russ. J. Inorg. Chem.*, **10**, 1239 (1963).
- (356) G. N. Kudryashova, L. A. Obzova, and D. V. Korolkov, *Russ. J. Inorg. Chem.*, **15**, 1697 (1970).
- (357) C. Cerny, M. Habes, M. Zelena, and E. Erdős, *Collect. Czech. Chem. Commun.*, **24**, 3836 (1959); *Chem. Abstr.*, **54**, 10465f (1960).
- (358) L. D. McGraw, H. Seltz, and P. E. Snyder, *J. Am. Chem. Soc.*, **69**, 329 (1947).
- (359) H. Baadsgaard and W. D. Treadwell, *Helv. Chim. Acta*, **38**, 1669 (1955).
- (360) C. H. Brubaker, Jr., *J. Am. Chem. Soc.*, **78**, 5762 (1956); K. O. Groves, J. L. Dye, and C. H. Brubaker, Jr., *ibid.*, **82**, 4445 (1960).
- (361) A. A. Opalovskii, V. E. Fedorov, E. U. Lobkov, and B. I. Tsikanovskii, *Russ. J. Phys. Chem.*, **45**, 1064 (1971).
- (362) N. A. Korotchenko, K. K. Fomicheva, G. A. Bogdanov, and R. I. Tereshkina, *Russ. J. Inorg. Chem.*, **18**, 646 (1973).
- (363) M. P. Morozova and G. P. Ovcharova, *Russ. J. Phys. Chem.*, **43**, 966 (1969).
- (364) P. G. Dickens and D. J. Neild, *J. Chem. Soc., Dalton Trans.*, 1074 (1973).
- (365) P. G. Dickens, J. H. Moore, and D. J. Neild, *J. Solid State Chem.*, **7**, 241 (1973); *Chem. Abstr.*, **79**, 140291b (1973); P. G. Dickens and D. J. Neild, *J. Solid State Chem.*, **7**, 474 (1973); *Chem. Abstr.*, **79**, 108788h (1973).
- (366) Y.-W. Kim and G. R. Belton, *Metall. Trans.*, **5**, 1811 (1974).
- (367) N. Sano and G. R. Belton, *Metall. Trans.*, **5**, 2151 (1974).
- (368) A. L. Jones, H. G. Linge, and I. R. Wilson, *Aust. J. Chem.*, **24**, 2005 (1971).
- (369) P. A. G. O'Hare, M. Ader, W. N. Hubbard, G. K. Johnson, and J. L. Settle, *Thermodyn. Nucl. Mater., Proc. Symp.*, **1974**, 2, 439-453 (1974); P. A. G. O'Hare and J. Boerio, submitted for publication. P. A. G. O'Hare, J. Boerio, and K. J. Jensen, submitted for publication.
- (370) L. Denielou, J.-P. Petitot, and C. Tequi, *Thermochim. Acta*, **12**, 209 (1975).
- (371) S. B. Tsyrenova, Yu. L. Suponitskii, and M. Kh. Karapet'yants, *Russ. J. Phys. Chem.*, **48**, 919 (1974).
- (372) I. Dellien, K. G. McCurdy, and L. G. Hepler, *J. Chem. Thermodyn.*, in press.
- (373) R. Gee and R. A. J. Shelton, *J. Less-Common Met.*, **41**, 347 (1975).
- (374) I. A. Ratykovskii, T. A. Pribytkova, and N. V. Galitskii, *High Temp. (Engl. Transl.)*, **12**, 638 (1974).
- (375) M. Aits and H. Schäfer, *Z. Anorg. Allg. Chem.*, **408**, 37 (1974).
- (376) I. Dellien and L. G. Hepler, *Can. J. Chem.*, in press.
- (377) B. Baranowski and K. Bojarski, *Rocz. Chem.*, **46**, 525 (1972); *ibid.*, **46**, 1403 (1972); B. Baranowski and M. Tkacz, *ibid.*, **48**, 713 (1974).
- (378) R. A. Vecher, G. S. Petrov, T. B. Zil'berman, T. A. Karpova, and A. A. Vecher, *Russ. J. Phys. Chem. (Engl. Transl.)*, **48**, 927 (1974).
- (379) E. Rosén, *Chem. Scr.*, **8**, 43 (1975).
- (380) R. Benz, J. F. Elliott, and J. Chipman, *Metall. Trans.*, **5**, 2235 (1974).
- (381) D. A. Pittam, G. Pilcher, D. S. Barnes, H. A. Skinner, and D. Todd, *J. Less-Common Met.*, **42**, 217 (1975).
- (382) G. Pilcher, M. J. Ware, and D. A. Pittam, *J. Less-Common Met.*, **42**, 223 (1975).
- (383) N. A. A. Mumallah and W. J. Popiel, *Anal. Chem.*, **46**, 2055 (1974), and papers cited by these authors; P. Cerutti, E. Shaskin, K. G. McCurdy, and L. G. Hepler, to be submitted for publication.
- (384) P. A. G. O'Hare and H. R. Hoekstra, *J. Chem. Thermodyn.*, **7**, 279 (1975).
- (385) D. R. Fredrickson and M. G. Chasanov, "Analytical Calorimetry", Vol. 3, R. S. Porter and J. F. Johnson, Ed., Plenum Publishing Corp., New York, N.Y., 1974, p 723.
- (386) W. G. Lyon and E. F. Westrum, Jr., *J. Chem. Thermodyn.*, **7**, 741 (1975).
- (387) I. Johnson, *J. Phys. Chem.*, **79**, 722 (1975).
- (388) H. L. Kiwia and E. F. Westrum, Jr., submitted for publication.
- (389) A. Kigoshi, *Thermochim. Acta*, **12**, 179 (1975).
- (390) G. F. McKnight and G. P. Haight, Jr., *Inorg. Chem.*, **12**, 1934 (1973).
- (391) T. G. Chart, *Met. Sci.*, **8**, 344 (1974).
- (392) V. N. Chentsov, V. A. Levitskii, V. K. Trunov, and Ya. I. Gerasimov, *Proc. Acad. Sci. USSR*, **217**, 751 (1974).
- (393) T. N. Rezukhina and T. A. Kashina, *Russ. J. Phys. Chem.*, **48**, 1702 (1974).
- (394) V. A. Levitskii and Y. Y. Skolis, *J. Chem. Thermodyn.*, **6**, 1181 (1974).
- (395) V. A. Levitskii, V. N. Chentsov, Yu. Ya. Skolis, and Yu. G. Golovanova, *Russ. J. Phys. Chem.*, **46**, 151 (1972).
- (396) A. V. Gusarov, V. S. Pervov, I. S. Gotkis, L. I. Klyuev, and V. D. Butskii, *Proc. Acad. Sci. USSR*, **216**, 445 (1974).
- (397) A. J. Kale and K. Sathianandan, *Curr. Sci.*, **39**, 58 (1970).
- (398) M. S. Whittingham, *J. Electrochem. Soc.*, **122**, 713 (1975).
- (399) B. E. Taylor and P. F. Weller, *J. Solid State Chem.*, **1**, 210 (1970).
- (400) D. S. Barnes, G. Pilcher, D. A. Pittam, H. A. Skinner, and D. Todd, *J. Less-Common Met.*, **28**, 53 (1974).